

1 **Oxidative stress in testes of rats exposed to n-butylparaben**

2

3 Elga Schreiber^a, Tània Garcia^a, Raju Prasad Sharma^{a,b}, Margarita Torrente^{a,c}, José Luis
4 Domingo^a, Mercedes Gómez^{a,*}

5 ^a *Laboratory of Toxicology and Environmental Health, School of Medicine, IISPV, Universitat Rovira i
6 Virgili, Reus, Catalonia, Spain*

7 ^b *Environmental Analysis and Management Group, Department of Chemical Engineering, Universitat
8 Rovira i Virgili, Catalonia, Spain*

9 ^c *Research Center in Behavioral Assessment (CRAMC), Department of Psychology, Universitat Rovira i
10 Virgili, Tarragona, Catalonia, Spain*

11

12

13

14

15

16

17

18

19 *Corresponding author: Dra Mercedes Gómez (e-mail: mariamercedes.gomez@urv.cat).
20

21


22

23

24

25

1 Graphical abstract



1 **Highlights**

2

- 3 - Peanut oil causes oxidative stress increasing the levels of TBARS and H₂O₂.
- 4 - High doses of n-ButP show a possible antioxidant effect in testes due to n-ButP
5 and its metabolites.
- 6 - Ca concentration is increased in testes after treatment due to peanut oil.
- 7 - A PBPK model shows that C_{max} in testes is slightly higher than in the plasma.

1 **Abstract**

2 This study was aimed at determining if oxidative stress imbalance in testes of rats
3 occurs after n-butylparaben (n-ButP) exposure. Young male Sprague-Dawley rats were
4 subcutaneously treated with n-ButP during one spermatogenic cycle (57 days) at 0
5 (control-oil), 150, 300 and 600 mg/kg/d with peanut oil as vehicle. A non-vehicle control
6 group was also included. Antioxidant enzyme activities (superoxide dismutase, catalase,
7 glutathione reductase, and glutathione peroxidase) and levels of reduced and oxidized
8 glutathione were measured in testes. Lipid peroxidation and H₂O₂ concentrations were
9 also assessed. Results showed an increase of oxidative stress in oil-treated groups,
10 excepting 600 mg/kg/d, suggesting oxidative stress due to peanut oil. A possible
11 antioxidant effect due to n-ButP and its metabolites was suggested at 600 mg/kg/d, the
12 only group not showing oxidative stress. An increase of calcium concentration in testes
13 was also observed. On the other hand, a physiologically-based pharmacokinetic (PBPK)
14 model was developed and the concentrations of n-ButP and its metabolites were simulated
15 in plasma and testes. The peak concentration (C_{max}) in testes was found slightly higher
16 than that in plasma. The current results indicate that peanut oil can cause oxidative stress
17 while high doses of n-ButP can act as antioxidant agent in testes.

18

19

20 *Keywords:* n-butylparaben (n-ButP), peanut oil, oxidative stress, testes, rats.

21

1 **1. Introduction**

2 Endocrine disruption chemicals (EDCs) can be found in a number of materials
3 such as pesticides, food, drugs or personal care products, etc. They can mimic or interfere
4 the hormone system (Monneret, 2017), having harmful consequences on the biological
5 systems. Nowadays, a wide range of substances have shown to possess some potential to
6 alter hormone homeostasis. These include dioxin and dioxin-like compounds (PCDD/Fs),
7 polychlorinated biphenyls (PCBs), bisphenol A (BPA) or parabens, among others
8 (Giulivo et al., 2016; Kabir et al., 2015; McKinlay et al., 2008; Monneret, 2017; Sweeney
9 et al., 2015; Waring and Harris, 2005).

10 Parabens (PBs) are alkyl esters of the p-hydroxybenzoic acid, which have been
11 widely used as preservative in cosmetics, foods, and pharmaceutical products for more
12 than 50 years (FDA, 2016; Moos et al., 2016). PBs are highly used due to their broad
13 spectrum of antimicrobial activity, low reactivity, high chemical stability (for a wide
14 temperature and pH range), odor and colorless, non-volatility and low costs of production
15 (Błedzka et al., 2014; Silva et al., 2018). In 1974, the FAO/WHO Joint Expert Committee
16 on Food Additives (JECFA), fixed the total acceptable daily intake (ADI) at a maximum
17 of 10 mg/kg body weight (bw) for the sum of methylparaben (MetP), ethylparaben (EthP)
18 and propylparaben (ProP) (JECFA, 1974). MetP and EthP are still currently permitted as
19 food additives, but ProP was recommended to be removed due to its estrogenic effects in
20 young male rats (EFSA, 2004; Sun et al., 2016). In pharmaceutical products, MetP and
21 ProP are often used together due to their synergistic effects (Samarasinghe et al., 2018).
22 In cosmetics, the maximum concentration of 0.4% is recommended for one paraben, and
23 0.8% for a mixture of parabens (Moos et al., 2016). However, the sum of ProP and
24 butylparaben (n-ButP) ester concentrations should not exceed 0.19% in mixtures (Darbre
25 and Harvey, 2014). In 2014, the European Union banned isopropyl-, isobutyl-, benzyl-,

1 pentyl- and phenyl- paraben (Sasseville et al., 2015). Nowadays, the most commonly PBs
2 used are methylparaben (MetP), ethylparaben (EthP), propylparaben (ProP) and
3 butylparaben (n-ButP) (Garcia et al., 2017).

4 *In vitro* studies have shown endocrine toxicity of PBs (Byford et al., 2002; Chen
5 et al., 2007; Kjærstad et al., 2010), while *in vivo* studies have reported adverse effects of
6 PBs such as uterotrophic effects (Lemini et al., 2003), estrogenic activity (Boberg et al.,
7 2016; Garcia et al., 2017; Kang et al., 2002; Zhang et al., 2014), and also on the
8 reproductive tract of male and female rats (Harvey and Everett, 2006). Carcinogenic
9 potential for some parabens has been also reported in human breast (Darbre et al., 2004).

10 A wide variety of EDCs have shown to cause oxidative stress, which targets the
11 endocrine system and can cause reproductive alterations (Sedha et al., 2015). In humans,
12 exposure to PBs through ingestion, inhalation or dermal contact can induce oxidative
13 stress (Kang et al., 2013; Watkins et al., 2015). In rodents (Docea et al., 2018; Shah and
14 Verma, 2011) and in fish models (Ateş et al., 2018; Brown et al., 2018; Silva et al., 2018),
15 it has been found that PBs can induce oxidative stress by altering antioxidants enzymes,
16 the levels of antioxidants molecules, and lipid peroxides.

17 One of the main factors for male infertility is oxidative stress. Reactive oxygen
18 species (ROS) can induce lipid peroxidation, sperm motility loss, DNA fragmentation at
19 spermatozoa level, disrupt steroidogenesis and membrane stabilization in testes (Aitken
20 et al., 2016). *In vivo* studies have demonstrated a link between abnormalities of male
21 reproductive function and the induction of oxidative stress (Aksu et al., 2017; Khan et al.,
22 2015; Rodríguez-González et al., 2015). Testes are very sensitive to oxidative stress due
23 to the abundance of highly unsaturated fatty acids, and the presence of potential ROS-
24 producing systems such as high rates of cells divisions, corresponding to spermatogenesis
25 process, and also Leydig cells steroidogenesis (Yuksel et al., 2012).

1 A possible involvement of trace elements in the dysfunction of male reproductive
2 system in terms of endocrine activity, gametes production and gamete quality, has been
3 reported in men and other animal species (Fallah et al., 2018; Kovacik et al., 2018;
4 Slivkova et al., 2009). Associations of trace metals with oxidative status parameters,
5 linked to sperm motility parameters have been also reported (Djuric et al., 2015; Tvrda et
6 al., 2013). Some trace elements are considered essential for reproduction (Dobrakowski
7 et al., 2018; Türk et al., 2014). However, combined higher levels of these elements in
8 seminal plasma can have adverse effects on sperm motility and oxidative stress
9 production, interfering with physiological processes responsible for successful
10 fertilization (Fallah et al., 2018; Kovacik et al., 2018).

11 On the other hand, physiologically based pharmacokinetics (PBPK) models have
12 been widely used to describe the distribution and to generate time course concentration
13 profiles of chemicals and their metabolites in different organs and tissues of the body
14 (Sharma et al., 2018a, 2018b). In the current study, a PBPK model has been developed
15 and used to determine the time course of n-ButP target tissue (testes) concentrations. The
16 time course concentrations profile has been used to fit the experimentally observed
17 response for the dose-response analysis.

18 Recently, we found reproductive abnormalities in young male rats subcutaneously
19 treated with n-ButP during one spermatogenic cycle (Garcia et al., 2017). The aim of the
20 present study was to determine if the reproductive abnormalities could be due to a
21 possible oxidative stress imbalance after n-ButP exposure in testes, as well as if trace
22 elements could be involved in this process. Moreover, we also used a PBPK model to
23 evaluate the time course concentration profile of n-ButP in testes and to relate it to the
24 antioxidant system.

1 **2. Materials and Methods**

2 *2.1. Chemicals*

3 N-butylparaben (n-ButP) (purity ≥ 99%, Sigma-Aldrich, St. Louis, MO, USA)
4 was dissolved in absolute ethanol and diluted with peanut oil (Sigma-Aldrich, St. Louis,
5 MO, USA) to obtain concentrations of 0 (control-oil), 150, 300 and 600 mg/kg/d. These
6 doses were selected based on data from the scientific literature, as well as the results of
7 previous studies (Garcia et al., 2017; Kim et al., 2015).

8 *2.2. Animals and experimental design*

9 Young male Sprague-Dawley rats (6-week old) (Charles River Laboratories, Sant
10 Germain-L'Arbresle, France) were housed in a room equipped with automatic light cycles
11 (12 h light/dark cycle) and maintained at $22 \pm 2^{\circ}\text{C}$ and 40 - 60% humidity. Food (Panlab
12 rodent chow, Barcelona, Spain) and tap water were offered *ad libitum*. The experimental
13 design was approved by the Ethics Committee of Animal Research, “Rovira i Virgili”
14 University (Tarragona, Spain).

15 Rats were randomly divided into five groups (10 animals per group): control (no
16 vehicle), control-oil (peanut oil), 150, 300 and 600 mg/kg/d of n-ButP. The chemical was
17 subcutaneously administered, in a volume of 0.5 mL, in neck and both sides of the
18 inguinal zone, for 3 alternative days per week, in order to ensure a correct penetration of
19 the vehicle, during one spermatogenic cycle (57 days). Clinical signs (abnormal behavior,
20 presence of wounds, infections and mortality) were daily monitored. At the middle of the
21 treatment period, after subcutaneous injection, urine samples were collected for 24 h and
22 stored at -80°C .

23 Three days after the exposure period, body weights were recorded, and animals

1 were anesthetized by an intraperitoneal injection of 75 mg/kg of ketamine and 0.5 mg/kg
2 of metedomidine. Subsequently, blood samples were obtained from the portal vein into
3 10 mL serum collection tubes, and maintained at room temperature (minimum 30 min).
4 Tubes were centrifuged at 1300g for 10 min at 4°C in order to obtain serum samples.
5 After centrifugation, serum was immediately separated and stored at -80°C for subsequent
6 Oxygen Radical Absorbance Capacity (ORAC) assay. Testes were removed, weighed,
7 placed in liquid nitrogen and stored at -80°C (Garcia et al., 2017).

8 *2.3. Oxidative stress biomarkers analyses*

9 Testes were cut in sagittal plane. Afterwards, 100 mg of each sample were
10 homogenized in 0.2 M cold sodium phosphate buffer at pH 6.25 in a Potter-Elvehjem
11 homogenizer fitted with a Teflon pestle. Homogenates were centrifuged at 105000g for 1
12 h at 4°C. The crude soluble fraction was divided into two parts. The first part was
13 precipitated with cold trichloroacetic acid (TCA) 70% (final concentration TCA 10%) for
14 20 min at 4 °C, being then centrifuged at 2300g for 10 min at 4°C. The supernatant was
15 stored at -80 °C for the subsequent reduced glutathione (GSH) and oxidized glutathione
16 (GSSG) determination. The second part was aliquoted and stored at -80 °C to determine
17 superoxide dismutase (SOD), catalase (CAT), glutathione reductase (GR) and glutathione
18 peroxidase (GPx) activities, hydrogen peroxide (H_2O_2) concentrations and protein
19 content. Total protein concentration was measured by the Bradford's method (1976),
20 using bovine serum albumin (BSA) (Sigma-Aldrich, St. Louis, MO, USA) as standard.

21 *2.3.1. Antioxidant enzyme activities*

22 Superoxide dismutase (SOD) and catalase (CAT) were determined according to
23 previously reported methods (Cohen et al., 1970; Misra and Fridovich, 1972). SOD
24 activity determination is based on autoxidation of epinephrine by superoxide anion.

1 Enzyme activity was expressed as the amount of enzyme required to inhibit the
2 epinephrine's autoxidation by 50%. Results are given as U/mg of protein. For CAT
3 activity, the assay measured for 0.5 min at 240 nm, the extinction of H₂O₂ substrate due
4 to the action of catalase. Results are expressed as μmol/min/mg protein.

5 *2.3.2. Glutathione dependent enzymes and reducing power*

6 Glutathione reductase (GR) and glutathione peroxidase (GPx) activities were
7 determined using a Cobas Mira automatic analyzer (Roche Pharmaceuticals, Basel,
8 Switzerland), which reads at 340 nm the consumption of NADPH following the reduction
9 of oxidized glutathione (GSSG) and t-butyl hydroperoxide (t-BuOOH), respectively. GR
10 activity was measured according to Goldberg and Spooner (1983), while GPx activity
11 was based on the Wheeler's method (1990). The enzyme activity is expressed as mU/mg
12 of protein.

13 Reduced (GSH) and oxidized (GSSG) glutathione concentrations (Sigma-Aldrich,
14 St. Louis, MO, USA) were measured according to the Hissin and Hilf (1976) method,
15 using a LS50 Perkin Elmer fluorimeter (Perkin Elmer, Baconfield, UK) at 350 and 420
16 nm (excitation and emission wavelength, respectively. GSH and GSSG concentrations
17 are given as nmol/mg of protein. Reducing power (GSSG/GSH ratio) was also
18 determined.

19 *2.3.3. Hydrogen peroxide (H₂O₂) concentration*

20 The concentrations of H₂O₂, a biomarker of oxidative stress in tissues, were
21 determined using a commercial kit (Peroxide assay kit, Sigma-Aldrich, St. Louis, MO,
22 USA), based on a colorimetric method, using different concentrations of pure H₂O₂ as
23 standard. Results are given as μM.

24 *2.3.4. Malondialdehyde assay: thiobarbituric acid reactive substances (TBARS)*

1 Malondialdehyde (MDA) concentration, a marker for lipid peroxidation, was also
2 determined. For MDA assay, testes were cut in sagittal plane and 100 mg of each sample
3 were homogenized in a cold solution of 1.15% KCl containing 0.01% butylated
4 hydroxytoluene (Sigma-Aldrich, St. Louis, MO, USA) (Gómez et al., 2004) in a Potter-
5 Elvehjem homogenizer fitted with Teflon pestle. 100 µL of homogenates were incubated
6 in 3 mL of buffer A (constituted for equal volumes of 0.8% thiobarbituric acid and 20%
7 acetic acid at pH 3.5, and supplemented with 200 µL 8.1% SDS and 700 µL H₂O milli-
8 Q) at 95°C for 1 h. After incubation, homogenates were centrifuged at 3225 g for 15 min
9 at 25 °C. The soluble fraction was read in a standard spectrophotometer at 532 nm
10 (Zingarelli et al., 1999). MDA (Sigma-Aldrich, St. Louis, MO, USA) was used as
11 standard being prepared with buffer B (1% H₂SO₄ and 20% absolute ethanol). Results are
12 given as pmol/mg of protein.

13 *2.3.4. Oxygen Radical Absorbance Capacity (ORAC) assay*

14 The ORAC assay is used to determine the total antioxidant capacity in different
15 matrices. The hydrophilic content of peanut oil (+ n-ButP), serum and urine was analysed.
16 Total antioxidant capacity from vehicle (oil) and vehicle + compound (150, 300 and 600
17 mg/kg/d of n-ButP) was determined. Briefly, vehicle, and vehicle with compound, were
18 diluted 1:1 with 80% of methanol and vortexed for 2 min at room temperature. The
19 solution was centrifuged at 5000 g for 10 min at room temperature and the supernatant
20 was collected. The steps were repeated, being the total supernatant measured and diluted
21 1:20 with 0.075 M Na-K phosphate buffer at pH 7.4. Serum samples were diluted 1/500
22 with 0.075 M Na-K phosphate buffer at pH 7.4 and then analysed. Before urine ORAC
23 assay, samples were centrifuged, and the supernatant analysed without dilutions.

24 This assay is based on a fluorimetric reaction that measures the antioxidant
25 capacity to scavenge peroxide radicals induced by dihydrochloride of 2,2 azobis (2-

1 aminopropane) at 485 nm excitation wave and 538 nm emission wave on a fluorescence
2 plate reader (Fluoroskan Ascent, Labsystems, Helsinki, Finland) (Cao et al., 1998).
3 Results are given as a μ mol of TE/mL. TE are equivalents of trolox, which is an analogue
4 of vitamin E, used as standard.

5 *2.4. Protein expression analyses*

6 CAT (MW: 60-65 kDa), GR (MW: 55 kDa) and GPx-4 (MW: 17-22 kDa, the
7 main isoform in testis) expressions were analysed by the Western blot technique. Samples
8 of testes were homogenized in RIPA buffer supplemented with a cocktail of proteases
9 and phosphatases inhibitors. Total protein concentration was determined using DC
10 Protein Assay kit (Bio-Rad, Hercules, CA, USA), which is based on Lowry's method
11 (1951). It was used due to its compatibility with RIPA buffer. BSA was used as standard.

12 Aliquots containing 30 μ g of total protein were mixed in buffer (0.5 M Tris-HCl
13 pH 6.8, 10% glycerol, 2% (wt/vol) SDS, 5% (vol/vol) 2- β -mercaptoethanol, 0.05%
14 bromophenol blue) and denatured by boiling at 95 °C for 10 min. Samples were
15 electrophoresed on SDS-polyacrylamide gel 12.5 (Bio-Rad, Hercules, CA, USA).
16 Proteins were transferred on a PVDF (Bio-Rad, Hercules, CA, USA) membrane using a
17 transblot apparatus (Bio-Rad, Hercules, CA, USA), being subsequently blocked with non-
18 fat milk diluted in TBS-T (50 mM Tris, 1.5% NaCl, 0.05% Tween 20 at pH 7.5) for 1h.
19 Membranes were incubated with primary antibody solution overnight at 4 °C, being
20 subsequently washed in TBS-T buffer, and incubated with a peroxidase-conjugated
21 immunoglobulin G antibody for 1 h 30 min at room temperature. Proteins were visualized
22 using a chemiluminescent substrate (Western ECL Blotting substrates, Bio-Rad,
23 Hercules, CA, USA). Specific labelled proteins were quantified by measuring the density
24 of the detected bands using Image Lab software (Bio-Rad, Hercules, CA, USA). β -actin
25 (MW: 43 kDa) was used as internal standard. Levels of protein expression were expressed

1 as the relative expression respect to β -actin.

2 *2.5. Analysis of trace elements*

3 Samples of testes were weighed in a Teflon microinsert and 65% nitric acid
4 (Suprapur, Merck, Darmstadt, Germany) was added to digest the samples. Teflon
5 microinserts were introduced in Teflon vessels, which were put into a microwave oven
6 Star D (Milestone, Sorisole, Italy) during 25 min at 200 °C and 1000 W, according to the
7 manufacturer's instructions. The digested solution was diluted at 10 mL with H₂O milli-
8 Q. All materials were previously washed with 10% nitric acid in order to avoid potential
9 contaminations (Gómez et al., 2008). The levels of Cd, Co, Cu, Mn, Pb, Se, and Zn were
10 determined by means of a computer-controlled sequential inductively coupled plasma-
11 mass spectrometer (ICP-MS) (PerkinElmer Elan 6000), while the concentrations of Ca,
12 Fe and Mg were determined by computer-controlled sequential inductively coupled
13 plasma optical emission spectrometer (ICP-OES) (PerkinElmer Optima 3200RL), both
14 according to DIN EN ISO 17294-2. Detection limits were the following: Cd, 0.05 ng/g;
15 Co, 0.05 ng/g; Cu, 0.25 ng/g; Mn, 0.05 ng/g; Pb, 0.05 ng/g; Se, 2 ng/g; Zn, 1.5 ng/g; Ca,
16 0.025 µg/g; Fe, 0.005 µg/g; and Mg, 0.025 µg/g. For quality control, DOLT-5 Standard
17 (Fisher liver, National Research Council Canada, Ottawa, ON, Canada) was used in each
18 assay.

19 *2.6. PBPK model*

20 The rat PBPK model developed for this study comprises six compartments: gut,
21 liver, plasma, kidney, fat, and testes, as well as a compartment representing the rest of the
22 body. The exchange of the chemical between blood and tissue in each organ is described
23 by a perfusion rate-limited PBPK model. A previously developed PBPK model
24 (Campbell et al., 2015) was modified to build the current model by adding the

1 subcutaneous and others several compartments such as testes. Subcutaneous dose was
2 considered to build the model, which was evaluated against the plasma concentrations in
3 the rat, after a dose of 100 mg/kg body weight (Aubert et al., 2012). The partition
4 coefficient was estimated using the QSAR approach of Poulin and Krishnan tissue
5 composition method (Poulin and Krishnan, 1996, 1995; Poulin and Theil, 2000). The
6 clearance of butyl paraben in liver, gut and skin was described using the Michaelis-
7 Menten equation. The PBPK model was developed using ordinary differential equations.
8 The equations were written in the GNU MCSim modeling language (Bois and Maszle,
9 1997) and solved by numerical integration with the GNU MCSim, using the R platform.
10 Monte Carlo simulations were performed to estimate the impact on model predictions of
11 uncertainty in all of the rate constants and the partition coefficients. The model output
12 simulation was in a range between 2.5th and 97.5th percentiles.

13 *2.7. Statistics*

14 Statistical analysis was performed using ANOVA, followed by the Tukey post-
15 hoc test when variances among treated groups were homogeneous. If variances were not
16 homogeneous, the Kruskal-Wallis test, followed by the Dunns post-hoc test, was used.
17 To determine differences due to vehicle, two-tailed t-test between control and control-oil
18 (0 mg/kg/d) groups was performed. If differences between control groups were not found,
19 ANOVA was used between treated groups and control-oil group. If control groups were
20 different, both were then included in the ANOVAs analysis. Statistical analysis was
21 performed using the software Statistical Package for the Social Sciences (SPSS v.25).
22 Significance was set at p < 0.05.

23 **3. Results**

24 *3.1. Oxidative stress markers*

1 Data on oxidative stress markers in testes are depicted in Figure 1. SOD activity
2 did not show differences among groups (Figure 1A) while a dose-related tendency to
3 decrease was noticed in CAT activity, being 600 mg/kg/d group the only group showing
4 differences with respect to the control-oil group (Figure 1B). Regarding glutathione
5 dependent enzymes activities, differences were noted. An overall effect on the decrease
6 in GR activity due to oil was observed (Figure 1C), while a significant increase between
7 the control-oil group and the 600 mg/kg/d group was due to n-ButP exposure. According
8 to GPx activity, a significant increase in control-oil group with respect to the control
9 group was noted. Moreover, all n-ButP-treated groups showed a significant decrease with
10 respect to the control-oil group (Figure 1D).

11 Glutathione reducing power was established through GSH and GSSG levels and
12 GSSG/GSH ratio. Only concentrations of GSH showed significant differences (Figure 2).
13 Control-oil and 300 mg/kg/d groups showed a significant decrease in comparison to
14 control group. Although the rest of groups did not show differences with respect to the
15 control group, a tendency to increase their levels in relation to the control-oil group was
16 noted. Moreover, GSH levels at 600 mg/kg/d were significantly different from those in
17 the control-oil group, showing an increase. Figure 3 shows H₂O₂ concentrations in testes.
18 A significant increase in all oil-treated groups, excepting 600 mg/kg/d, was observed with
19 respect to the control group. The results of the malondialdehyde assay showed a
20 significant increase (in comparison to the control group) for the oil-treated groups,
21 excepting at 600 mg/kg/d (Figure 4).

22 As a consequence of the possible oxidizing effects due to peanut oil, the ORAC
23 assay was conducted to determine total antioxidant capacity of the hydrophilic fraction
24 of the peanut oil. Analysis showed some antioxidant capacity in groups treated with n-
25 ButP, while control-oil group did not show antioxidant capacity by itself (Table 1). In

1 relation to serum samples, differences between the control-oil group and the 150 mg/kg/d
2 group were found. Serum samples were collected 3 days after the last subcutaneous
3 injection, being well known that n-ButP is removed after 48 h (Figure 6B). Finally, urine
4 samples in both control and 600 mg/kg/d groups presented significant high μ mol of
5 equivalents of trolox/mL with respect to all oil-treated groups. In addition, the 600
6 mg/kg/d group showed the same levels that the control group.

7 *3.2. Protein expression of antioxidant enzymes*

8 The relative protein expression levels of antioxidant enzymes CAT, GR and GPx-
9 4 in testes were quantified by Western Blot (Figure 5). CAT enzyme is minority in testes
10 and, consequently, showed a great variability between the different groups, being 600
11 mg/kg/d group the only one showing a significant decrease with respect to the control
12 group (Figures 5A and B). GR expression presented a significant decrease at 300 and 600
13 mg/kg/d with respect to the control and control-oil groups (Figures 5A and C). No
14 differences in GPx-4 expression were found (Figures 5A and D).

15 *3.3. Concentrations of trace elements*

16 Trace elements levels in testes are summarized in Table 2. Concentrations of Cd,
17 Co and Pb were not detected. Those of Cu, Mn, Se, Zn, Fe and Mg were detected, but no
18 significant differences among groups were found. Only Ca levels showed significant
19 differences, when a significant increase due to oil was found in oil-treated groups with
20 respect to the control group, excepting at 300 mg/kg/d. This group was also the only n-
21 ButP treated group that showed reduced Ca levels compared to the control-oil group.

22 *3.4. PBPK model*

23 The model prediction for ButP plasma concentration is depicted in Figure 6A with
24 median (red) and their two extremes, corresponding to 2.5 and 97.5 percentiles (blue).

1 The model was evaluated against the experimental *in vivo* data, where the green points
2 represent the experimentally observed mean concentrations and the black bars represents
3 \pm SD. The model predicted very adequately the experimental data with the rate of
4 appearance; i.e. initial observed experimental points are at the 50th percentile of the modelling
5 percentile with slight deviation at the tail end 24 h time course curve. The maximum
6 plasma concentration (C_{max}) was reached at approx. 4 h (t_{max}) and decreased within 24 h
7 to a significant lower level, not showing signs of accumulation. In Figure 6B, the model
8 is simulated for a scenario of the current experimental set-up for the subcutaneous dosing.
9 The simulation of n-ButP concentration in plasma after a subcutaneous dose of 100 mg/kg
10 body weight, at alternative days for 57 days, did not show accumulation (data not shown).
11 The phenomena of no accumulation can be observed in the Figure 6B, which represents
12 the time course of n-ButP plasma concentrations (median) for 72 h for the current
13 experimental dosing scenario.

14 The model was further simulated for 24 h at the doses used in the current study:
15 150, 300 and 600 mg/kg body weight. The results were recorded for plasma and testis
16 compartments being shown in Figures 7 and 8. Figure 7(A, B and C) represents n-ButP
17 plasma concentrations at the different doses. We observed a dose-dependent increase in
18 C_{max} value, with chemical concentrations decreasing within 24 h to a significantly lower
19 level. It indicates a dose-dependent metabolic activity with higher volume of elimination
20 at higher doses, showing no saturation of enzymatic activities (Figure 7C). PBPK
21 predicted peak plasma concentration to dosing ratio of n-ButP is approximately 10% in
22 case of the administered subcutaneous dose (Figure 7). The t_{max} (time at which C_{max}
23 occurs) for concentration in the testes was observed at approx. 5 h of treatment, while the
24 plasma t_{max} occurs well before 5 h, which might be due to the distribution delay. The
25 corresponding C_{max} values in testes were slightly higher than those in the plasma at the

1 given subcutaneous doses, suggesting that testes are exposed to a higher concentration
2 than the plasma (Figure 8). However, the elimination profile in testes followed the similar
3 pattern to that of plasma, being almost eliminated within 24 h (Figure 8).

4 **4. Discussion**

5 In the present study, rats were subcutaneously treated with peanut oil during one
6 spermatogenic cycle. The present results show that TBARS and H₂O₂ levels increased
7 with respect to these in the control group, excepting at 600 mg/kg/d, which indicates
8 oxidative stress in testes. It may be due to the peanut oil effect and not to the
9 administration route. In the current investigation, oil-treated rats started to show little
10 depots under the skin and Levoe et al. (2014) found depots and skin abnormalities after
11 subcutaneous treatment with peanut oil during 21 days in rats. They suggested that an
12 activation of immune system was due to the peanut oil and no to the own injection. It is
13 well known that immune response is highly related to oxidative stress (Marri and Richner,
14 2015) and that subcutaneous route is painless (Nebendahl, 2000). In addition, peanut oil
15 has a high proportion of unsaturated fatty acids: 35-58% of oleic (18:1, n-9) and 20-43%
16 of linoleic (18:2, n-6,9), being the rest saturated fatty acids (Carrín and Carelli, 2010).
17 Unsaturated fatty acids can be oxidized faster and are related with elevated TBARS levels
18 (Samarasinghe et al., 2018; Verma and Asnani, 2007; Yun and Surh, 2012). We found
19 that oxidative stress in testes was mainly caused by peanut oil administration, but the 600
20 mg/kg/d group seemed to act independently on the other oil-treated groups. The high dose
21 of n-ButP showed an antioxidant effect with a tendency to restore the normal TBARS
22 levels. This effect might be due to exposure to n-ButP.

23 P-hydroxybenzoic acid and p-hydroxyhippuric acid are the metabolites of n-ButP
24 hydrolysis. Both are widely known as antioxidants being established that phenolic
25 compounds have the property to stabilize some oils (Gómez-Alonso et al., 2002). It can

1 suggest the possibility that the antioxidant effects due to n-ButP (or its metabolites) are
2 higher than the oxidant effects due to peanut oil at 600 mg/kg/d and, as result, no oxidative
3 stress was found. Popa and co-workers (2014) treated rats orally with BPA + MetP,
4 reporting that p-hydroxybenzoic acid had the capacity to trapping hydroxyl radicals and,
5 in consequence, can inhibit lipid peroxidation in plasma. In turn, Kopalli et al. (2013)
6 reported that MetP could inhibit LPO in mice neuronal cells. These studies would support
7 our suggestion on the role of n-ButP and its metabolites to help to keep a non-oxidative
8 state at 600 mg/kg/d. In contrast, Riad et al. (2018) found that butylparaben can cause
9 reproductive abnormalities and oxidative stress in testis of weanling rats treated orally
10 with a dose of n-ButP lower than 150 mg/kg/d. Some studies have also reported oxidative
11 stress due to n-ButP (Brown et al., 2018; Hegazy et al., 2015; Shah and Verma, 2011;
12 Yang et al., 2018). These apparently contradictory results could be due to the different
13 animal species, the analysed tissues, the duration of the treatment, as well as the route of
14 administration, among other experimental conditions.

15 In relation to ORAC analyses, the equivalents trolox/mL were found at all doses
16 of n-ButP, whereas the vehicle group did not show this antioxidant capacity. This is in
17 accordance with our previous suggestion that n-ButP could have antioxidant properties.
18 However, ORAC in serum did not show differences in antioxidant capacity, which can
19 be due to the fact that n-ButP is almost removed after 48 h, just like PBPK model shows
20 (Aubert et al., 2012). Regarding urine samples, all groups with an increase of ROS had
21 less antioxidant capacity with respect to the control group. These results would support
22 our hypothesis, which suggests a little antioxidant capacity due to n-ButP and their
23 metabolites, being this effect only effective at the highest dose of the chemical.

24 On the other hand, SOD, CAT and GPx are the first defence line against oxygen
25 radicals. In the present study, SOD did not show activity differences. Animals were

1 sacrificed 3 days after the last exposure, being n-ButP completely metabolized and
2 eliminated after 48 h. Therefore, we suggested that SOD activity could be restored after
3 the treatment. By contrast, CAT and GPx levels were altered, suggesting that their
4 concentrations were not restored. CAT and GPx are implied in H₂O₂ degradation, while
5 GPx also acts removing all kind of peroxides. Increasing levels of TBARS and H₂O₂ can
6 generate an activation of CAT and GPx, being GPx predominant in testes (Bauché et al.,
7 1994). The current results show an increase of GPx activity in the control-oil group, which
8 suggests a reactivation of GPx activity as a consequence of oxidative stress. However, all
9 n-ButP treated groups showed a significant decrease in this activity with respect to the
10 control-oil group. We suggest on scenario in which GPx activity of control-oil group
11 increases due to H₂O₂ levels, while possible n-ButP antioxidant effects would help to
12 restore normal activity values. Only the highest n-ButP dose showed a decrease of CAT
13 activity in comparison to that in the control-oil group. At the highest dose, no differences
14 in TBARS and H₂O₂ levels were found, which suggests that the lower activity of CAT
15 respecting that in the control-oil group, might be due to the antioxidant n-ButP effects.

16 With respect to GR, it is an enzyme implied in a second line of antioxidant
17 defence. All n-ButP treated groups showed a decrease in GR activity (regarding the
18 control group) related with an oxidative state. GR depletion was observed in all groups
19 with elevated levels of TBARS and H₂O₂. Moreover, a decrease in GR activity was also
20 observed at 600 mg/kg/d. However, this group showed a significant increase in GR
21 activity with respect to the rest of oil-treated groups. This increase could be related with
22 a previous oxidative stress, which is possibly being restored. Shah and Verma (2011)
23 reported a decrease in antioxidant enzymes activities, depletion of GSH, and an increase
24 of TBARS levels due to exposure to n-ButP in mice liver respecting a control-oil group.
25 In the current investigation, similar results due to peanut oil were found in testes.

1 According to glutathione concentrations, a depletion in GSH levels were found in
2 the control-oil and 300 mg/kg/d groups, which indicates oxidative stress in testes. In
3 contrast, the 600 mg/kg/d group showed an increase in GSH concentrations, with a
4 tendency to restore normal levels. These results are in accordance with our hypothesis
5 that peanut oil can generate ROS. In contrast, Nakagawa and Moldeus (1998) reported
6 GSH depletion due to n-ButP in isolated hepatocytes. These authors suggested that a
7 suppression of GR could be related to GSH depletion, and the significant decreased
8 enzyme activities might be due to protein oxidation induced by n-ButP.

9 Protein expression was analysed using western blot technique. It is important to
10 remark that GPx-4 is the major isoform in testes (Liang et al., 2009). GPx-4 expression
11 did not show differences among n-ButP treated groups with respect to control-oil group.
12 It could mean that there is only an enhancement in control-oil activity due to oxidative
13 stress, but this effect was not enough to affect GPx-4 expression. With respect to CAT
14 and GR expressions, all results would suggest that ROS levels due to peanut oil can affect
15 protein activity, but were not sufficient to damage protein expression. Probably, the lower
16 expression of CAT and GR at the highest doses was due to an elevated levels of n-ButP
17 in testes for 3 alternative days per week during 57 days (Figure 8C), which would affect
18 other metabolic pathways as response to its endocrine disruptor potential. It is known that
19 in some cases, the activity and expression of a same protein does not change in the same
20 way. Shah and Verma (2011) suggested that protein expression reduction might be due
21 to n-ButP.

22 Trace elements have an important role in cell homeostasis. In this study, Cu, Mn,
23 Se, Zn Fe and Mg concentrations did not show significant differences between groups.
24 Only Ca showed a significant increase with respect to the control group, with the
25 exception of the 300 mg/kg/d group. Yang et al. (2018) treated trophoblast cells with

1 parabens reporting that ROS induced a cytoplasmatic Ca increase while Popa et al. (2014)
2 also reported disruption of cytosolic Ca homeostasis in liver and kidney due to MetP. In
3 contrast, in the present study, peanut oil induced ROS, and consequently, Ca in testes
4 could increase in the oil-treated groups with respect to the control group.

5 On the other hand, delivery, distribution and residence time of n-ButP was
6 simulated with a PBPK model for all the three dose scenarios. The C_{max} was slightly
7 higher in testes than in plasma at the corresponding doses, suggesting a higher exposure
8 in testes of n-ButP than in the circulatory fluid of the rat. However, the elimination profile
9 of n-ButP in both plasma and testes was similar with no accumulation. Time course
10 concentration profile in the testes was directly proportional to the animal dose, validating
11 the delivery, distribution and elimination of n-ButP, as also reported in a recent previous
12 (Aubert et al., 2012).

13 **5. Conclusion**

14 To the best of our knowledge, this is the first study evaluating alterations in
15 oxidant-antioxidant status, as well as the concentrations of trace elements in rat testes
16 subcutaneously treated with n-ButP. Rats subcutaneously treated with peanut oil + n-ButP
17 can develop oxidative stress due to oil, but not to n-ButP. These results suggest that either
18 peanut oil can mask the effects of n-ButP, or the chemical can have not sufficient
19 antioxidant effects. High doses of n-ButP are metabolized in high quantities of p-
20 hydroxybenzoic acid and p-hydroxyhippuric acid. These metabolites can then act as
21 antioxidant compounds. In addition, they can stabilize peanut oil, reducing its oxidant
22 effects.

23 **Conflict of interest**

24 The authors declare that there are no conflicts of interest.

1 **Acknowledgments**

2 This work was supported by grant from European Commission: European Test and Risk
3 Assessment Strategies for Mixtures (EuroMix) No. 633172 (H2020-SFS-2014-2).

4

5

6

7

8

9 **REFERENCES**

10

11 Aitken, R.J., Gibb, Z., Baker, M.A., Drevet, J., Gharagozloo, P., 2016. Causes and
12 consequences of oxidative stress in spermatozoa. *Reprod. Fertil. Dev.* 28, 1–10.

13 Aksu, E.H., Kandemir, F.M., Özkaraca, M., Ömür, A.D., Küçükler, S., Çomaklı, S., 2017.

14 Rutin ameliorates cisplatin-induced reproductive damage via suppression of oxidative
15 stress and apoptosis in adult male rats. *Andrologia* 49, 1–8.

16 Ateş, P.S., Ünal, İ., Üstündağ, Ü.V., Alturfan, A.A., Yiğitbaşı, T., Emekli-Alturfan, E.,
17 2018. Methylparaben induces malformations and alterations on apoptosis, oxidant–
18 antioxidant status, ccnd1 and myca expressions in zebrafish embryos. *J. Biochem. Mol.*
19 Toxicol.

20 Aubert, N., Ameller, T., Legrand, J.J., 2012. Systemic exposure to parabens:
21 Pharmacokinetics, tissue distribution, excretion balance and plasma metabolites of [¹⁴C]-
22 methyl-, propyl- and butylparaben in rats after oral, topical or subcutaneous

- 1 administration. Food Chem. Toxicol. 50, 445–454.
- 2 Bauché, F., Fouchard, M.H., Jégou, B., 1994. Antioxidant system in rat testicular cells.
- 3 FEBS Lett. 349, 392–396.
- 4 Błedzka, D., Gromadzińska, J., Wasowicz, W., 2014. Parabens. From environmental
- 5 studies to human health. Environ. Int. 67, 27–42.
- 6 Boberg, J., Axelstad, M., Svingen, T., Mandrup, K., Christiansen, S., Vinggaard, A.M.,
- 7 Hass, U., 2016. Multiple endocrine disrupting effects in rats perinatally exposed to
- 8 butylparaben. Toxicol. Sci. 152, 244–255.
- 9 Bois, F.Y., Maszle, D.R., 1997. MCSim : A Monte Carlo Simulation Program. J. Stat.
- 10 Softw. 2.
- 11 Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram
- 12 quantities of protein utilizing the principle of protein-dye binding. Anal. Biochem. 72,
- 13 248–254.
- 14 Brown, S.E., Sant, K.E., Fleischman, S.M., Venezia, O., Roy, M.A., Zhao, L., Timme-
- 15 Laragy, A.R., 2018. Pancreatic beta cells are a sensitive target of embryonic exposure to
- 16 butylparaben in zebrafish (*Danio rerio*). Birth Defects Res. 110, 933–948.
- 17 Byford, J.R., Shaw, L.E., Drew, G., Sauer, M.J., Darbre, P.D., 2002. Oestrogenic activity
- 18 of parabens in MCF7 human breast cancer cells. J. Steroid. Biochem. Mol. Biol. 80, 49–
- 19 60.
- 20 Campbell, J.L., Yoon, M., Clewell, H.J., 2015. A case study on quantitative in vitro to in
- 21 vivo extrapolation for environmental esters: Methyl-, propyl- and butylparaben.
- 22 Toxicology 332, 67–76.
- 23 Cao, G., Booth, S.L., Sadowski, J.A., Prior, R.L., 1998. Increases in human plasma

- 1 antioxidant capacity after consumption of controlled diets high in fruit and vegetables.
- 2 Am. J. Clin. Nutr. 68, 1081–1087.
- 3 Carrín, M.E., Carelli, A.A., 2010. Peanut oil: Compositional data. Eur. J. Lipid Sci.
- 4 Technol. 112, 697–707.
- 5 Chen, J., Ahn, K.C., Gee, N.A., Gee, S.J., Hammock, B.D., Lasley, B.L., 2007.
- 6 Antiandrogenic properties of parabens and other phenolic containing small molecules in
- 7 personal care products. Toxicol. Appl. Pharmacol. 221, 278–284.
- 8 Cohen, G., Dembiec, D., Marcus, J., 1970. Measurement of catalase activity in tissue
- 9 extracts. Anal. Biochem. 34, 30–38.
- 10 Darbre, P.D., Aljarrah, A., Miller, W.R., Coldham, N.G., Sauer, M.J., Pope, G.S., 2004.
- 11 Concentrations of parabens in human breast tumours. J. Appl. Toxicol. 24, 5–13.
- 12 Darbre, P.D., Harvey, P.W., 2014. Parabens can enable hallmarks and characteristics of
- 13 cancer in human breast epithelial cells: A review of the literature with reference to new
- 14 exposure data and regulatory status. J. Appl. Toxicol. 34, 925–938.
- 15 Djuric, A., Begic, A., Gobeljic, B., Stanojevic, I., Ninkovic, M., Vojvodic, D., Pantelic,
- 16 A., Zebic, G., Prokic, V., Dejanovic, B., Stojanovic, I., Pavlica, M., Djukic, D., Saso,
- 17 L., Djurdjevic, D., Pavlovic, M., Topic, A., Vujanovic, D., Stevnovic, I., Djukic, M.,
- 18 2015. Oxidative stress, bioelements and androgen status in testes of rats subacutely
- 19 exposed to cadmium. Food Chem. Toxicol. 86, 25–33.
- 20 Dobrakowski, M., Kaletka, Z., Machoń-Grecka, A., Kasperczyk, S., Horak, S., Birkner,
- 21 E., Zalejska-Fiolka, J., Kasperczyk, A., 2018. The role of oxidative stress, selected
- 22 metals, and parameters of the immune system in male fertility. Oxid. Med. Cell. Longev.
- 23 2018, 1–8.

- 1 Docea, A.O., Gofita, E., Goumenou, M., Calina, D., Rogoveanu, O., Varut, M., Olaru,
2 C., Kerasioti, E., Fountoucidou, P., Taitzoglou, I., Zlatian, O., Rakitskii,
3 V.N., Hernandez, A.F., Kouretas, D., Tsatsakis, A., 2018. Six months exposure to
4 a real life mixture of 13 chemicals' below individual NOAELs induced non monotonic
5 sex-dependent biochemical and redox status changes in rats. Food. Chem. Toxicol. 115,
6 470–481.
- 7 European Food Safety Authority (EFSA), 2004. Opinion of the Scientific Panel on food
8 additives, flavourings, processing aids and materials in contact with food (AFC) related
9 to para hydroxybenzoates (E 214-219). EFSA J. 83, 1–26.
- 10 Fallah, A., Mohammad-Hasani, A., Colagar, A.H., 2018. Zinc is an essential element for
11 male fertility: A review of Zn roles in men's health, germination, sperm quality, and
12 fertilization. J. Reprod. Infertil. 19, 69–81.
- 13 Food and drug administration (FDA), 2016. Parabens in Cosmetics.
- 14 Garcia, T., Schreiber, E., Kumar, V., Sharma, R.P., Sirvent, J.J., Domingo, J.L., Gómez,
15 M., 2017. Effects on the reproductive system of young male rats of subcutaneous
16 exposure to n-butylparaben. Food Chem. Toxicol. 106, 47–57.
- 17 Giulivo, M., Lopez de Alda, M., Capri, E., Barcel, D., 2016. Human exposure to
18 endocrine disrupting compounds: their role in reproductive systems, metabolic syndrome
19 and breast cancer. A. Rev. Environ. Res. 151, 251–264.
- 20 Goldberg, D.M., Spooner, R.J., 1983. Assay of glutathione reductase, in: Bergmeyen,
21 H.V. (Ed.), Methods of Enzymatic Analysis. Verlag Chemie, Weinheim, pp. 258–265.
- 22 Gómez, J.A., Molero, X., Vaquero, E., Alonso, A., Salas, A., Malagelada, J.R., 2004.
23 Vitamin E attenuates biochemical and morphological features associated with

- 1 development of chronic pancreatitis. Am. J. Physiol. Gastrointest. Liver Physiol. 287,
2 G162–G169.
- 3 Gómez, M., Esparza, J.L., Cabré, M., García, T., Domingo, J.L., 2008. Aluminum
4 exposure through the diet: metal levels in A β PP transgenic mice, a model for Alzheimer's
5 disease. Toxicology 249, 214–219.
- 6 Gómez-Alonso, S., Salvador, M., Fragapan, G., 2002. Phenolic compound profile of
7 cornicabra virgin olive oil. J. Agric. Food Chem. 50, 6812–6817.
- 8 Harvey, P.W., Everett, D.J., 2006. Regulation of endocrine-disrupting chemicals: critical
9 overview and deficiencies in toxicology and risk assessment for human health. Best.
10 Pract. Res. Clin. Endocrinol. Metab. 20, 145–165.
- 11 Hegazy, H.G., Ali, E.H.A., Elgoly, A.H.M., 2015. Interplay between pro-inflammatory
12 cytokines and brain oxidative stress biomarkers: Evidence of parallels between butyl
13 paraben intoxication and the valproic acid brain physiopathology in autism rat model.
14 Cytokine 71, 173–180.
- 15 Hissin, P.J., Hilf, R., 1976. A fluorometric method for determination of oxidized and
16 reduced glutathione in tissues. Anal. Biochem. 74, 214–226.
- 17 Joint Expert Committee on Food Additives (JECFA), 1974. Evaluación Toxicológica de
18 ciertos aditivos alimentarios con un examen de los principios generales y de las normas.
- 19 Kabir, E.R., Rahman, M.S., Rahman, I., 2015. A review on endocrine disruptors and their
20 possible impacts on human health. Environ. Toxicol. Pharmacol. 40, 241–258.
- 21 Kang, K.S., Che, J.H., Ryu, D.Y., Kim, T.W., Li, G.X., Lee, Y.S., 2002. Decreased sperm
22 number and motile activity on the F1 offspring maternally exposed to butyl p-
23 hydroxybenzoic acid (butyl paraben). J. Vet. Med. Sci. 64, 227–235.

- 1 Kang, S., Kim, S., Park, J., Kim, H.J., Lee, J., Choi, G., Choi, S., Kim, S.Y., Moon, H.B.,
2 Kho, Y.L., Choi, K., 2013. Urinary paraben concentrations among pregnant women and
3 their matching newborn infants of Korea, and the association with oxidative stress
4 biomarkers. *Sci. Total Environ.* 461, 214–221.
- 5 Khan, S., Jan, M.H., Kumar, D., Telang, A.G., 2015. Firpronil induced spermotoxicity is
6 associated with oxidative stress, DNA damage and apoptosis in male rats. *Pestic.*
7 *Biochem. Physiol.* 124, 8–14.
- 8 Kim, J.K., Kwack, S.J., Lim, S.K., Kim, Y.J., Roh, T.H., Choi, S.M., Kim, H.S., Lee,
9 B.M., 2015. Toxicological evaluation of isopropylparaben and isobutylparaben mixture
10 in Sprague-Dawley rats following 28 days of dermal exposure. *Regul. Toxicol.*
11 *Pharmacol.* 73, 544–551.
- 12 Kjærstad, M.B., Taxvig, C., Andersen, H.R., Nellemann, C., 2010. Mixture effects of
13 endocrine disrupting compounds in vitro. *Int. J. Androl.* 33, 425–433.
- 14 Kopalli, S.R., Noh, S.J., Koppula, S., Suh, Y.H., 2013. Methylparaben protects 6-
15 hydroxydopamine-induced neurotoxicity in SH-SY5Y cells and improved behavioral
16 impairments in mouse model of Parkinson's disease. *Neurotoxicology* 34, 25–32.
- 17 Kovacik, A., Tirpaka, F., Tomkab, M., Miskejec, M., Tvrdaa, E., Arvayd, J., Andrejie, J.,
18 Slaninaa, T., Gaborf, M., Hlebag, L., Fike, T., Jambora, M., Cisarovah, M., Massanyia,
19 P., 2018. Trace elements content in semen and their interactions with sperm quality and
20 RedOx status in freshwater fish *Cyprinus carpio*: A correlation study. *J. Trace Elem. Med.*
21 *Biol.* 50, 399–407.
- 22 Lemini, C., Jaimez, R., Avila, M.E., Franco, Y., Larrea, F., Lemus, A.E., 2003. In vivo
23 and in vitro estrogen bioactivities of alkyl parabens. *Toxicol. Ind. Health* 19, 69–79.

- 1 Levoe, S.N., Flannery, B.M., Brignolo, L., Imai, D.M., Koehne, A., Austin, A.T., Bruun,
2 D.A., Tancredi, D.J., Lein, P.J., 2014. Factors influencing adverse skin responses in rats
3 receiving repeated subcutaneous injections and potential impact on neurobehavior. *Curr.*
4 *Neurobiol.* 5, 1–10.
- 5 Liang, H., Yoo, S.E., Na, R., Walter, C.A., Richardson, A., Ran, Q., 2009. Short form
6 glutathione peroxidase 4 is the essential isoform required for survival and somatic
7 mitochondrial functions. *J. Biol. Chem.* 284, 30836–30844.
- 8 Lowry, O.H., Rosebrough, N.J., Farr, A.L., Randall, R.J., 1951. Protein measurement
9 with the folin phenol reagent. *J. Biol. Chem.* 193, 265–275.
- 10 Marri, V., Richner, H., 2015. Immune response, oxidative stress and dietary antioxidants
11 in great tit nestlings. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 179, 192–196.
- 12 McKinlay, R., Plant, J.A., Bell, J.N.B., Voulvoulis, N., 2008. Endocrine disrupting
13 pesticides: implications for risk assessment. *Environ. Int.* 34, 168–183.
- 14 Misra, H.P., Fridovich, I., 1972. The role of superoxide anion in the autoxidation of
15 epinephrine and a simple assay for superoxide dismutase. *J. Biol. Chem.* 247, 3170–3175.
- 16 Monneret, C., 2017. What is an endocrine disruptor? *Comptes Rendus - Biol.* 340, 403–
17 405.
- 18 Moos, R.K., Angerer, J., Dierkes, G., Brüning, T., Koch, H.M., 2016. Metabolism and
19 elimination of methyl, iso- and n-butyl paraben in human urine after single oral dosage.
20 *Arch. Toxicol.* 90, 2699–2709.
- 21 Nakagawa, Y., Moldeus, P., 1998. Mechanism of p-hydroxybenzoate ester-induced
22 mitochondrial dysfunction and cytotoxicity in isolated rat hepatocytes. *Biochem.*
23 *Pharmacol.* 55, 1907–1914.

- 1 Nebendahl, K., 2000. Routes of administration, in: Krinke, G.J. (Ed.), The laboratory
2 rat. Academic Press, San Diego, pp. 463–483.
- 3 Popa, D.S., Bolfa, P., Kiss, B., Vlase, L., Păltinean, R., Pop, A., Cătoi, C., Crișan, G.,
4 Loghin, F., 2014. Influence of *Genista tinctoria* L or methylparaben on subchronic
5 toxicity of bisphenol A in rats. Biomed. Env. Sci. 27, 85–96.
- 6 Poulin, P., Krishnan, K., 1996. Molecular structure-based prediction of the partition
7 coefficients of organic chemicals for physiological pharmacokinetic models. Toxicol.
8 Mech. Methods 6, 117–137.
- 9 Poulin, P., Krishnan, K., 1995. A biologically-based algorithm for predicting human
10 tissue: blood partition coefficients of organic chemicals. Hum. Exp. Toxicol. 14, 273–
11 280.
- 12 Poulin, P., Theil, F.P., 2000. A priori prediction of tissue: Plasma partition coefficients of
13 drugs to facilitate the use of physiologically-based pharmacokinetic models in drug
14 discovery. J. Pharm. Sci. 89, 16–35.
- 15 Riad, M.A., Abd-Rabo, M.M., Abd El Aziz, S.A., El Behairy, A.M., Badawy, M.M.,
16 2018. Reproductive toxic impact of subchronic treatment with combined butylparaben
17 and triclosan in weanling male rats. J. Biochem. Mol. Toxicol. 32, 1–8.
- 18 Rodríguez-González, G.L., Vega, C.C., Boeck, L., Vázquez, M., Bautista, C.J., Reyes-
19 Castro, L.A., Saldaña, O., Lovera, D., Nathanielsz, P.W., Zambrano, E., 2015. Maternal
20 obesity and overnutrition increase oxidative stress in male rat offspring reproductive
21 system and decrease fertility. Int. J. Obes. (Lond.) 39, 549–556.
- 22 Samarasinghe, S.V.A.C., Krishnan, K., Naidu, R., Megharaj, M., Miller, K., Fraser, B.,
23 Aitken, R.J., 2018. Parabens generate reactive oxygen species in human spermatozoa.

- 1 Andrology 6, 532–541.
- 2 Sasseville, D., Alfalah, M., Lacroix, J.P., 2015. “Parabenoia” Debunked, or “Who’s
3 Afraid of Parabens?” Dermatitis 26, 254–259.
- 4 Sedha, S., Kumar, S., Shukla, S., 2015. Role of oxidative stress in male reproductive
5 dysfunctions with reference to phthalate compounds. Urol. J. 12, 2304–2316.
- 6 Shah, K.H., Verma, R.J., 2011. Butyl p-hydroxybenzoic acid induces oxidative stress in
7 mice liver-an in vivo study. Acta Pol. Pharm. 68, 875–879.
- 8 Sharma, R.P., Schuhmacher, M., Kumar, V., 2018a. Development of a human
9 physiologically based pharmacokinetic (PBPK) model for phthalate (DEHP) and its
10 metabolites : A bottom up modeling approach. Toxicol. Lett. 296, 152–162.
- 11 Sharma, R.P., Schuhmacher, M., Kumar, V., 2018b. The development of a pregnancy
12 PBPK Model for Bisphenol A and its evaluation with the available biomonitoring data.
13 Sci. Total Environ. 624, 55–68.
- 14 Silva, D.C., Serrano, L., Oliveira, T.M.A., Mansano, A.S., Almeida, E.A., Vieira, E.M.,
15 2018. Effects of parabens on antioxidant system and oxidative damages in Nile tilapia
16 (*Oreochromis niloticus*). Ecotoxicol. Environ. Saf. 162, 85–91.
- 17 Slivkova, J., Popelkova, M., Massanyi, P., Toporcerova, S., Stawarz, R., Formicki, G.,
18 Lukac, N., Putała, A., Guzik, M., 2009. Concentration of trace elements in human semen
19 and relation to spermatozoa quality. J. Environ. Sci. Health Part A 44, 370–375.
- 20 Sun, L., Yu, T., Guo, J., Zhang, Z., Hu, Y., Xiao, X., Sun, Y., Xiao, H., Li, J., Zhu, D.,
21 Sai, L., Li, J., 2016. The estrogenicity of methylparaben and ethylparaben at doses close
22 to the acceptable daily intake in immature Sprague-Dawley rats. Sci. Rep. 6, 1–9.

- 1 Sweeney, M.F., Hasan, N., Soto, A.M., 2015. Environmental endocrine disruptors: effects
2 on the human male reproductive system. *Rev. Endocr. Metab. Disord.* 16, 341–357.
- 3 Türk, S., Mändar, R., Mahlapuu, R., Viitak, A., Punab, M., Kullisaar, T., 2014. Male
4 infertility: decreased levels of selenium, zinc and antioxidants. *J. Trace Elem. Med. Biol.*
5 28, 179–185.
- 6 Tvrďá, E., Kňažická, Z., Lukáčová, J., Schneidgenová, M., Goc, Z., Greń, A., Szabó, C.,
7 Massányi, P., Lukáč, N., 2013. The impact of lead and cadmium on selected motility,
8 prooxidant and antioxidant parameters of bovine seminal plasma and spermatozoa. *J.*
9 *Environ. Sci. Health Part A* 48, 1292–1300.
- 10 Verma, R.J., Asnani, V., 2007. Ginger extract ameliorates paraben induced biochemical
11 changes in liver and kidney of mice. *Acta Pol. Pharm. - Drug Res.* 64, 217–220.
- 12 Waring, R.H., Harris, R.M., 2005. Endocrine disrupters: a human risk? *Mol. Cell.*
13 *Endocrinol.* 244, 2–9.
- 14 Watkins, D.J., Ferguson, K.K., Anzalota Del Toro, L.V., Alshawabkeh, A.N., Cordero,
15 J.F., Meeker, J.D., 2015. Associations between urinary phenol and paraben
16 concentrations and markers of oxidative stress and inflammation among pregnant women
17 in Puerto Rico. *Int. J. Hyg. Environ. Health* 218, 212–219.
- 18 Wheeler, C.R., Salzman, J.A., Elsayed, N.M., Omaye, S.T., Korte, D.W., 1990.
19 Automated assays for superoxide dismutase, catalase, glutathione peroxidase, and
20 glutathione reductase activity. *Anal. Biochem.* 184, 193–199.
- 21 Yang, C., Lim, W., Bazer, F.W., Song, G., 2018. Butyl paraben promotes apoptosis in
22 human trophoblast cells through increased oxidative stress-induced endoplasmic
23 reticulum stress. *Environ. Toxicol.* 33, 436–445.

- 1 Yuksel, M.B., Kavak, S., Gecit, I., Basel, H., Gümrükçüoğlu, H.A., Demir, H., Meral, I.,
- 2 2012. Short-term levosimendan treatment protects rat testes against oxidative stress.
- 3 Brazilian J. Med. Biol. Res. 45, 716–720.
- 4 Yun, J.M., Surh, J., 2012. Fatty acid composition as a predictor for the oxidation stability
- 5 of korean vegetable oils with or without induced oxidative stress. Prev. Nutr. Food Sci.
- 6 17, 158–165.
- 7 Zhang, L., Dong, L., Ding, S., Qiao, P., Wang, C., Zhang, M., Zhang, L., Du, Q., Li, Y.,
- 8 Tang, N., Chang, B., 2014. Effects of n-butylparaben on steroidogenesis and
- 9 spermatogenesis through changed E₂ levels in male rat offspring. Environ. Toxicol.
- 10 Pharmacol. 37, 705–717.
- 11 Zingarelli, B., Szabó, C., Salzman, A.L., 1999. Reduced oxidative and nitrosative damage
- 12 in murine experimental colitis in the absence of inducible nitric oxide synthase. Gut 45,
- 13 199–209.

TABLE 1

ORAC assay (μmol of equivalents of trolox/mL) into three different matrices.

	Control	Control-oil	150 mg/kg/d	300 mg/kg/d	600 mg/kg/d
Hydrophilic fraction ¹	---	ND	D (4.00)	D (3.85)	D (4.49)
Serum (<i>n</i> =7)	14.00 \pm 3.70	19.11 \pm 7.56 ^a	10.56 \pm 1.23 ^b	18.70 \pm 7.18 ^{ab}	14.35 \pm 4.18 ^{ab}
Urine (<i>n</i> =5)	0.170 \pm 0.003	0.160 \pm 0.003 ^{**,a}	0.157 \pm 0.003 ^{***,a}	0.158 \pm 0.003 ^{***,a}	0.168 \pm 0.005 ^b

ND: non-detected; D: detected. Hydrophilic extraction values are qualitative.

¹ Hydrophilic fraction of vehicle (peanut oil) and vehicle with three different doses of n-ButP extracted with 80% methanol.

Values are expressed as mean \pm SD.

* Significantly different from the control (** p < 0.01; *** p < 0.005)

^{ab} Different superscripts in the same row indicate significant differences between groups at p < 0.05

TABLE 2

Levels of trace elements in testes of male rats treated with n-ButP (n=8)

		Control	Control-oil	150 mg/kg/d	300 mg/kg/d	600 mg/kg/d
Cd	ng/g	ND	ND	ND	ND	ND
Co	ng/g	ND	ND	ND	ND	ND
Cu	ng/g	2654 ± 656	2389 ± 249	2463 ± 497	2520 ± 497	2388 ± 395
Mn	ng/g	409.0 ± 95.1	368.4 ± 47.4	409.0 ± 98.3	374.4 ± 71.8	361.3 ± 69.3
Pb	ng/g	ND	ND	ND	ND	ND
Se	ng/g	1513 ± 164	1571 ± 228	1422 ± 138	1725 ± 244	1475 ± 255
Zn	ng/g	27276 ± 12492	35655 ± 6891	32063 ± 8925	34878 ± 13267	31602 ± 6931
Ca	µg/g	22.28 ± 9.77	61.13 ± 20.00 ^{***,a}	50.8 ± 14.94 ^{**,ab}	35.1 ± 7.93 ^b	51.67 ± 14.53 ^{**,ab}
Fe	µg/g	26.80 ± 6.75	25.14 ± 3.20	25.34 ± 1.87	27.52 ± 4.32	24.04 ± 3.45
Mg	µg/g	190.8 ± 49.5	179.8 ± 17.5	192.1 ± 50.5	183.0 ± 36.6	161.7 ± 19.4

ND: non-detected

Values are expressed as mean ± SD

* Significantly different from the control (*p < 0.05; **p < 0.01; ***p < 0.005)


^{a,b} Different superscripts in the same row indicate significant differences between groups at p < 0.05

1

2

1 **FIGURES**

2




3

4 Figure 1. Antioxidant enzyme activities. A: Superoxid dismutase (SOD) activity (U/mg
5 protein); B: Catalase (CAT) activity ($\mu\text{mol}/\text{min}/\text{mg}$ protein); C: Glutathione Reductase
6 (GR) activity (mU/mg protein); D: Glutathione Peroxidase (GPx) activity (mU/mg
7 protein). All values are expressed as mean \pm SD. Asterisk (*) means significantly different
8 from the control group (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.005$). Different superscripts (ab)
9 indicate significant differences between groups at $p < 0.05$.

10

11

12




1

2 Figure 2: Glutathione levels (nmol/mg protein). GSH: reduced glutathione; GSSG:
 3 oxidized glutathione. GSSG/GSH: ratio between oxidized/reduced glutathione (no units).


4 All values are expressed as mean \pm SD. Asterisk (*) means significantly different from
 5 the control group (${}^*p < 0.05$; ${}^{**}p < 0.01$; ${}^{***}p < 0.005$). Different superscripts (ab)
 6 indicate significant differences between groups at $p < 0.05$.

7



8


9 Figure 3: Hydrogen Peroxide (H₂O₂) concentration (μM). All values are expressed as
 10 mean \pm SD. Asterisk (*) means significantly different from the control group (${}^*p < 0.05$).



1

2 Figure 4: Thiobarbituric Acid Reactive Substances (TBARS) concentration. All values
3 are expressed as mean \pm SD. Asterisk (*) means significantly different from the control
4 group (* $p < 0.05$; ** $p < 0.01$).
5

6




2 Figure 5: Protein expression of catalase (CAT, MW: 60-65 kDa), glutathione reductase
3 (GR, MW: 55 kDa) and glutathione peroxidase (GPx-4, MW: 17-22 kDa); β-actin (MW:
4 43 kDa) as a control. A: detected bands corresponding interested proteins where lane 1:

1 control; lane 2: control-oil; lane 3: 150 mg/kg/d; lane 4: 300 mg/kg/d and lane 5: 600
2 mg/kg/d. B: CAT/β-actin relative expression. C: GR/β-actin relative expression. D: GPx-
3 4/β-actin relative expression. Data were expressed as a percentage of relative expression
4 respect β-actin (% ± SD). Asterisk (*) means significantly different from the control group
5 (*p < 0.05; **p < 0.01; ***p < 0.005). Different superscripts (^{ab}) indicate significant
6 differences between groups at p < 0.05.

7

8


9



1


2 Figure 6: PBPK model. A: Simulation of butylparaben (n-ButP) plasma concentrations in
 3 the rat after a subcutaneous dosing of 100 mg/kg body weight. Blue lines: 2.5 and 97.5
 4 percentiles; Gray lines: 20 simulations chosen at random from the 15000 iterations; Red
 5 lines: median prediction; the green dots indicate the mean concentrations and black lines
 6 indicate the mean \pm SD reported in (Aubert et al., 2012). B: Simulation of n-ButP plasma
 7 concentrations in the rat after a subcutaneous dosing of 100 mg/kg body weight at
 8 alternative days.

9



1

2 Figure 7: Simulation of n-ButP plasma concentrations according to PBPK model in the
 3 rats after a subcutaneous dosing of different scenarios: A. 150 mg/kg; B. 300 mg/kg; C.
 4 600 mg/kg. Blue lines: 2.5 and 97.5 percentiles; Gray lines: 20 simulations selected at
 5 random from the 15000 iterations; Red lines: median prediction.



1

2 Figure 8: Simulation of n-ButP testes concentrations according to PBPK model in the rats
 3 after a subcutaneous dosing of different scenarios: A. 150 mg/kg; B. 300 mg/kg; C. 600
 4 mg/kg. Blue lines: 2.5 and 97.5 percentiles; Gray lines: 20 simulations selected at random
 5 from the 15000 iterations; Red lines: median prediction.