

# Accepted Manuscript

Epidemiology of rare cancers and inequalities in oncologic outcomes

G. Gatta, A. Trama, R. Capocaccia

PII: S0748-7983(17)30685-6

DOI: [10.1016/j.ejso.2017.08.018](https://doi.org/10.1016/j.ejso.2017.08.018)

Reference: YEJSO 4744

To appear in: *European Journal of Surgical Oncology*


Received Date: 10 August 2017

Revised Date: 0748-7983 0748-7983

Accepted Date: 17 August 2017

Please cite this article as: Gatta G, Trama A, Capocaccia R, Epidemiology of rare cancers and inequalities in oncologic outcomes, *European Journal of Surgical Oncology* (2017), doi: 10.1016/j.ejso.2017.08.018.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Title:

**Epidemiology of rare cancers and inequalities in oncologic outcomes**

Authors:

G.Gatta <sup>1</sup>, A. Trama<sup>1</sup>, R. Capocaccia<sup>1</sup>

Affiliation:

1. Evaluative Epidemiology Unit, Fondazione IRCCS, Istituto Nazionale dei Tumori, Milan, Italy

Corresponding author:

Gemma Gatta,

Evaluative Epidemiology Unit, Via Venezian, 1, 20131 Milano, Italy;

e-mail: [gemma.gatta@istitutotumori.mi.it](mailto:gemma.gatta@istitutotumori.mi.it);

telephone number +39 02 23903518

## Summary

Rare cancers epidemiology is better known compared to the other rare diseases. Thanks to the long history of the European population-based cancer registries and to the EURO CARE huge database, the burden of rare cancers has been estimated the European (EU28) population. A considerable fraction of all cancers is represented by rare cancers (24%). They are a heterogeneous group of diseases, but they share similar problems: uncertainty of diagnosis, lack of therapies, poor research opportunities, difficulties in clinical trials, lack of expertise and of centres of reference. This paper analyses the major epidemiological indicators of frequency (incidence and prevalence) and outcome (5-year survival) of all rare cancers combined and of selected rare cancers that will be in depth treated in this monographic issue. Source of the results is the RARECAREnet search, a database publicly available. Disparities both in incidence and survival, and consequently in prevalence of rare cancers were reported across European countries. Major differences were shown in outcome: 5-year relative survival for all rare cancers together, adjusted by age and case-mix, varied from 55% or more (Italy, Germany, Belgium and Iceland) and less than 40% (Bulgaria, Lithuania and Slovakia). Similarly, for all the analyzed rare cancers, a large survival gap was observed between the Eastern and the Nordic and Central European regions. Dramatic geographical variations were assessed for curable cancers like testicular and non epithelial ovarian cancers. Geographical difference in the annual age-adjusted incidence rates for all rare cancers together varied between >140 per 100,000 (Italy, Scotland, France, Germany, and Switzerland) and < 100 (Finland, Portugal, Malta, and Poland). Prevalence, the major indicator of public health resources needs, was about 7-8 times larger than incidence. Most of rare cancers require complex surgical treatment, thus a multidisciplinary approach is essential and treatment should be provided in centres of expertise and/or in networks including expert centres. Networking is the most appropriate answer to the issues pertaining to rare cancers. Actually, in Europe, an opportunity to improve outcome and reduce disparities is provided by the creation of the European Reference Networks for rare diseases (ERNs). The Joint Action of rare cancers (JARC) is a major European initiative aimed to support the mission of the ERNs. The role of population based cancer registries still remains crucial to describe rare cancers management and outcome in the real world and to evaluate progresses made at the country and at the European level.

## Keywords:

Rare cancers, Europe, population-based cancer registry

## Introduction

In Europe, about 1 out of 4 new cancer patients has a diagnosis of rare cancer. For these patients 5-year survival is 49% and no major improvements in outcome were observed until the beginning of 2000's [1]. Rare cancers include an heterogeneous group of tumours difficult to treat because of the difficulties in performing clinical trials which lead to limited evidence to develop clinical guidelines and therefore to treat patients in a standardised and evidence-based way. In addition, because of the rarity of these tumours, it is difficult for professionals to develop the expertise to treat them especially in small size countries. The timely and correct diagnosis is not always easy to reach and treatment, even if most of rare cancers are surgically treated [2], may be complex. Thus, a multidisciplinary approach is essential as well as the treatment of these cancers in centres of expertise and/or in networks including expert centres. It is well known that survival for common cancers is better in young than in old patients, in females than in males, and that survival disparities exists across European countries [3]. One might expect the same for rare cancers. An analysis of the burden of rare cancers in Europe was provided by the RARECARE and RARECAREnet projects [4,1]. A country-based analysis of the burden of rare cancers including survival differences across EU countries is needed to suggest the best health care organisation for rare cancers. The point is crucial, because an effective organisation of the health care system could avoid unnecessary inequalities in quality of care and survival.

This paper has 2 main aims:

- 1) to analyse possible variations and inequalities in incidence, survival and prevalence across European countries for all rare cancers combined and,
- 2) to analyse the rare cancers addressed in this monographic issue to report on their incidence, survival and possible differences by age, sex and geographic regions.

The rare cancers addressed are: the epithelial tumours of head and neck (H&N), the gastroenteropancreatic (GEP) neuroendocrine tumours, the urogenital and ovarian rare cancers, soft tissue sarcomas, malignant thymoma and mesothelioma. They are relevant for different specific issues: availability of new treatments, complex treatment management, multidisciplinary approach, low outcome because none therapeutic novelties in use.

## Material and methods

All the data presented in this paper come from the online analysis tool [5] developed by the 'Information Network of Rare Cancers - RARECAREnet' project. This database contains aggregated data and is publicly available. Data source and methods are described in detail in the RARECAREnet web site. Briefly, the RARECAREnet database is a subset of EUROCARE-5, the wider collaborative study on cancer patients' survival in Europe ([www.eurocare.it](http://www.eurocare.it)). Ninety-four European population-based cancer registries (CRs) adhered to the RARECAREnet project. They provided information on cancer patients diagnosed up to 2007 and followed-up for vital status ascertainment to the end of 2008 or later. Data on incidence, survival and prevalence were produced for 198 rare cancers. They were defined combining topography and morphology codes of the International Classification of Diseases for Oncology, 3rd edition (ICD-O-3) [6]. Rare entities are grouped according to clinical management and research point of views. In this monograph we consider 41 rare cancers grouped into 7 families:

- 8 epithelial cancers of the H&N (epithelial tumours of nasal cavity and sinuses, nasopharynx and, oropharynx, tumours of major salivary glands and salivary gland type tumours, squamous cell carcinoma of hypopharynx, larynx and, oral cavity;
- 2 tumours of the thoracic tract (epithelial tumours of thymus and mesothelioma of pleura and pericardium),
- 4 ovarian cancers (rare adenocarcinomas and non epithelial tumours),
- 2 male genital tumours (epithelial tumour of penis, testicular and paratesticular tumours)
- 18 soft tissue sarcomas (STS) (head and neck, limbs, superficial trunk, mediastinum, hearth, breast, uterus, genitourinary tract, viscera, paratestis, retroperitoneum and peritoneum, pelvis, skin, periorbit, brain, embryonal rhabdomyosarcoma, alveolar rhabdomyosarcoma and Ewin's sarcoma),
- 4 GEP neuroendocrine tumours (well differentiated not functioning endocrine carcinoma, well differentiated functioning endocrine carcinoma, poorly differentiated endocrine carcinoma, mixed endocrine-exocrine carcinoma ).

Rare cancers are those with an incidence rate  $<6$  per 100,000 per year in the European population [4], thus all the selected cancer have an incidence rate  $<6$ . This paper also presents incidence, prevalence and survival for all rare cancer together, which have an interest for the public health point of view. Rational and list of all rare cancers are at <http://www.rarecarenet.eu/rarecarenet/index.php/cancerlist>.

Data are presented according to age, sex, country and 5 European regions: Northern Europe (Finland, Iceland, Norway); United Kingdom and Ireland (England, Northern Ireland, Republic of Ireland, Scotland, Wales); Central Europe (Austria, Belgium, France, Germany, Switzerland, the Netherlands); Eastern Europe (Bulgaria, Czech Republic, Estonia, Latvia, Lithuania, Poland, Slovakia) and Southern Europe (Croatia, Italy, Malta, Portugal, Slovenia, Spain).

Countries were covered by 94 CRs: 19 countries were covered by national CRs (Austria, Bulgaria, Czech Republic, Croatia, Estonia, England, Finland, Iceland, Northern Ireland, Latvia, Lithuania, Malta, Norway, Republic of Ireland, Slovakia, Slovenia, Scotland, the Netherlands, Wales); 8 countries were covered by regional CRs partially representing the population of their country (Belgium, France, Germany, Switzerland, Italy, Portugal, Poland, Spain).

The mean European population covered, over the period 2000-2007, was about 207,942,000, corresponding to 48% of the population of countries participating in RARECAREnet and 46% of the European Union population (excluding Norway, Switzerland, and Iceland, which are not EU members).

We analysed incidence and survival for rare cancers for the period 2000-2007. Crude and age-adjusted incidence rates were calculated from 83 CRs data, excluding 11 CRs collecting data only for specific cancer sites. The estimated numbers of new diagnoses in 2013 was calculated applying 2000-2007 incidence rates to the 2013 population. Relative survival estimates were obtained by the cohort approach considering patients diagnosed with a rare cancer in the period 2000-2007 and followed-up to 31st December 2008. Since all patients were included (not only those followed up for 5 years) this method is also called 'complete' cohort analysis. All the 94 CRs contributed data to survival analysis. More details of the calculation of incidence and survival are reported in the RARECAREnet website (<http://app.rarecarenet.eu/index.php>).

For the category of all rare cancers together the distribution of patients by age and rare cancers type may differ by country. Adjusted country-specific 5-year survival was then estimated by a multivariable survival model including country, age class and cancer entity as covariates.

## Results

All rare cancers together.

Table 1 shows for all rare cancers combined, the crude incidence rate, the expected number of new cases/year and the number of prevalent cases, by country. According to the population size of a country, the estimated new diagnoses per year ranged from 500 or less (Malta and Iceland) to more than 80,000 (France, Italy and Germany). The annual incidence rates (per 100,000 people) depend on the prevalence of risk factors across European countries. Among risk factors, age is the most important, therefore country variation are better compared when rates are age-adjusted (Figure 1). The age-adjusted rates varied from less than 90 (Bulgaria, Malta, Finland, Northern Ireland) to more than 130 (Germany, Belgium, Switzerland, Scotland). For incidence, all adjusted rates were higher than the crude rates (Table 1), especially in countries with young population. However, rare cancers incidence ranking of countries did not change after age adjustment. The estimated number of prevalent cases were about 8 times higher than the number of new cases per year (Table 1). Prevalent cases were more than 500,000 in Germany, France, Italy and England and less than 4,000 in Iceland and Malta. Five-year survival adjusted by age and case mix, for all rare cancer together, showed figures < 40% in Bulgaria, Lithuania and Slovakia and > 50% in the Northern countries, Germany, Switzerland, Italy, Belgium, Austria, France and, Spain (Figure 2).

Incidence of selected rare cancers.

The majority of the rare cancers studied had an annual incidence rates lower than 1 per 100,000 people, with a an estimated annual number of cases in EU28 up to about 5500 (epithelial tumour of major salivary glands) (Table 2). Less rare tumours, with incidence rate > 3, had an estimated number of new diagnoses ranging between 16,000 and 26,000 in EU28 (epithelial tumours of larynx and, epithelial tumours of oropharynx and, testicular cancers).

Survival of selected rare cancers.

Five-year relative survival was very good (>80%) for testicular cancers, STS of paratestis and skin, and non-epithelial tumours of ovary (Table 3). They accounted for only 12% of the all rare cancers here considered. The majority of these had a 5-year survival lower than 50%. Very poor outcome ( $\leq 25\%$ ) characterised several rare cancers like the epithelial tumour of hypopharynx, STS of mediastinum and hearth, undifferentiated carcinoma of thymus, and thoracic mesothelioma.

Survival by age. Five-year survival reduced with increasing age (Table 4). The decrease was marked for patients with epithelial tumours of nasopharynx, testicular cancers, visceral STS and embrional rhabdomyosarcoma of soft tissue.

Survival by sex. For the majority of epithelial rare cancers, outcome was better in women than men (Table 5), with the exception of epithelial tumours of renal pelvis and ureter and rare epithelial tumours of bladder. For STS, outcome was similar between sexes. However, women experienced a worse survival than men for STS of head and neck, breast and heart.

Geographical variation of survival. With some exception, and similarly to what shown in Figure 2, 5-years survival figures for the considered rare tumours were lower in the East Europe and highest in North and

Central Europe (Table 6). Dramatic variation was observed for testicular cancer, a highly curable tumour, in which all region presented 5-year survival higher than 95% whereas in the Eastern region survival was 88%. Similar variability emerged for non-epithelial tumour of ovary with a wide 5-year survival range between 86% (UK and Ireland) and 77% (East Europe).

ACCEPTED MANUSCRIPT



## Discussion

With this analysis, based on about 2 million rare cancer patients from 27 European countries, we outlined large disparities in the frequency of and outcome for rare cancers. Age-adjusted incidence rates (per 100,000 people) of all rare cancers combined ranged from less than 100 (Iceland, Finland, Poland, Malta, Portugal) to more than 150 in Italy. The latter high figure is due to a high incidence of thyroid, larynx and liver cancers [7], with an incidence in Italy higher than the conventional threshold of  $6 \times 100,000$  [4]. Different prevalence in the exposure to risk factors, mainly explains geographical incidence variation. However, differences in pathological diagnosis skill (NET), or in overdiagnosis (thyroid) cannot be excluded. Finally, we should consider that part of the incidence geographical variability is likely due to problems in rare cancer registration [1,4] which in turn may depend of difficulties in reaching a correct diagnosis from the pathologists [1,8]. We have estimated that a not negligible burden of new diagnoses is due to rare cancers, an important fraction among all cancers requiring specific investment from the public health organization. Prevalence is another important indicator of the burden of rare cancers for public health, representing how many people with a diagnosis of rare cancers are alive in the country population. Some of prevalent cases are in their major treating phase, others in treatment for their recurrences. Part of prevalent people can be considered cured and part of them are suffering for late effect of treatment. Prevalence indicates an heterogeneous population group who deserves different needs. Furthermore, prevalence is strongly influenced by incidence [9], which in turn means by prevalence of risk factors in the populations. In order to reduce rare cancers prevalence, investment in preventive public health programs have to be implemented.

Variation in survival for all rare cancers together appears relevant. Five-year adjusted survival of all cancers published by EUROCARE-5 [3], showed a similar geographical pattern of figure 2, for the overall rare cancers. Interestingly, the paper by Baili et al. [3] shows that adjusted survival correlated with macro-economic variables like the GDP and the TNEH. Since, in our study, survival discrepancy was only partially explained by different rare cancer incidence by country (case mix) or by different age distributions of various populations, we can speculate that investment in health care system may explain part of survival variation. Actually, five years relative survival adjusted for age distribution and case mix, is one of the most succinct indicator of cancer control performance.

International collaboration in research on and cure of rare cancers is crucial. Since most rare cancers are particularly rare [1], trial are impossible at national level and the expertise is low because of the low number of the disease in small populations. Reference centres should be strictly connected between them: patients movement between one centre to another are not so unusual in small countries or in countries without facilities for complex treatments. Strong collaboration is also relevant for experience exchanging and training as for twinning programs.

Most countries do not consider specifically rare cancers in their cancer plans. In Europe, a brilliant experience comes from the French second National Plan for Cancers, which included specific actions dedicated to the development of a network of regional centres for rare cancers patients, labelled 'reference centre for rare cancers', linked to national reference centres [10].

Survival disparities across countries were important for specific rare cancers considered by this monograph. Examples are H&N and penis cancers, NET. For H&N and penis cancers treatments are available, even if complex, an early diagnosis is crucial. Recognising borderline lesions, especially in the high risk population, is relatively simple, and greater awareness should be diffused among GPs, health care providers and general

public. Actually, low-income and disadvantaged groups are generally more exposed to avoidable risk factors such as environmental carcinogens, alcohol, tobacco use, diet and infectious agents. These groups have less access to the health services and health education programme that would empower them to make decisions to protect and improve their own health [11].

Age affects survival and about 50% of all rare cancers occurs in the elderly (65+) [5]. This was relevant for some rare cancers chosen to be described by this monograph. Actually, 50% or more cases occurred in the elderly for the epithelial tumours of nasal cavities, STS of H&N, viscera and paratestis. This point makes difficult to plan clinical trials, in which elderly are not usually included, and finally the question of disparities in accessing to proper treatment remains quite unsolved in the elderly.

There is general agreement that treatment of rare cancers should be concentrated in specialised multidisciplinary centres. However, there is scarce knowledge about the extent of centralization of rare cancers treatment at the population level. In a recent publication [1], the extent of hospital centralisation for rare cancers, at the beginning of 2000, has been described for 7 European countries and the data showed that, overall, centralisation of rare cancer treatment was generally low and varied widely between countries. Treatment of patients with epithelial tumours of the ovary, urinary tract, penis, STS, GEP neuroendocrine tumours were spread among a large number of hospitals. Slovenia and the Netherlands had, among the considered countries, the highest centralisation patterns.

To conclude, even if rare cancers account for 24% of all cancer diagnoses [1], their management do not reach optimal standards in most EU countries, and the overall survival could highly improve if all the countries could offer the best available treatments. Provision of inadequate treatments is more likely when healthcare is delivered by institutions with limited expertise, suboptimal multidisciplinary organisation of cancer care and/or low case volumes. However, since survival appears to correlate with macro-economic determinants [3], investments in the health care system may explain part of the difference in outcome and decision in the allocation of resources at national level could be facilitated with the European positive scenario. Actually, in Europe, an opportunity to reduce disparities is provided by the creation of the European Reference Networks for rare diseases (ERNs). Networking is the most appropriate answer to the issues pertaining to rare cancers. The Joint Action of rare cancers (JARC) is another major European initiative that will help the mission of the ERNs. The JARC objectives are to prioritise rare cancers in the agendas of the European Member States and to optimise the functioning of the ERNs, providing operational solutions and professional guidance in quality of care, epidemiology, research, education and state of the art definition on prevention, diagnosis and treatment of rare cancers. All of these objectives should contribute to reduce the survival gap across countries. The role of population based CRs still remain crucial to describe the real word and to evaluate progresses made at country and European level.

**Conflict of Interest**

The authors of the present study have no conflict of interest to declare.

**Funding**

This research was funded by the European Commission through the Consumers, Health, Agriculture and Food Executive Agency (Chafea); Grant No. 2000111201; Information network on rare cancers—RARECARENet. We also recognize the private donation provided by Guarneri Nicoletti for rare cancer epidemiological research.

**Acknowledgments**

We thank Lucia Buratti for technical assistance (Evaluative Epidemiology Unit, Fondazione IRCCS Istituto Nazionale Tumori, Milan, Italy).

## References


1. Gatta G, Capocaccia R, Botta L et al Burden and centralized treatment in Europe of rare tumors: results of RARECAREnet - a population-based study. *Lancet Oncol* 2017 July 4.
2. <http://www.rarecarenet.eu/rarecarenet/images/indicators> ; Report on the health care pathway study and on the updated epidemiological indicators, 2011
3. Baili P, Di Salvo F, Marcos-Gragera R, et al. Age and case mix-standardised survival for all cancer patients in Europe 1999–2007: Results of EURO CARE-5, a population-based study. *European Journal of Cancer* (2015) 51, 2120– 2129
4. Gatta G, van der Zwan JM, Casali PG, et al. Rare cancers are not so rare: the rare cancer burden in Europe. *Eur J Cancer* 2011;47: 2493-511
5. <http://app.rarecarenet.eu/analysis.php>
6. Percy C, Fritz A, Jack A, et al. International classification of diseases for oncology (ICD-O). 3rd ed. Geneva: World Health Organisation; 2000
7. Trama A, Mallone S, Ferretti S, et al. The burden of rare cancers in Italy: the surveillance of rare cancers in Italy (RITA) project, *Tumori* 2012;98(5):550-558
8. Trama A, Marcos-Gragera R, Sánchez Pérez MJ, et al. Data quality in rare cancers registration: the report of the RARECARE data quality study. *Tumori* 2017; 103: 22–32
9. Micheli A, Mugno E, Krogh V, Quinn MJ, Coleman M, Hakulinen T, Gatta G, Berrino F, Capocaccia R; EUROPREVAL Working Group. Cancer prevalence in European registry areas. *Ann Oncol.* 2002 Jun; 13(6):840-65.
10. Isabelle Ray-Coquard , Eric Pujade Lauraine , Axel Le Cesne, et al. Improving treatment results with reference centres for rare cancers: where do we stand? *European Journal of Cancer* 77 (2017).
11. Nutting CM1, Robinson M, Birchall M. Survival from laryngeal cancer in England and Wales up to 2001. *Br J Cancer.* 2008 Sep 23.

**Table 1. Crude incidence rate with 95% confidence intervals (95%CI), estimated number of new diagnoses per year and prevalence (estimated number) for all rare cancers, by European countries**

Country	All rare cancers		
	<i>incidence</i>		<i>prevalence</i>
	crude rate (95%CI)	No. of expected new cases 2013	No. of expected prevalent cases at 2008
Austria	102.280 (101.505-103.059)	10,576	84,302
Belgium	132.780 (131.754-133.812)	13,696	108,654
Bulgaria	84.363 (83.645-85.085)	9,496	79,676
Croatia	95.545 (94.530-96.568)	5,394	44,542
Czech Republic	110.483 (109.764-111.205)	12,708	101,917
Estonia	115.560 (113.544-117.603)	1,608	13,469
Finland	88.876 (87.974-89.784)	6,941	54,642
France	125.333 (124.224-126.450)	80,009	635,085
Germany	130.240 (129.645-130.837)	112,617	899,961
Iceland	104.677 (100.579-108.900)	327	2,635
Ireland	91.026 (89.991-92.070)	4,564	35,652
Italy	142.048 (141.370-142.728)	81,617	648,661
Latvia	83.440 (82.128-84.768)	2,523	22,056
Lithuania	97.684 (96.441-98.938)	3,664	31,713
Malta	86.090 (82.897-89.375)	509	3,949
Norway	113.766 (112.678-114.863)	5,746	45,097
Poland	76.161 (75.305-77.025)	43,773	351,229
Portugal	87.983 (87.173-88.799)	13,604	108,447
Slovakia	109.182 (108.130-110.243)	5,855	47,492
Slovenia	122.627 (120.916-124.356)	2,567	20,337
Spain	118.766 (117.602-119.938)	57,601	453,534
Switzerland	134.887 (133.185-136.606)	9,896	77,205
The Netherlands	122.006 (121.406-122.609)	20,349	159,546
United Kingdom	117.162 (116.856-117.469)	76,601	601,597
UK England	116.163 (115.830-116.497)	64,063	503,160
UK Northern Ireland	88.079 (86.515-89.664)	2,024	16,113
UK Scotland	134.286 (133.162-135.417)	6,577	51,738
UK Wales	121.553 (120.149-122.970)	3,937	30,586
EU28		636,753	5,085,137


Data from [www.rarecarenet.eu](http://www.rarecarenet.eu)

Figure 1. Age-adjusted annual incidence rate (per 100,000) for all rare cancers by European country and in Europe (EU). Error bars are 95% Confidence Intervals



Data elaborated from [www.rarecarenet.eu](http://www.rarecarenet.eu)

Figure 2. Five-year relative survival (%) for all rare cancers by European country and in Europe (EU). Error bars are 95% confidence intervals



note: survival adjusted by age and case-mix  
 elaboration from [www.rarecarenet.eu](http://www.rarecarenet.eu)

Table 2. No. of observed cases (obs.), crude rate with (95% CI) and No. of estimated new cases EU28 in 2013 (est.) by selected rare cancer

Family	Rare cancer entity	obs.	crude rate	est.
head and neck rare cancers	Epithelial tumours of nasal cavity and sinuses	7.046	0.450 (0.439-0.460)	2.564
	Epithelial tumours of nasopharynx	7.439	0.475 (0.464-0.486)	2580
	Epithelial tumours of major salivary glands	15.053	0.961 (0.946-0.976)	5.454
	Salivary gland type tumours of head and neck	6.741	0.430 (0.420-0.441)	2.384
	Squamous cell carcinoma with variants of hypopharynx	19.828	1.266 (1.248-1.283)	7.074
	Squamous cell carcinoma with variants of larynx	72.210	4.609 (4.576-4.643)	26.153
	Epithelial tumours of oropharynx	52.016	3.320 (3.292-3.349)	18.437
rare thoracic cancers	Squamous cell carcinoma with variants of oral cavity	54.930	3.506 (3.477-3.536)	19.670
	Epithelial tumours of thymus	2.795	0.178 (0.172-0.185)	975
rare epithelial and not epithelial ovarian cancers	Mesothelioma of pleura and pericardium	28.676	1.830 (1.809-1.852)	10.703
	Mucinous adenocarcinoma of ovary	12.066	0.770 (0.757-0.784)	4.144
	Clear cell adenocarcinoma of ovary	4.753	0.303 (0.295-0.312)	1.648
	Mullerian mixed tumour of ovary	2.255	0.144 (0.138-0.150)	798
rare male genital cancers	Non epithelial tumours of ovary	3.977	0.254 (0.246-0.262)	1.288
	Testicular and paratesticular cancers	51.605	3.294 (3.266-3.323)	16.061
soft tissue sarcomas	Epithelial tumours of penis	10.368	0.662 (0.649-0.675)	3.887
	Soft tissue sarcoma	73.795	4.711 (4.677-4.745)	25.851
	Soft tissue sarcoma of head and neck	4.087	0.261 (0.253-0.269)	1.488
	Soft tissue sarcoma of limbs	17.177	1.096 (1.080-1.113)	6.047
	Soft tissue sarcoma of superficial trunk	7.813	0.499 (0.488-0.510)	2.763
	Soft tissue sarcoma of mediastinum	465	0.030 (0.027-0.033)	160
	Soft tissue sarcoma of heart	216	0.014 (0.012-0.016)	72
	Soft tissue sarcoma of breast	2.865	0.183 (0.176-0.190)	987
	Soft tissue sarcoma of uterus	8.657	0.553 (0.541-0.564)	2.984
	Other soft tissue sarcomas of genitourinary tract	3.160	0.202 (0.195-0.209)	1.114
	Soft tissue sarcoma of viscera	6.004	0.383 (0.374-0.393)	2.145
	Soft tissue sarcoma of paratestis	510	0.033 (0.030-0.036)	185
	Soft tissue sarcoma of retroperitoneum and peritoneum	4.911	0.313 (0.305-0.322)	1.740
	Soft tissue sarcoma of pelvis	3.090	0.197 (0.190-0.204)	1.088
	Soft tissue sarcoma of skin	4.737	0.302 (0.294-0.311)	1.642
	Soft tissue sarcoma of paraorbit	117	0.007 (0.006-0.009)	40
	Soft tissue sarcoma of brain and other parts of nervous system	2.723	0.174 (0.167-0.180)	921
	Embryonal rhabdomyosarcoma of soft tissue	836	0.053 (0.050-0.057)	257
	Alveolar rhabdomyosarcoma of soft tissue	519	0.033 (0.030-0.036)	160
	Ewing's sarcoma of soft tissue	998	0.064 (0.060-0.068)	315
gastroenteropancreatic (GEP) neuroendocrine tumours	GEP - well differentiated not functioning endocrine carcinoma	15.852	1.012 (0.996-1.028)	5.599
	GEP - well differentiated functioning endocrine carcinoma	411	0.026 (0.024-0.029)	142
	GEP - poorly differentiated endocrine carcinoma	10.421	0.665 (0.652-0.678)	3.742
	GEP - mixed endocrine-exocrine carcinoma	147	0.009 (0.008-0.011)	51

Data from [www.rarecarennet.eu](http://www.rarecarennet.eu)



**Table 3. No. of cases (No.) and 5-year Relative Survival (RS%) with (95% CI) by selected rare cancers**

Family	Rare cancer entity	All ages	
		No.	5-year RS %
head and neck rare cancers	Epithelial tumours of nasal cavity and sinuses	6867	<b>47 (46-49)</b>
	Epithelial tumours of nasopharynx	7276	<b>49 (48-50)</b>
	Epithelial tumours of major salivary glands	14703	<b>61 (60-62)</b>
	Salivary gland type tumours of head and neck	6683	<b>67 (66-69)</b>
	Squamous cell carcinoma of hypopharynx	19878	<b>25 (24-26)</b>
	Squamous cell carcinoma of larynx	71928	<b>61 (60-61)</b>
	Epithelial tumours of oropharynx	50843	<b>41 (40-41)</b>
	Squamous cell carcinoma of oral cavity	54229	<b>48 (48-49)</b>
rare thoracic cancers	Epithelial tumours of thymus	2729	<b>64 (62-67)</b>
	Mesothelioma of pleura&pericardium	27893	<b>5 (4-5)</b>
rare epithelial and not epithelial ovarian cancers	Mucinous adenocarcinoma of ovary	12.010	<b>60 (59-61)</b>
	Clear cell adenocarcinoma of ovary	4.761	<b>56 (54-57)</b>
	Mullerian mixed tumour of ovary	2.242	<b>21 (20-24)</b>
	Non epithelial tumours of ovary	3.970	<b>82 (81-84)</b>
rare male genital cancers	Testicular and paratesticular cancers	51011	<b>95 (95-95)</b>
	Epithelial tumours of penis	10210	<b>68 (66-69)</b>
soft tissue sarcomas	Soft tissue sarcoma (STS)	72696	<b>57 (56-57)</b>
	STS of head and neck	4062	<b>60 (58-62)</b>
	STS of limbs	17094	<b>68 (67-69)</b>
	STS of superficial trunk	7723	<b>48 (47-50)</b>
	STS of mediastinum	457	<b>23 (19-28)</b>
	STS of heart	203	<b>14 (10-21)</b>
	STS of breast	2864	<b>75 (73-77)</b>
	STS of uterus	8568	<b>52 (51-53)</b>
	Other STSs of genitourinary tract	3107	<b>50 (48-53)</b>
	STS of viscera	5915	<b>42 (41-44)</b>
	STS of paratestis	510	<b>87 (82-92)</b>
	STS of retroperitoneum and peritoneum	4854	<b>39 (37-41)</b>
	STS of pelvis	3064	<b>47 (45-50)</b>
	STS of skin	4728	<b>90 (89-92)</b>
	STS of paraorbit	115	<b>63 (53-76)</b>
	STS of brain and other nervous system	2695	<b>55 (52-57)</b>
	gastroenteropancreatic (GEP) neuroendocrine tumours	Embryonal rhabdomyosarcoma of soft tissue	825
Alveolar rhabdomyosarcoma of soft tissue		515	<b>36 (32-41)</b>
Ewin's sarcoma of soft tissue		992	<b>45 (42-49)</b>
GEP - well differentiated not functioning endocrine carcinoma		15656	<b>72 (71-73)</b>
GEP - well differentiated functioning endocrine carcinoma	407	<b>61 (56-67)</b>	
GEP - poorly differentiated endocrine carcinoma	10456	<b>35 (34-36)</b>	
GEP - mixed endocrine-exocrine carcinoma	141	<b>26 (18-37)</b>	

Data from [www.rarecarenet.eu](http://www.rarecarenet.eu)

Table 4. Number of cases (No.) and 5-year Relative Survival (RS%) with (95% CI) for selected rare cancers by age

Family	Rare cancer entity	age class							
		0-14		15-24		25-64		65+	
		No.	5-year RS	No.	5-year RS	No.	5-year RS	No.	5-year RS
head and neck rare cancers	Epithelial tumours of nasal cavity and sinuses	8	100 (NE)	27	63 (45-88)	3,055	53 (51-55)	3,777	42 (40-45)
	Epithelial tumours of nasopharynx	69	84 (75-93)	259	73 (67-79)	4,791	55 (53-56)	2,157	31 (29-34)
	Epithelial tumours of major salivary glands	97	97 (94-100)	278	95 (92-98)	6,703	70 (69-71)	7,625	50 (49-52)
	Salivary gland type tumours of head and neck	16	94 (83-100)	101	93 (87-98)	3,574	71 (70-73)	2,992	60 (58-63)
	Squamous cell carcinoma of hypopharynx	1	0 (NE)	2	100 (NE)	13,136	27 (26-27)	6,739	22 (21-24)
	Squamous cell carcinoma of larynx	3	(NE)	24	96 (87-100)	38,749	62 (61-62)	33,155	59 (58-60)
	Epithelial tumours of oropharynx	3	100 (NE)	37	67 (53-84)	35,409	44 (43-44)	15,411	34 (33-35)
rare thoracic cancers	Squamous cell carcinoma of oral cavity	11	79 (57-100)	106	64 (55-75)	31,440	50 (49-50)	22,688	46 (45-47)
	Epithelial tumours of thymus	5	60 (30-100)	54	65 (53-80)	1,596	70 (67-73)	1,074	55 (52-60)
rare epithelial and not epithelial ovarian cancers	Mesothelioma of pleura&pericardium	1	(NE)	5	(NE)	8,936	6 (6-7)	18,951	4 (3-4)
	Mucinous adenocarcinoma of ovary	2	50 (13-100)	227	88 (84-93)	7,114	68 (67-69)	4,667	46 (44-48)
	Clear cell adenocarcinoma of ovary	0	(NE)	8	74 (49-100)	3,150	58 (56-60)	1,603	51 (48-54)
	Mullerian mixed tumour of ovary	0	(NE)	3	67 (30-100)	904	27 (24-31)	1,335	17 (15-20)
rare male genital cancers	Non epithelial tumours of ovary	302	95 (92-98)	675	92 (89-94)	2,231	84 (82-85)	762	63 (59-68)
	Testicular and paratesticular cancers	206	97 (94-100)	7,406	94 (94-95)	42,103	96 (96-96)	1,308	73 (69-77)
soft tissue sarcomas	Epithelial tumours of penis	0	(NE)	6	84 (59-100)	4,400	73 (71-74)	5,805	63 (61-65)
	Soft tissue sarcoma (STS)	2,222	68 (66-71)	2,631	64 (62-66)	37,226	64 (63-64)	30,625	46 (45-47)
	STS of head and neck	174	71 (65-79)	128	64 (55-74)	1,553	67 (64-70)	2,207	53 (50-56)
	STS of limbs	257	87 (83-92)	651	78 (74-81)	8,541	74 (73-75)	7,645	58 (57-60)
	STS of superficial trunk	145	59 (51-68)	234	59 (53-66)	3,775	57 (55-59)	3,569	37 (35-39)
	STS of mediastinum	11	63 (40-100)	20	23 (8-63)	262	27 (22-33)	164	15 (9-23)
	STS of heart	5	40 (14-100)	14	(NE)	143	15 (10-24)	41	6 (2-23)
	STS of breast	2	50 (13-100)	60	85 (76-96)	1,794	81 (79-83)	1,008	61 (57-66)
	STS of uterus	1	100 (NE)	38	78 (66-93)	5,829	59 (58-60)	2,700	36 (34-38)
	Other STSs of genitourinary tract	132	60 (52-70)	97	71 (62-81)	1,447	60 (57-63)	1,431	38 (34-41)
	STS of viscera	65	66 (55-79)	69	53 (42-67)	2,587	45 (43-47)	3,194	39 (37-41)
	STS of paratestis	5	100 (NE)	15	52 (30-91)	222	92 (87-97)	268	85 (77-94)
	STS of retroperitoneum and peritoneum	11	73 (51-100)	48	42 (28-61)	2,444	46 (44-48)	2,351	31 (28-33)
	STS of pelvis	63	62 (50-76)	93	53 (44-65)	1,569	55 (53-58)	1,339	36 (33-40)
	STS of skin	67	98 (94-100)	258	99 (97-100)	2,731	96 (95-97)	1,672	77 (73-81)
	STS of paraorbit	34	74 (59-92)	11	46 (22-96)	32	65 (49-88)	38	57 (37-88)
	STS of brain and other nervous system	147	49 (40-58)	209	59 (52-66)	1,573	59 (56-62)	766	44 (40-49)
	Embryonal rhabdomyosarcoma of soft tissue	584	76 (72-79)	148	48 (40-58)	80	40 (29-53)	13	11 (2-75)
	Alveolar rhabdomyosarcoma of soft tissue	238	49 (43-57)	128	24 (17-34)	117	24 (17-34)	32	29 (14-61)
	Ewin's sarcoma of soft tissue	167	67 (60-76)	270	51 (45-58)	423	40 (35-45)	132	21 (14-31)
gastroenteropancreatic (GEP) neuroendocrine tumours	GEP - well differentiated not functioning endocrine carcinoma	112	98 (96-100)	447	97 (95-99)	7,798	81 (79-82)	7,299	60 (59-62)
	GEP - well differentiated functioning endocrine carcinoma	2	100 (NE)	4	100 (NE)	278	68 (61-74)	123	44 (34-57)
	GEP - poorly differentiated endocrine carcinoma	15	79 (60-100)	83	71 (61-83)	4,627	45 (43-46)	5,731	26 (25-28)
	GEP - mixed endocrine-exocrine carcinoma	0	(NE)	2	100 (NE)	68	25 (15-41)	71	27 (17-44)

Data from www.rarecarenet.eu

**Table 5. Number of cases (No.) and 5-year Relative Survival (RS%) with (95% CI) for selected rare cancers, by sex**

Family	Rare cancer entity	sex	
		Male	Female
		No.	5-year RS
head and neck rare cancers	Epithelial tumours of nasal cavity and sinuses	4430	<b>46 (44-48)</b>
	Epithelial tumours of nasopharynx	5,205	<b>47 (45-49)</b>
	Epithelial tumours of major salivary glands	8,249	<b>54 (52-55)</b>
	Salivary gland type tumours of head and neck	3,701	<b>58 (56-61)</b>
	Squamous cell carcinoma of hypopharynx	17,272	<b>25 (24-25)</b>
	Squamous cell carcinoma of larynx	63,906	<b>60 (60-61)</b>
	Epithelial tumours of oropharynx	40,212	<b>38 (38-39)</b>
	Squamous cell carcinoma of oral cavity	36,623	<b>44 (44-45)</b>
rare thoracic cancers	Epithelial tumours of thymus	1,482	<b>62 (59-65)</b>
	Mesothelioma of pleura&pericardium	22,915	<b>4 (4-5)</b>
soft tissue sarcomas	Soft tissue sarcoma (STS)	32,964	<b>57 (56-57)</b>
	STS of head and neck	2,659	<b>62 (60-65)</b>
	STS of limbs	9,029	<b>67 (66-69)</b>
	STS of superficial trunk	4,236	<b>48 (47-50)</b>
	STS of mediastinum	285	<b>21 (16-27)</b>
	STS of heart	105	<b>17 (11-26)</b>
	STS of breast	35	<b>9 (61-100)</b>
	STS of uterus	-	-
	Other STSs of genitourinary tract	1,316	<b>53 (50-57)</b>
	STS of viscera	3,2	<b>38 (36-40)</b>
	STS of paratestis	510	<b>87 (82-92)</b>
	STS of retroperitoneum and peritoneum	2,311	<b>34 (32-37)</b>
	STS of pelvis	1,562	<b>48 (45-51)</b>
	STS of skin	2,595	<b>91 (89-93)</b>
	STS of paraorbit	62	<b>58 (44-77)</b>
	STS of brain and other nervous system	1,399	<b>54 (51-57)</b>
	Embryonal rhabdomyosarcoma of soft tissue	507	<b>67 (63-72)</b>
	Alveolar rhabdomyosarcoma of soft tissue	291	<b>34 (29-41)</b>
Ewing's sarcoma of soft tissue	551	<b>45 (40-50)</b>	
gastroenteropancreatic (GEP) neuroendocrine tumours	GEP - well differentiated not functioning endocrine carcinoma	7,888	<b>70 (68-71)</b>
	GEP - well differentiated functioning endocrine carcinoma	195	<b>57 (50-66)</b>
	GEP - poorly differentiated endocrine carcinoma	5,719	<b>33 (31-34)</b>
	GEP - mixed endocrine-exocrine carcinoma	77	<b>23 (14-37)</b>

Data from [www.rarecarenet.eu](http://www.rarecarenet.eu)

Table 6. Five- year relative survival (RS%) for selected rare cancers by European region

Family	Cancer entity		Northern	UK&Ireland	Centre	South	East
		No.	382	2,009	2,007	1,243	1,226
<b>H&amp;N</b>	Epithelial tumours of nasal cavity and sinuses	RS (95%CI)	<b>49 (43-56)</b>	<b>48 (45-51)</b>	<b>54 (52-57)</b>	<b>47 (43-50)</b>	<b>35 (32-39)</b>
	Epithelial tumours of nasopharynx	No.	218	1,793	1,764	2,03	1,471
		RS (95%CI)	<b>55 (47-63)</b>	<b>51 (48-54)</b>	<b>55 (52-58)</b>	<b>51 (48-53)</b>	<b>36 (33-39)</b>
	Epithelial tumours of major salivary glands and salivary-gland type tumours	No.	1,132	6,583	6,027	3,963	3,659
		RS (95%CI)	<b>74 (71-78)</b>	<b>65 (64-67)</b>	<b>65 (64-67)</b>	<b>62 (60-64)</b>	<b>52 (50-54)</b>
	Epithelial tumours of major salivary glands	No.	789	4,503	3,884	2,749	2,778
		RS (95%CI)	<b>70 (66-75)</b>	<b>63 (61-65)</b>	<b>64 (62-66)</b>	<b>61 (59-63)</b>	<b>51 (48-53)</b>
	Salivary gland type tumours of H&N	No.	343	2,088	2,154	1,216	882
		RS (95%CI)	<b>84 (79-90)</b>	<b>70 (68-73)</b>	<b>68 (65-70)</b>	<b>65 (62-69)</b>	<b>55 (51-60)</b>
	Squamous cell carcinoma of hypopharynx	No.	402	3,749	8,371	3,949	3,407
		RS (95%CI)	<b>21 (17-26)</b>	<b>26 (24-28)</b>	<b>29 (28-30)</b>	<b>26 (24-27)</b>	<b>15 (14-17)</b>
	Squamous cell carcinoma of larynx	No.	1,813	17,660	19,070	18,216	15,169
		RS (95%CI)	<b>65 (62-68)</b>	<b>64 (63-65)</b>	<b>63 (62-64)</b>	<b>63 (62-64)</b>	<b>51 (50-52)</b>
	Epithelial tumours of oropharynx	No.	1,335	12,697	19,330	8,192	9,289
		RS (95%CI)	<b>55 (52-58)</b>	<b>50 (49-51)</b>	<b>43 (42-44)</b>	<b>35 (34-37)</b>	<b>28 (27-29)</b>
	Squamous cell carcinoma oral cavity	No.	2,558	16,685	17,794	9,993	7,199
		RS (95%CI)	<b>56 (54-59)</b>	<b>53 (52-54)</b>	<b>51 (50-52)</b>	<b>47 (46-48)</b>	<b>28 (27-29)</b>
	Epithelial tumours of thymus	No.	93	778	935	595	328
<b>THORACIC</b>		RS (95%CI)	<b>50 (39-64)</b>	<b>63 (59-68)</b>	<b>67 (63-71)</b>	<b>65 (61-70)</b>	<b>62 (55-68)</b>
	Thoracic mesothelioma	No.	1,113	14,570	7,685	3,590	935
		RS (95%CI)	<b>6 (4-8)</b>	<b>3 (3-4)</b>	<b>5 (5-6)</b>	<b>7 (6-8)</b>	<b>7 (5-9)</b>
<b>GEP</b>	GEP - well diff not funct endocrine ca	No.	1,984	4,937	5,306	1,789	1,640
<b>ENDOCRINE</b>		RS (95%CI)	<b>78 (76-81)</b>	<b>67 (65-68)</b>	<b>76 (74-77)</b>	<b>75 (72-77)</b>	<b>65 (62-68)</b>
	GEP - well diff funct endocrine ca	No.	51	103	139	57	57
		RS (95%CI)	<b>40 (27-60)</b>	<b>61 (51-74)</b>	<b>71 (62-81)</b>	<b>67 (55-82)</b>	<b>50 (36-69)</b>
	GEP - poorly diff endocrine ca	No.	589	3,422	4,282	1,446	717
		RS (95%CI)	<b>35 (30-40)</b>	<b>24 (22-26)</b>	<b>44 (42-45)</b>	<b>39 (37-42)</b>	<b>26 (22-30)</b>
	GEP - mixed endocrine-exocrine ca	No.	0	6	40	32	63
		RS (95%CI)	<b>(-)</b>	<b>70 (40-100)</b>	<b>34 (20-59)</b>	<b>20 (8-49)</b>	<b>19 (10-38)</b>
<b>UROGENITAL</b>							
	Testicular & paratesticular ca	No.	2,937	16,899	16,864	6,589	7,722
		RS (95%CI)	<b>97 (97-98)</b>	<b>97 (96-97)</b>	<b>96 (96-96)</b>	<b>95 (94-95)</b>	<b>88 (88-89)</b>
	Epithelial tum penis	No.	477	3,655	2,723	1,727	1,628
		RS (95%CI)	<b>77 (71-83)</b>	<b>69 (67-72)</b>	<b>68 (66-71)</b>	<b>67 (64-70)</b>	<b>60 (57-64)</b>
		No.	3,561	23,273	21,435	13,053	11,374
<b>STS</b>	Soft tissue sarcoma	RS (95%CI)	<b>55 (53-57)</b>	<b>55 (55-56)</b>	<b>58 (57-59)</b>	<b>60 (59-61)</b>	<b>53 (52-54)</b>
		No.	6,767	38,943	32,858	15,685	21,411
<b>OVARY</b>	Epithelial tum ovary & fallopian tube	RS (95%CI)	<b>47 (46-49)</b>	<b>36 (36-37)</b>	<b>43 (43-44)</b>	<b>46 (45-47)</b>	<b>44 (43-44)</b>
	Non epithelial tum ovary	No.	179	1,069	1,038	487	1,197
		RS (95%CI)	<b>85 (79-92)</b>	<b>86 (83-88)</b>	<b>82 (80-85)</b>	<b>85 (81-88)</b>	<b>77 (74-80)</b>

Data from [www.rarecarennet.eu](http://www.rarecarennet.eu)