



# **WORKING PAPERS**

## Col·lecció "DOCUMENTS DE TREBALL DEL DEPARTAMENT D'ECONOMIA - CREIP"

An axiomatic characterization of the strong constrained egalitarian solution

Francesc Llerena Cori Vilella

Document de treball n.16 - 2012

DEPARTAMENT D'ECONOMIA – CREIP Facultat d'Economia i Empresa





### Edita:

Departament d'Economia www.fcee.urv.es/departaments/economia/publi c\_html/index.html Universitat Rovira i Virgili Facultat d'Economia i Empresa Avgda. de la Universitat, 1 43204 Reus Tel.: +34 977 759 811 Fax: +34 977 300 661 Email: <u>sde@urv.cat</u>

### CREIP

www.urv.cat/creip Universitat Rovira i Virgili Departament d'Economia Avgda. de la Universitat, 1 43204 Reus Tel.: +34 977 558 936 Email: creip@urv.cat

Adreçar comentaris al Departament d'Economia / CREIP

Dipòsit Legal: T - 886 - 2012

ISSN edició en paper: 1576 - 3382 ISSN edició electrònica: 1988 - 0820

> DEPARTAMENT D'ECONOMIA – CREIP Facultat d'Economia i Empresa

## An axiomatic characterization of the strong constrained egalitarian solution<sup>\*</sup>

Francesc Llerena and Cori Vilella Dep. de Gestió d'Empreses, CREIP and GRODE Universitat Rovira i Virgili. Av. Universitat, 1 E-43204 Reus, Spain

#### Abstract

In this paper we axiomatize the strong constrained egalitarian solution (Dutta and Ray, 1991) over the class of weak superadditive games using *constrained egalitarianism*, order-consistency, and converse order-consistency.

JEL classification: C71, C78

**keywords:** Cooperative TU-game, strong constrained egalitarian solution, axiomatization.

## 1 Introduction

On the space of transferable utility cooperative games, several solution concepts have been motivated by the idea of egalitarianism. One of the best known is the **weak constrained egalitarian solution** (WCES for short), introduced by Dutta and Ray (1989). This solution is defined under the assumption that agents believe in egalitarianism as a social value, but their individual preferences dictate selfish behavior. On the domain of convex games, the WCES has many desirable properties and it has been axiomatized by different authors (see Dutta, 1990; Klijn et al., 2000; Hougaard et al., 2001; Arin et al., 2003). Nevertheless, existence is only guaranteed for the class of convex games. In order to widen the potential class of applications, Dutta and Ray (1991) introduced the **strong constrained egalitarian** 

<sup>\*</sup>We want to thank Antonio Quesada, Carles Rafels and William Thomson for helpful comments in previous versions of this work. Financial support from research grants ECO2011-22765 (Ministerio de Ciencia e Innovación and FEDER) and 2009SGR900 (Generalitat de Catalunya) is gratefully acknowledged.

**solution** (SCES for short), a parallel concept that exists for a large class of games. However, there is no a proper characterization result for the SCES. Thus, it seems a worthwhile exercise to provide an axiomatic characterization for this solution concept. With this objective in mind, we begin introducing some notation and terminology.

The set of natural numbers  $\mathbb{N}$  denotes the universe of potential players. By  $N \subset \mathbb{N}$  we denote a finite set of players, in general  $N = \{1, \ldots, n\}$ . A **transferable utility coalitional game (a game)** is a pair (N, v) where  $v : 2^N \longrightarrow \mathbb{R}$  is the characteristic function with  $v(\emptyset) = 0$  and  $2^N$  denotes the set of all subsets (coalitions) of N. Here we only consider games with  $|N| \ge 2$ . Let  $\Gamma$  denote the set of all games. We use  $S \subset T$  to indicate strict inclusion, that is  $S \subseteq T$  but  $S \neq T$ . By |S| we denote the cardinality of the coalition  $S \subseteq N$ .

The set of **feasible payoff vectors** of a game (N, v) is defined by  $X^*(N, v) := \{x \in \mathbb{R}^N | x(N) \leq v(N)\}$ . A **solution** on a class of games  $\Gamma' \subseteq \Gamma$ , is a mapping  $\sigma$  which associates with each game  $(N, v) \in \Gamma'$  a subset  $\sigma(N, v)$  of  $X^*(N, v)$ . Notice that  $\sigma(N, v)$  is allowed to be empty. The **pre-imputation set** of a game (N, v) is defined by  $X(N, v) := \{x \in \mathbb{R}^N | x(N) = v(N)\}$ , and the set of **imputations** by  $I(N, v) := \{x \in X(N, v) | x(i) \geq v(\{i\})\}$ , for all  $i \in N\}$ . The core of (N, v) is defined by  $C(N, v) = \{x \in X(N, v) | x(S) \geq v(S)$  for all  $S \subseteq N\}$ . A game (N, v) is **convex** (Shapley, 1971) if, for every  $S, T \subseteq N, v(S) + v(T) \leq v(S \cup T) + v(S \cap T)$ . We denote by  $\Gamma_{vex}$  the class of all convex games. A game (N, v) is **weakly superadditive** if for all partition,  $\{S_1, \ldots, S_m\}$  of  $N, v(S_1) + \cdots + v(S_m) \leq v(N)$ . We denote by  $\Gamma_{ws}$  the class of all weak superadditive games. Notice that  $\Gamma_{vex} \subseteq \Gamma_{ws}$ .

Let  $\mathbb{R}^N$  stand for the space of real-valued vectors indexed by  $N, x = (x_i)_{i \in N}$ , and for all  $S \subseteq N$ ,  $x(S) = \sum_{i \in S} x_i$ , with the convention  $x(\emptyset) = 0$ . For each  $x \in \mathbb{R}^N$  and  $T \subseteq N$ ,  $x_T$  denotes the restriction of x to T:  $x_T = (x_i)_{i \in T} \in \mathbb{R}^T$ . Given two vectors  $x, y \in \mathbb{R}^N$ ,  $x \ge y$  if  $x_i \ge y_i$  for all  $i \in N$ . We say that x > y if  $x \ge y$  and for some  $j \in N, x_j > y_j$ . Moreover,  $x \gg y$  if  $x_i > y_i$  for all  $i \in N$ . For any  $x \in \mathbb{R}^N$ , denote by  $\hat{x} = (\hat{x}_1, \cdots, \hat{x}_n)$ the vector obtained from x by rearranging its coordinates in a non-decreasing order, that is,  $\hat{x}_1 \leq \hat{x}_2 \leq \ldots \leq \hat{x}_n$ . For any two vectors  $y, x \in \mathbb{R}^N$  with y(N) = x(N), we say that y **Lorenz dominates** x, denoted by  $y \succ_L x$ , if  $\sum_{j=1}^k \hat{y}_j \ge \sum_{j=1}^k \hat{x}_j$ , for all  $k \in \{1, \ldots, |N|\}$ , with at least one strict inequality. Given a coalition  $\emptyset \neq S \subseteq N$  and a set  $A \subseteq \mathbb{R}^S$ , EAdenotes the set of allocations that are Lorenz undominated within A. That is,  $EA := \{x \in A\}$ A | there is no  $y \in A$  such that  $y \succ_L x$ . Given a game (N, v), the strong Lorenz core (Dutta and Ray, 1989) is defined in a recursive way as follows: the strong Lorenz core of a singleton coalition is  $L^*(\{i\}, v) = \{v(\{i\})\}$ . Now suppose that the strong Lorenz core for all coalitions of cardinality k or less have been defined, where 1 < k < |N|. The strong Lorenz core of a coalition  $S \subset \mathbb{N}$  of size (k+1) is defined by  $L^*(S, v) = \{x \in \mathbb{R}^S \mid x(S) = \{x \in \mathbb{R}^S \mid x(S) = x(S) \}$ v(S), and there is no  $T \subset S$  and  $y \in EL^*(T, v)$  such that  $y \gg x_T$ . The weak Lorenz core is defined similarly, but replacing  $\gg$  by >. The strong constrained egalitarian

solution (Dutta and Ray, 1991) selects the Lorenz-undominated vectors within the strong Lorenz core. The **weak constrained egalitarian** solution (Dutta and Ray, 1989), denoted by **DR**, selects the Lorenz-undominated vectors within the weak Lorenz core. For  $(N, v) \in \Gamma$ ,  $|DR(N, v)| \leq 1$ .

## 2 An axiomatic characterization of the strong constrained egalitarian solution

In this section, we provide an axiomatization of the SCES based on the **consistency** principle and its converse, together with **constrained egalitarianism** (Dutta, 1990), a prescriptive property which fix the solution for two person games.

Before introducing consistency we need to define the concept of a reduced game. The terminology is taken from Thomson (2006a).

**Definition 1.** Let  $(N, v) \in \Gamma$ ,  $y \in \mathbb{R}^N$  and  $\emptyset \neq T \subset N$ . The **max reduced game** (Davis and Maschler, 1965) **relative to** T **at** y is the game  $\left(T, r_{y, DM}^T(v)\right)$  defined by

$$r_{y,\,DM}^T(v)(S) := \begin{cases} 0 & \text{if } S = \emptyset, \\ \max_{Q \subseteq N \setminus T} \left\{ v(S \cup Q) - y(Q) \right\} & \text{if } S \subset T, \\ v(N) - y(N \setminus T) & \text{if } S = T. \end{cases}$$

For an interpretation of the max reduced game see Peleg (1986).

Roughly speaking, consistency says that there is no difference in what the players of the reduced game will get in both the original game and in the reduced game. The dual of consistency is named **converse consistency**. This property states that if an efficient payoff vector is accepted for every pair of players, then it is accepted for the set of all players.<sup>1</sup> Let us now introduce formally these properties.

Let  $(\{i, j\}, v)$  be a 2-person weak-superadditive game. Without loss of generality, assume  $v(\{i\}) \leq v(\{j\})$ . The **constrained egalitarian solution** of the game  $(\{i, j\}, v)$ , denoted by  $CE(\{i, j\}, v)$ , is defined as follows:  $CE_j(\{i, j\}, v) := \max\left\{\frac{v(\{i, j\})}{2}, v(\{j\})\right\}$ , and  $CE_i(\{i, j\}, v) := v(\{i, j\}) - CE_j(\{i, j\}, v)$ .

A solution  $\sigma$  on  $\Gamma_{ws}$  satisfies

• **constrained egalitarianism** if it coincides with the constrained egalitarian solution for all 2-person weak-superadditive games.

<sup>&</sup>lt;sup>1</sup>See Thomson (2006a) for a survey on consistency and its converse

A solution  $\sigma$  on  $\Gamma'$  satisfies

- max consistency if for each  $(N, v) \in \Gamma'$ , each  $\emptyset \neq T \subset N$ , and each  $y \in \sigma(N, v)$ , then  $\left(T, r_{y, DM}^{T}(v)\right) \in \Gamma'$  and  $y_{T} \in \sigma\left(T, r_{y, DM}^{T}(v)\right)$ .
- converse max consistency if for each  $(N, v) \in \Gamma'$ ,  $y \in X(N, v)$ , and for each  $T = \{i, j\} \subseteq N$ ,  $\left(T, r_{y, DM}^{T}(v)\right) \in \Gamma'$  and  $y_{T} \in \sigma\left(T, r_{y, DM}^{T}(v)\right)$ , then  $y \in \sigma(N, v)$ .

These properties have been widely studied and appear in axiomatizations of several solution concepts. For instance, they appear among other properties in Peleg's (1986) characterization of the core. On the domain of convex games, Dutta (1990) characterizes the WCES by means of constrained egalitarianism and max consistency. The next example shows that the SCES is not *max consistent*.

#### Example 1. Dutta and Ray (1991)

Let  $(N, v) \in \Gamma$  where  $N = \{1, 2, 3\}$ , v(S) = v(N) = 1 if |S| = 2, and v(S) = 0 if |S| = 1. Here  $EL^*(N, v) = \{(0.5, 0.5, 0), (0.5, 0, 0.5), (0, 0.5, 0.5)\}$  (Dutta and Ray, 1991). Consider the max reduced game  $(\{1, 2\}, r_{x, DM}^{\{1, 2\}}(v))$ , where x = (0.5, 0.5, 0). As the reader can easy check,  $I(\{1, 2\}, r_{x, DM}^{\{1, 2\}}(v)) = \emptyset$ , thus  $EL^*(\{1, 2\}, r_{x, DM}^{\{1, 2\}}(v)) = \emptyset$ .

Hence, to characterize the SCES by means of consistency we introduced a different notion of a reduced game. To do this, we need additional definitions.

An ordering  $\theta = (i_1, i_2, \dots, i_n)$  of N where |N| = n, is a bijection from  $\{1, 2, \dots, n\}$  to N. We denote by  $\Theta_N$  the set of all orderings of N.

**Definition 2.** Given  $(N, v) \in \Gamma$  and  $\theta = (i_1, \ldots, i_n) \in \Theta_N$ , let  $x^{\theta} \in \mathbb{R}^N$  be defined as follows:

(1) 
$$x_{i_k}^{\theta} := \max_{S \in P_{i_k}} \left\{ \frac{v(S)}{|S|} \right\}, \text{ for } k = 1, \dots, n,$$

where  $P_{i_1} := \{S \subseteq N \mid i_1 \in S\}$  and  $P_{i_k} := \{S \subseteq N \mid i_1, \dots, i_{k-1} \notin S, i_k \in S\}$ , for  $k = 2, \dots, n$ .

Observe that in the construction of  $x^{\theta}$  underlines the principle of equal division. As Selten (1972) showed by a great number of experimental games, this principle is a strong distributive norm which influences the behavior of players. Thus, we can consider  $x^{\theta}$  as a vector of "natural" claims that players can require in a sequential way.

Given  $(N, v) \in \Gamma$  and  $\theta \in \Theta_N$ , let

$$\Delta^{\theta}(v) := \{ y \in X(N, v) \text{ such that } y \ge x^{\theta} \}, \text{ and}$$

$$\Theta_N^{\preceq}(v) := \{ \theta' \in \Theta_N \, | \, \nexists \, \theta \in \Theta_N \, \text{such that} \, \theta \succ_L \theta' \}$$

where  $\theta \succ_L \theta'$  means that there is  $x \in \Delta^{\theta}(v)$  such that  $x \succ_L y$ , for all  $y \in \Delta^{\theta'}(v)$ .

**Definition 3.** Let  $(N, v) \in \Gamma$ ,  $\emptyset \neq T \subset N$ ,  $\theta = (i_1, i_2, \ldots, i_n) \in \Theta_N^{\preceq}(v)$  and  $y \in \mathbb{R}^N$ . The order-reduced game on T at y is the game  $\left(T, r_{y,\theta}^T(v)\right)$ , where

$$r_{y,\theta}^{T}(v)(S) := \begin{cases} 0 & \text{if } S = \emptyset \\ \max_{Q \subseteq N \setminus T} \left\{ x^{\theta}(S \cup Q) - y(Q) \right\} & \text{if } S \subset T, \\ v(N) - y(N \setminus T) & \text{if } S = T. \end{cases}$$

In the definition of the order-reduced game, we suppose that the worth of a non empty coalition  $S \subset T$  is revaluated under the assumption that the members of S can choose the best partners in  $N \setminus T$  in order to maximize the "natural" requirements given by  $x^{\theta}$ , provided that it pays them their components of y. This is as in the Davis and Maschler (1965) definition. Assuming that all the members of N agree that the members of  $N \setminus T$  will get  $y_{N\setminus T}$ , the members of T may get  $v(N) - y(N \setminus T)$ .

A solution  $\sigma$  on  $\Gamma_{ws}$  satisfies

- order-consistency if for each  $(N, v) \in \Gamma_{ws}$  and each  $y \in \sigma(N, v)$ , there is  $\theta \in \Theta_N^{\preceq}(v)$  such that, for all  $\emptyset \neq T \subset N$ ,  $\left(T, r_{y,\theta}^T(v)\right) \in \Gamma_{ws}$  and  $y_T \in \sigma\left(T, r_{y,\theta}^T(v)\right)$ .
- converse order-consistency if for each  $(N, v) \in \Gamma_{ws}$  and  $y \in X(N, v)$ , each  $\theta \in \Theta_N^{\preceq}(v)$  and for each  $T = \{i, j\} \subseteq N$ ,  $\left(T, r_{y,\theta}^T(v)\right) \in \Gamma_{ws}$  and  $y_T \in \sigma\left(T, r_{y,\theta}^T(v)\right)$ , then  $y \in \sigma(N, v)$ .

Now we characterize the SCES.

**Theorem 2.1.** On the domain of weak superadditive games, the SCES is the only solution satisfying constrained egalitarianism, order-consistency, and converse order-consistency.

PROOF: Constrained egalitarianism follows directly from the fact that for 2-person games both the strong and the weak constrained egalitarian solution coincide. To prove the remaining axioms, we need a geometrical decomposition of the strong Lorenz core. Let  $(N, v) \in \Gamma$ , we claim that

(2) 
$$L^*(N,v) = \bigcup_{\theta \in \Theta_N} \Delta^{\theta}(v)$$

Indeed, let  $y \in L^*(N, v)$  and  $S_1 \in \arg \max_{S \subseteq N} \left\{ \frac{v(S)}{|S|} \right\}$ . Since  $y \in L^*(N, v)$ , there is  $i_1 \in S_1$ such that  $y_{i_1} \geq \frac{v(S_1)}{|S_1|}$ . Now let  $S_2 \in \arg \max_{S \subseteq N \setminus \{i_1\}} \left\{ \frac{v(S)}{|S|} \right\}$  and  $i_2 \in S_2$  such that  $y_{i_2} \geq \frac{v(S_2)}{|S_2|}$ . Following this process step by step, we can generate an order  $\theta = (i_1, i_2, \dots, i_n) \in \Theta_N$ . Let  $x^{\theta} \in \mathbb{R}^N$  as given in Definition 2. Since  $y \geq x^{\theta}$  and y(N) = v(N), we have that  $y \in \Delta^{\theta}(v)$ . To show the reverse inclusion, let  $\theta \in \Theta_N$  and  $y \in \Delta^{\theta}(v)$ . Let  $S \subset N$  be a non-empty coalition and  $i_k \in S$  the first player in S with respect to  $\theta$ . Then,  $y_{i_k} \geq x_{i_k}^{\theta} \geq \frac{v(S)}{|S|}$  and so  $y \in L^*(N, v)$ .

Next we prove order-consistency. Let  $(N, v) \in \Gamma_{ws}$  and  $y \in EL^*(N, v)$ . From expression (2) we know that there is  $\theta = (i_1, i_2, \dots, i_n) \in \Theta_N$  such that  $y \in \Delta^{\theta}(v)$ . Suppose that  $\theta \notin \Theta_N^{\preceq}(v)$ . Then, there are  $\theta' \in \Theta_N$  and  $y' \in \Delta^{\theta'}(v)$  such that  $y' \succ_L y$ , a contradiction because, from (2),  $\Delta^{\theta'}(v) \subseteq L^*(N, v)$ . Hence,  $\theta \in \Theta_N^{\preceq}(v)$ . Let  $\emptyset \neq T \subset N$ , and consider the order-reduced game  $\left(T, r_{y,\theta}^T(v)\right)$ . Now define the game  $(N, v_{\theta})$  as follows:

(3) 
$$v_{\theta}(R) := \begin{cases} x^{\theta}(R) & \text{if } R \neq N, \\ \\ v(N) & \text{if } R = N. \end{cases}$$

Since  $x^{\theta}(N) \leq v(N)$ , we have  $(N, v_{\theta}) \in \Gamma_{vex}$ . Consider the max reduced game  $\left(T, r_{y, DM}^{T}(v_{\theta})\right)$ . Notice that

(4) 
$$\left(T, r_{y,\theta}^T(v)\right) = \left(T, r_{y,DM}^T(v_{\theta})\right).$$

Clearly  $C(N, v_{\theta}) = \Delta^{\theta}(v)$ , and so  $y = DR(N, v_{\theta})$  (Dutta-Ray, 1989). Since the WCES is max consistent on the domain of convex games (Dutta, 1990),  $\left(T, r_{y,DM}^{T}(v_{\theta})\right) \in \Gamma_{ex} \subseteq \Gamma_{ws}$ and  $y_{T} = DR\left(T, r_{y,DM}^{T}(v_{\theta})\right)$ . Observe that  $C\left(T, r_{y,DM}^{T}(v_{\theta})\right) = L^{*}\left(T, r_{y,DM}^{T}(v_{\theta})\right)$ . From Dutta-Ray (1989) we know that  $y_{T} \in C\left(T, r_{y,DM}^{T}(v_{\theta})\right)$ . Thus, from (4) we conclude that  $\left(T, r_{y,\theta}^{T}(v)\right) \in \Gamma_{ws}$  and  $y_{T} \in EL^{*}\left(T, r_{y,\theta}^{T}(v)\right)$ .

Next we prove converse order-consistency. Let  $y \in X(N, v)$  and  $\theta = (i_1, i_2, \ldots, i_n) \in \Theta_N^{\leq}(v)$  such that, for all  $T = \{i_k, i_l\} \subseteq N$ ,  $\left(T, r_{y,\theta}^T(v)\right) \in \Gamma_{ws}$  and  $y_T \in EL^*\left(T, r_{y,\theta}^T(v)\right)$ . Thus, for all  $i \in N$ ,  $y_i \geq r_{y,\theta}^T(v)(\{i\}) \geq x_i^{\theta}$ . Since y(N) = v(N), we have that  $y \in \Delta^{\theta}(v)$ . Consider the game  $(N, v_{\theta})$  as defined in (3). Let  $T = \{i, j\} \subseteq N$ . As we have seen before  $\left(T, r_{y,\theta}^T(v)\right) = \left(T, r_{y,DM}^T(v_{\theta})\right)$ . For 2-person games, the SCES and WCES coincide, so  $y_T = DR\left(T, r_{y,DM}^T(v_{\theta})\right)$ . Since the WCES is converse max consistent on the domain of convex games (Dutta, 1990),  $y = DR(N, v_{\theta})$ . Moreover,  $y \in C(N, v_{\theta}) = \Delta^{\theta}(v) \subseteq L^*(N, v)$ (expression (2)). Suppose that  $y \notin EL^*(N, v)$ . Then, there is  $y' \in L^*(N, v)$  such that  $y' \succ_L y$ . From (2) there is  $\theta' \in \Theta_N$  such that  $y' \in \Delta^{\theta'}(v)$ . Since  $y' \succ_L y \succ_L x$ , for all  $x \in \Delta^{\theta}(v)$ , we get  $\theta' \succ_L \theta$ , a contradiction. Thus,  $y \in EL^*(N, v)$ . Finally, to show uniqueness, suppose there is a another solution  $\sigma$  on  $\Gamma_{ws}$  satisfying the above three axioms. For 2-person games, constrained egalitarianism implies  $\sigma = EL^*$ . Let  $(N, v) \in \Gamma_{ws}$  with  $|N| \geq 3$ . First note that constrained egalitarianism and order-consistency imply efficiency. Let  $y \in \sigma(N, v)$ . By order-consistency there is  $\theta \in \Theta_N^{\preceq}(v)$  such that, for each  $T \subset N$ , |T| = 2,  $y_T \in \sigma\left(T, r_{y,\theta}^T(v)\right)$ . By constrained egalitarianism,  $y_T \in EL^*\left(T, r_{y,\theta}^T(v)\right)$ . Now applying converse order-consistency we get  $y \in EL^*(N, v)$ . Following a symmetric argument we obtain the reverse inclusion,  $EL^*(N, v) \subseteq \sigma(N, v)$ .<sup>2</sup>

The following examples show that in Theorem 2.1 the axioms are independent:

- Let  $\sigma^1(N, v)$  be defined as follows:  $\sigma^1(N, v) := \emptyset$ , for each  $(N, v) \in \Gamma_{ws}$ . Then,  $\sigma^1$  satisfies order-consistency, converse order-consistency, but not constrained egalitarianism.
- Let  $\sigma^2(N, v)$  be defined as follows:  $\sigma^2(N, v) := CE(N, v)$  if |N| = 2 and  $\sigma^2(N, v) := X(N, v)$  if |N| > 2, for each  $(N, v) \in \Gamma_{ws}$ . Then,  $\sigma^2$  satisfies constrained egalitarianism, converse order-consistency, but not order-consistency.
- Let  $\sigma^3(N, v)$  be defined as follows:  $\sigma^3(N, v) := CE(N, v)$  if |N| = 2, and  $\sigma^3(N, v) := \emptyset$ if |N| > 2, for each  $(N, v) \in \Gamma_{ws}$ . Then,  $\sigma^3$  satisfies constrained egalitarianism, orderconsistency, but not converse order-consistency.

### References

- Arin, J., Kuipers, J., Vermeulen, D., 2003. Some characterizations of egalitarian solutions on classes of TU-games. Mathematical Social Sciences 46, 327-345.
- [2] Davis, M., Maschler, M., 1965. The kernel of a cooperative game. Naval Research Logistics Quarterly 12, 223-259.
- [3] Dutta, B., 1990. The egalitarian solution and reduced game properties in convex games. International Journal of Game Theory 19, 153-169.
- [4] Dutta, B., Ray, D., 1989. A concept of egalitarianism under participation constraints. Econometrica 57, 615-635.
- [5] Dutta, B., Ray, D., 1991. Constrained Egalitarian Allocations. Games and Economic Behavior 3, 403-422.
- [6] Hougaard, J.L., Peleg, B., Thorlund-Petersen, L., 2001. On the set of Lorenz-maximal imputations in the core of a balanced game. International Journal of Game Theory 30, 147-165.

 $<sup>^{2}</sup>$ Notice that this is a standard application of the Elevator Lemma (for more details see Thomson (2006b)).

- [7] Klijn, F., Sliker, M., Tijs, S., Zarzuelo, J., 2000. The egalitarian solutin on convex games: some characterizations. Mathematical Social Sciences 40, 111-121.
- [8] Peleg, B., 1986. On the reduced game property and its converse. International Journal of Game Theory 15, 187-200.
- [9] Selten, R., 1972. Equal share analysis of characteristic function experiments. Contributions to Experimental Economics III, J.C.B. Mohr, 130-165, In: Sauermann H (eds), Tubingen.
- [10] Shapley, L.S., 1971. Cores of convex games. International Journal of Game Theory 1, 11-16.
- [11] Thomson, W., 2006a. Consistent allocation rules. Discussion Paper. Department of Economics, University of Rochester, USA.
- [12] Thomson, W., 2006b. How to divide when there isn't enough. From the Talmud to game theory. Mimeo.