



## Changes in extreme temperature and precipitation in the Arab region: long-term trends and variability related to ENSO and NAO

Journal:	<i>International Journal of Climatology</i>
Manuscript ID:	JOC-12-0334.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	12-Mar-2013
Complete List of Authors:	<p>Donat, Markus; Climate Change Research Centre, University of New South Wales          Peterson, Thomas; National Climatic Data Center,          Brunet, M          King, Andrew; University of New South Wales, Climate Change Research Centre          Almazroui, Mansour; King Abdulaziz University, Meteorology          Kolli, Rupa Kumar; World Meteorological Organization, World Climate Applications &amp; CLIPS Division          Boucherf, Djamel; National Climatological Office Algeria,          Al-Mulla, Anwar          Nour, Abdourahman          Aly, Ahmed          Nada, Tamer          Semawi, Muhammad Mfddi; Jordan Meteorological Department,          Al Dashti, Hasan          Salhab, Tarek          El Fadli, Khalid          Muftah, Mohamed          Dah Eida, Sidaty          BADI, wafae; National meteorological service, National center for Meteorological Research          Driouech, Fatima          El Rhaz, Khalid          Abubaker, Mohammed          Ghulam, Ayman          Sanhoury Erayah, Amani          Ben Mansour, Maher          Alabdouli, Waleed          Al Dhanhani, Jemie Salem          Al Shekaili, Majed Naser; National Center for Meteorology and Seismology,</p>
Keywords:	climate extremes, climate change, observations, temperature, precipitation, ENSO, NAO

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

SCHOLARONE™  
Manuscripts

Peer Review Only

1  
2  
3  
4 **Changes in extreme temperature and precipitation in the Arab region: long-term**  
5 **trends and variability related to ENSO and NAO**  
6  
7  
8  
9

10 **(Short Title: Extreme temperature and precipitation in the Arab region)**  
11  
12  
13

14  
15 M.G. Donat<sup>1, 23</sup>, T.C. Peterson<sup>2</sup>, M. Brunet<sup>3, 22</sup>, A. D. King<sup>1, 23</sup>, M. Almazroui<sup>4</sup>, R.K.  
16 Kolli<sup>5</sup>, Djamel Boucherf<sup>6</sup>, Anwar Yousuf Al-Mulla<sup>7</sup>, Abdourahman Youssef Nour<sup>8</sup>,  
17 Ahmed Attia Aly<sup>9</sup>, Tamer Ali Ali Nada<sup>9</sup>, Muhammad M. Semawi<sup>10</sup>, Hasan Abdullah Al  
18 Dashti<sup>11</sup>, Tarek G. Salhab<sup>12</sup>, Khalid I. El Fadli<sup>13</sup>, Mohamed K. Muftah<sup>13</sup>, Sidaty Dah  
19 Eida<sup>14</sup>, Wafae Badi<sup>15</sup>, Fatima Driouech<sup>15</sup>, Khalid El Rhaz<sup>15</sup>, Mohammed J. Y.  
20 Abubaker<sup>16</sup>, Ayman S. Ghulam<sup>17</sup>, Amani Sanhoury Erayah<sup>18</sup>, Maher Ben Mansour<sup>19</sup>,  
21 Waleed O. Alabdouli<sup>20</sup>, Jemie Salem Al Dhanhani<sup>21</sup>, Majed N. Al Shekaili<sup>21</sup>  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

35 <sup>1</sup> Climate Change Research Centre, University of New South Wales, Sydney, Australia  
36

37 <sup>2</sup> NOAA's National Climatic Data Center, Asheville, NC, USA  
38

39 <sup>3</sup> Centre for Climate Change, Dep. Geography, Universitat Rovira i Virgili, Tarragona,  
40 Spain  
41  
42

43 <sup>4</sup> Center of Excellence for Climate Change Research / Dept of Meteorology, King  
44 Abdulaziz University, Jeddah, Saudi Arabia  
45 Center of Excellence for Climate Change  
46 Research, King Abdulaziz University, Jeddah, Saudi Arabia  
47  
48

49 <sup>5</sup> World Meteorological Organisation, Geneva, Switzerland  
50

51 <sup>6</sup> National Climatological Center, Algiers, Algeria  
52

53 <sup>7</sup> Transport/Civil Aviation Affairs, Manama, Kingdom of Bahrain  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3  
4<sup>8</sup> Climate Unit, Ministry of Transport, Djibouti, Republic of Djibouti  
5  
6<sup>9</sup> Ministry of Civil Aviation, Cairo, Arab Republic of Egypt  
7  
8  
9<sup>10</sup> Jordan Meteorological Department, Amman, Jordan  
10  
11<sup>11</sup> Department of Meteorology, DGCA, State of Kuwait  
12  
13<sup>12</sup> Ministry of Public Works and Transport, Beirut, Lebanon  
14  
15<sup>13</sup> Libyan National Meteorological Center, Tripoli, Libya  
16  
17<sup>14</sup> Office National de la Météorologie, Nouakchott, Mauritania  
18  
19  
20<sup>15</sup> Direction de la Meteorologie Nationale, Casablanca, Morocco  
21  
22<sup>16</sup> Palestinian Meteorological Office, Ramallah, Palestine  
23  
24<sup>17</sup> Presidency of Meteorology and Environment, Jeddah 21431, Saudi Arabia  
25  
26<sup>18</sup> Sudan Meteorological Authority (SMA), Khartoum, Sudan  
27  
28  
29<sup>19</sup> National Weather Institute, Tunis, Tunisia  
30  
31<sup>20</sup> UAE Airforce and Air Defence, Fujairah, United Arab Emirates  
32  
33<sup>21</sup> National Center of Meteorology and Seismology, Abu Dhabi, United Arab Emirates  
34  
35<sup>22</sup> Climatic Research Unit, School of Environmental Sciences, University of East  
36  
37 Anglia, Norwich, UK  
38  
39<sup>23</sup> ARC Centre of Excellence for Climate System Science, University of New South  
40  
41  
42 Wales, Sydney, Australia  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*Corresponding Author:*

Markus Donat,

Climate Change Research Centre, University of New South Wales, Sydney, Australia

Email: m.donat@unsw.edu.au

**Abstract**

A workshop was held in Casablanca, Morocco, in March 2012, to enhance knowledge of climate extremes and their changes in the Arab region. This workshop initiated intensive data compilation activities of daily observational weather station data from the Arab region. After conducting careful control processes to ensure the quality and homogeneity of the data, climate indices for extreme temperatures and precipitation were calculated.

This study examines the temporal changes in climate extremes in the Arab region with regard to long-term trends and natural variability related to ENSO and NAO. We find consistent warming trends since the middle of the 20<sup>th</sup> Century across the region. This is evident in the increased frequencies of warm days and warm nights, higher extreme temperature values, fewer cold days and cold nights and shorter cold spell durations. The warming trends seem to be particularly strong since the early 1970s. Changes in precipitation are generally less consistent and characterised by a higher spatial and temporal variability; the trends are generally less significant. However, in the western part of the Arab region, there is a tendency towards wetter conditions. In contrast, in the eastern part, there are more drying trends, although, these are of low significance.

We also find some relationships between climate extremes in the Arab region and certain prominent modes of variability, in particular El Niño-Southern Oscillation (ENSO) and North Atlantic Oscillation (NAO). The relationships of the climate extremes with NAO are stronger, in general, than those with ENSO, and are particularly strong in the western part of the Arab region (closer to the Atlantic Ocean). The

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

relationships with ENSO are found to be more significant towards the eastern part of the area of study.

Peer Review Only

## 1. Introduction

Climatic extreme events may have major impacts on society, economy, ecosystems, and on human health; they drive natural and human systems much more than the average climate (Parmesan et al., 2000). This is particularly relevant given the expectation of continuing changes in extremes. Assessing the state of the climate and science, the IPCC (Field et al., 2012) concluded that “it is likely that anthropogenic influences have led to warming of extreme daily minimum and maximum temperatures at the global scale” and that “there is medium confidence that anthropogenic influences have contributed to intensification of extreme precipitation at the global scale”. Additionally, Peterson et al. (2012) documented how anthropogenic climate change is altering the odds of extreme events occurring. Whilst some of the major extreme events that occurred in 2011 were not unusual in the context of natural variability, there were other events where an anthropogenic signal could be detected.

When monitoring observed changes in climate and, in particular, climate extremes, for many regions (particularly in Africa, South America and parts of Asia) we are still lacking suitable and comparable data (Alexander et al., 2006). To address this and try to fill the data gaps, regional workshops coordinated by the WMO Expert Team on Climate Change Detection and Indices (ETCCDI)<sup>1</sup> are regularly organised in different regions of the world (Peterson and Manton, 2008), mostly in developing countries. The workshops include hands-on sessions where participants assess the time series of daily

---

<sup>1</sup> Officially the team is the Joint-World Meteorological Organization (WMO) Commission for Climatology (CCI), Climate Variability and Predictability (CLIVAR), and Joint Technical Commission for Oceanography and Marine Meteorology (JCOMM) Expert Team on Climate Change Detection and Indices.

1  
2  
3  
4 station observations of maximum and minimum temperature as well as daily  
5  
6 precipitation data that they brought with them (once selected and extracted from their  
7  
8 national databases), under the guidance of international experts. Hence, the workshops  
9  
10 provide several benefits, including capacity-building in terms of training the participants  
11  
12 regarding data quality control, homogeneity testing and climate analysis and also in  
13  
14 gaining high-quality observational data for climate analysis. This helps to complete our  
15  
16 knowledge about how the climate is changing both regionally and globally.  
17  
18  
19

20  
21 Also, for much of the Arab region, availability of both observational data for research  
22  
23 and studies focused on analysis of changes in climate extremes is limited. Although  
24  
25 observations from a few stations are available in international archives, such as the  
26  
27 Global Historical Climatology Network (GHCN)-Daily dataset (Menne et al., 2012) or  
28  
29 the European Climate Assessment and Dataset (Klok and Klein Tank, 2009), many of  
30  
31 the records are short or have a large number of missing values. Direct personal contact  
32  
33 with local participants, both during and after the ETCCDI climate workshops, allows  
34  
35 researchers to verify suspicious values or specific station characteristics.  
36  
37  
38  
39  
40

41 From March 13<sup>th</sup> to March 16<sup>th</sup> 2012, a workshop organised by the United Nations  
42  
43 Economic and Social Commission for Western Asia (ESCWA), the League of Arab  
44  
45 States (LAS), the Arab Centre for the Studies of Arid Zones and Dry Lands (ACSAD),  
46  
47 the Swedish Meteorological and Hydrological Institute (SMHI), the World  
48  
49 Meteorological Organization (WMO) and the United Nations International Strategy for  
50  
51 Disaster Risk Reduction (UNISDR), on the topic of regional climate in the Arab region,  
52  
53 was held in Casablanca, Morocco. Representatives of the meteorological services of all  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Arab countries were invited, and participants from 17 countries were present at the  
5  
6 workshop. The participants were asked to bring long-term daily observational data  
7  
8 from their countries to analyse during the workshop, and more stations were analysed  
9  
10 afterwards.

11  
12  
13  
14  
15 The ETCCDI provides standardised software for quality control and calculation of  
16  
17 climate indices, such as the RCLimDex platform freely available at  
18  
19 <http://etccdi.pacificclimate.org/software.shtml>. The RHtestV3 program for homogeneity  
20  
21 testing temperature and precipitation series is also freely available at the same website.  
22  
23 Documentation of the indices calculation software is available in different languages,  
24  
25 such as English and French; in preparation for this workshop it was also translated into  
26  
27 Arabic.  
28  
29  
30  
31

32  
33 The Arab region extends from the Maghreb in Northwest Africa to the Arabian  
34  
35 Peninsula. While in previous climate assessments (such as IPCC), this region is usually  
36  
37 treated as parts of two continents (Africa and Asia), there are a number of common  
38  
39 climatic characteristics across the region. Most of the Arab region is located in the  
40  
41 northern hemisphere sub-tropics and thus is characterised by semi-arid to arid climate  
42  
43 conditions with generally dry and hot summers and mild winters. There are, however,  
44  
45 also a number of differences between the Arab sub-regions associated with differing  
46  
47 atmospheric circulation and rainfall patterns across the region. The Arab region is also  
48  
49 vulnerable to meteorological extreme events. Heavy rain may lead to disastrous  
50  
51 flooding, such as the devastating flood in Algier, Algeria, in November 2001 which  
52  
53 caused more than 800 deaths, or the extreme rainfall events in Jeddah, Saudi Arabia, in  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 November 2009 and January 2011 (Almazroui, 2011). Drought is also a recurring issue  
5  
6 throughout the region (e.g. Al-Qinna et al., 2011; Kaniewski et al., 2012).  
7  
8  
9

10  
11 There have not been previous Arab region-wide analyses of how the area's climate is  
12  
13 changing. One regional study focussed on changes in climate extremes over the Middle  
14  
15 East region (Zhang et al., 2005), which forms the eastern part of our investigation area.  
16  
17 This investigation reported consistent changes towards more warm and less cold  
18  
19 extremes in this region. Precipitation was found to be characterised by strong  
20  
21 interannual variability without any significant trend. In addition to this study, there  
22  
23 have been some local analyses focussing in more detail on the Arabian Peninsula, for  
24  
25 example, AlSarmi and Washington (2011) and Almazroui et al. (2012), which also  
26  
27 found warming trends in most of the station data. Almazroui et al. (2012) showed that  
28  
29 the warming during the dry season (June to September) is faster compared to the  
30  
31 warming during the wet season (November to April). In contrast, examining four  
32  
33 stations in Saudi Arabia since 1961, Alkolibi (2002) found "no discernible signs of  
34  
35 climatic change". Regional warming trends have also been reported for stations in Libya  
36  
37 (El Tantawi, 2005; El Kenawy et al., 2009; El Fadli, 2012), Sudan (Sanhoury, 2011),  
38  
39 Djibouti (Ozer and Mahamoud, 2012), and Tunisia (Dahech and Beltrando, 2012).  
40  
41 Driouech et al. (2010) documented a decrease in precipitation over the period 1958-  
42  
43 2000 in the Moulouya watershed in Morocco.  
44  
45  
46  
47  
48

49  
50 Model simulations of climate extremes in this region during the historical period show  
51  
52 similar warming trends over the past century (Sillmann et al., 2013a). These trends  
53  
54 continue in multi-model future climate projections (Sillmann et al., 2013b). The signal  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 in extreme precipitation projections is incoherent, however there is a tendency towards  
5  
6 drying. Projections of changes in the nearby Mediterranean region point to a warming  
7  
8 trend with more frequent extreme warm events (Giorgi and Lionello, 2008).  
9

10  
11  
12 This study provides a comprehensive analysis of climate extremes in the entire Arab  
13  
14 region, which has not been previously studied as a whole. It is the result of major efforts  
15  
16 regarding data collection and data quality control. After a description of the Data and  
17  
18 Methods used in this study (Section 2), we present results regarding observed long-term  
19  
20 changes in temperature and precipitation extremes in Section 3. We also investigate  
21  
22 internal variability related to teleconnections with large-scale internal variations of the  
23  
24 climate system (Section 4). We discuss our results and formulate conclusions in Section  
25  
26  
27  
28 5.  
29  
30  
31  
32

## 33 **2. Data and Methods**

34  
35 To assess changes in extremes, daily station data were used to calculate a suite of 27  
36  
37 indices developed by the ETCCDI (Frich et al., 2002; Peterson, 2005). These  
38  
39 internationally coordinated indices continue to undergo refinement (Zhang et al., 2011).  
40  
41 These climate extreme indices are calculated based on daily observational data of  
42  
43 precipitation totals and daily minimum (i.e. night-time) and maximum (i.e. daytime)  
44  
45 temperatures. Daily observations from a number of weather stations in the participating  
46  
47 countries were brought to the workshop, and additional data were also retrieved from  
48  
49 some countries during the post-workshop analysis. For some countries there were  
50  
51 restrictions regarding the sharing of daily observational data. If daily data could not be  
52  
53 provided, the country participants performed the complete analysis for all their stations  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 in close discussion with the experts, so that it was possible to share at least the  
5  
6 calculated climate indices, based on the quality-controlled daily time series.  
7  
8

9  
10  
11 Many countries in the Arab region have recently suffered from conflict and war. This  
12  
13 has affected the availability of suitable data in parts of the region. For some areas  
14  
15 affected by armed conflicts, there are long gaps of up to several years in the available  
16  
17 time series, and in some instances older data are also unavailable. For example, for the  
18  
19 Palestinian stations of Hebron, Jericho and Nablus, the data are only available to the  
20  
21 Palestinian Meteorological Office after 1996; former data are still in Jordanian,  
22  
23 Egyptian or British archives, but the Palestinian Meteorological Office experiences  
24  
25 difficulties in accessing those data. Similarly, all the stations from Lebanon that were  
26  
27 available for this study have long breaks in the 1980s and 1990s, which makes it  
28  
29 difficult to assess homogeneity. There is a high probability that the instruments were  
30  
31 changed between the periods when observations are available. For this reason, we could  
32  
33 not include data from either Palestine or Lebanon in this study. Access to suitable data  
34  
35 for Syria, Oman, Yemen, Somalia and Comoros could not be obtained for this study. To  
36  
37 avoid gaps in some sub-regions from affecting our results for the whole region, we  
38  
39 included some station data for Syria from Zhang et al. (2005), which unfortunately end  
40  
41 in 2003; an update for the past decade would be desirable if data could be made  
42  
43 available.  
44  
45  
46  
47  
48

49  
50  
51 In total, more than 100 station time series were collected and analysed during and after  
52  
53 the workshop. After careful quality control and testing for homogeneity of each time  
54  
55 series, we finally used data from 60 stations for this study. Table 1 provides an overview  
56  
57  
58  
59  
60

1  
2  
3  
4 of the stations used, which are well distributed across the northern parts of Africa, and  
5  
6 the Middle East/Arabian Peninsula region (Fig. 1).  
7  
8  
9

10 All ETCCDI workshops use standardised software to quality-control the data, test the  
11 time series for homogeneity, and calculate the extreme climate indices. This ensures  
12 straightforward comparison of results across different regions of the globe. As a first  
13 step, the station time series were subjected to a quality control algorithm, closely  
14 following the guidance outlined in Klein Tank et al. (2009), to identify suspicious values  
15 such as outliers from the seasonal climatological norm, or unreasonable values (e.g.  
16 negative precipitation amounts, or minimum temperature higher than maximum  
17 temperature). The suspicious values were then corrected or verified by the country  
18 participants, based on reviewing entries in their local climate archives and also their  
19 expertise of local weather and climate conditions.  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

35 After quality control, as a second step, all time series were carefully tested for  
36 homogeneity using the RHTestV3 program (Wang and Feng, 2009). We generally used  
37 a penalised maximal F-test (e.g. Wang, 2008a) to identify potential change points in the  
38 time series. This procedure was applied to the monthly means of daily maximum and  
39 minimum temperatures and to monthly total precipitation amounts; it is based on two-  
40 phase regression models for the detection of shifts in individual station time series  
41 (Wang, 2008b). Due to the generally large distances between the stations, we could not  
42 use testing methods that make use of reference stations. The identified potential change  
43 points were then compared with documented changes to the station, to assess whether  
44 the changes had been artificially introduced, e.g. by changes in station location or  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 instruments, or whether they may reflect natural shifts in climate. If the identified  
5  
6 significant change points occurred concurrently with documented station changes, we  
7  
8 assumed the time series to be inhomogeneous and restricted the usage of data to  
9  
10 homogeneous time periods (column 5 in Table 1). Many of the time series showed  
11  
12 potential change points that had not been documented, particularly around 1983, 1993  
13  
14 and 1998 - all years with strong El-Niño conditions. This suggests that El Niño  
15  
16 Southern Oscillation (ENSO) may have had an influence on the local climate conditions  
17  
18 in the Arab region. Therefore, the relationship between ENSO and local climate  
19  
20 extremes is also investigated in this study (Section 4).  
21  
22  
23  
24

25  
26 Adjustment for homogeneity is a complex problem (e.g. Domonkos, 2011), related to a  
27  
28 number of uncertainties. It is particularly problematic if no nearby homogeneous  
29  
30 reference station is available. Therefore, we decided not to adjust data for homogeneity  
31  
32 in the context of this study. Instead, for this paper, we used only stations that provide at  
33  
34 least 30 consecutive years of homogeneous data (Table 1) and discarded  
35  
36 inhomogeneous stations or periods. However, as 34 out of the 60 stations used here  
37  
38 exhibit inhomogeneities, which limit the length of time series for investigation, it would  
39  
40 be desirable for future work to apply a suitable method to correct for inhomogeneities in  
41  
42 these data sparse regions. Additional data from about 40 stations have been made  
43  
44 available for this study but had to be disregarded because they did not contain any  
45  
46 period of 30 years of homogeneous data. Careful adjustment for homogeneity would  
47  
48 thus also serve to include more stations in such a study.  
49  
50  
51  
52  
53  
54

55 As a final step, a set of climate indices, as recommended by the ETCCDI, was  
56  
57  
58  
59  
60

1  
2  
3  
4 calculated for each time series. Most of the indices represent different types of extreme  
5  
6 events related to particularly high or low temperature and precipitation values. The  
7  
8 indices represent different aspects of extreme climate events, such as intensity,  
9  
10 frequency and duration. An overview of the indices calculated is provided in Table 2  
11  
12 (for more detailed information, also refer to Zhang et al., 2011). Some of the indices are  
13  
14 calculated relative to certain climatic percentile values. For temperatures, the percentiles  
15  
16 are calculated relative to the time of year, i.e. they follow an annual cycle. This means  
17  
18 that, for example, a warm percentile extreme, such as TX90p or TN90p as the ETCCDI  
19  
20 defines them, is just as likely to occur in winter as in summer. We used 1981-2010 as  
21  
22 the base period for calculating the percentile values, as for this period data from most of  
23  
24 the stations were available. If more than 25 % of data during the base period were  
25  
26 missing, no percentile-based indices were calculated.  
27  
28  
29  
30  
31

32  
33 Linear trends were fitted to the time series using ordinary least squares regression. We  
34  
35 present the calculated trends at each station for two periods: (i) for 1981 to present (i.e.,  
36  
37 as long as the station provides data; see Table 1) when data from all stations are  
38  
39 available, and (ii) for 1966 to present, to show changes over a longer period but still  
40  
41 with reasonable spatial coverage derived from a satisfactory number of long-term  
42  
43 stations. To gain a more integrated picture on how the climate is changing across the  
44  
45 whole region, we also present station average time series. To account for the different  
46  
47 time periods covered by each station, first the anomalies from the 1980-2000 common  
48  
49 reference period were calculated for each station, and then the different anomaly time  
50  
51 series were averaged. Note that this 21-year reference period is slightly different from  
52  
53 the 30-year base period used for calculating the percentile values.  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 To investigate the relationships of the extreme climate indices with large-scale internal  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

To investigate the relationships of the extreme climate indices with large-scale internal variability in the climate system, we calculated correlations with ENSO- and NAO-indices. The Southern Oscillation Index (SOI) was used to represent ENSO and was calculated as the difference in standardised pressure between Tahiti and Darwin (Trenberth, 1984). Indices based on sea-surface temperatures (SSTs) could also have been used for ENSO but these produced similar and slightly weaker correlations than those found using SOI. The Hurrell NAO index (NCAR, 2012; hereafter NAOI) was used to examine the NAO relationship with the climate extremes. This index is calculated using the difference in sea level pressure between Lisbon in Portugal and Reykjavik/Stykkisholmur in Iceland. Correlations (Spearman's rank) of these two variability indices with several of the climate extreme indices were calculated for individual stations and area-averages. The climate extremes indices were de-trended before the correlations were calculated. Maps of the resulting correlation coefficients were plotted. Scatter plots of the SOI and NAOI versus the climate extreme indices were also plotted in order to examine for possible asymmetries in the relationships. The correlations were generally stronger for seasonal indices, particularly during the months December to February when both ENSO and NAO are most pronounced, rather than for annual values. We also calculated lag-correlations, but found them to be less significant.

### **3. Long-term changes in extreme climate indices**

#### **3.1. Changes in mean and extreme temperatures**

The data provide evidence for significant warming trends throughout the entire Arab

1  
2  
3  
4 region, generally reflected by more and stronger warm extremes and fewer and weaker  
5  
6 cold extremes.  
7

8  
9  
10 The annual averages of daily maximum and minimum temperatures show upward trends  
11 for most stations (Figure 2), some of them significantly. The warming trends are  
12 generally stronger during the most recent 30 years (since 1981) than for the longer  
13 period (since 1966). This is also confirmed by the region-averaged time series, which  
14 show that most of the warming has happened since the early 1970s. Averaging the  
15 temperature anomalies of all stations reveals significant warming trends for both  
16 variables across the Arab region.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 Regarding climate extreme indices, the warming trends are most significantly reflected  
29 by the frequencies of cool and warm nights and days (Figure 3). These indices count the  
30 occurrences of temperatures below the 10<sup>th</sup> percentile (cool nights/days) and above the  
31 90<sup>th</sup> percentile (warm nights/days), respectively. They are thus representative of the  
32 upper and lower tails of the distribution functions of daytime and night-time  
33 temperatures, which have experienced significant shifts in most regions of the globe  
34 (Donat and Alexander, 2012). Consistently, throughout the entire Arab region, we find  
35 decreasing numbers of cool nights and cool days and increasing numbers of warm  
36 nights and warm days. These changes are significant for most stations at the 5% level,  
37 and seem to be somewhat weaker at locations in lower latitudes, such as some Sudanese  
38 stations. Again, most of the warming appears to have happened during the past 40 years  
39 since the early 1970s. The magnitudes of the negative trends in the cool extremes are,  
40 on average, larger than the magnitudes of the positive trends in warm extremes.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 The index representing the warmest day of the year shows a tendency towards higher  
7 temperatures at most stations (Fig. 4a). Note that the annual maximum value represents  
8 only one value per year at the very upper tail of the distributions, and is therefore based  
9 on a much smaller sample, undergoing a higher variability compared to the extremes  
10 indices discussed previously. Still, the increases are significant at a number of stations  
11 and also tend to be stronger during the most recent 30 years. On average across the  
12 region, TXx values have increased by about 1°C since the 1960s. Similarly, the coldest  
13 night of the year (TNn) displays warming trends at the majority of stations (not shown).  
14 The regional average of TNn has increased by about 0.5°C since the 1960s. Changes are  
15 also found in the duration of warm and cold spells (Figs. 4b, c). While the warm spell  
16 duration index (WSDI) shows some significant increases, strong decreases are found for  
17 the cold spell duration index (CSDI).  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

### 35 **3.2. Changes in precipitation indices**

36  
37 The occurrence of extreme precipitation is characterised by much stronger temporal and  
38 spatial variability than seen in the temperature extremes. Therefore, changes in the  
39 precipitation extremes are generally less consistent between the different stations and  
40 regions, and the trends are also mostly less significant. A lower signal-to-noise ratio for  
41 precipitation in comparison to temperature indices has been found in numerous other  
42 studies for other regions around the world (e.g. Frich et al., 2002; Hegerl et al., 2004;  
43 Alexander et al., 2006)  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54

55 On average over the whole Arab region, the 1960s were wetter than any of the more  
56  
57  
58  
59  
60

1  
2  
3  
4 recent decades. So trends starting in 1960 show drying, while trends starting in 1970s  
5  
6 show no change or perhaps even a slight wetting trend. This is apparent from both the  
7  
8 total annual precipitation (PRCPTOT, Fig. 5a) and also from the frequency of days with  
9  
10 more than 10 mm of rainfall (R10mm, Fig. 5b). While particularly in the western part of  
11  
12 the region (Algeria, Morocco, Mauritania), there is a consistent tendency towards wetter  
13  
14 conditions during the past 30 years, for most of the rest of the Arab region the changes  
15  
16 in precipitation indices are generally not significant. However, another consistent  
17  
18 feature seems to be (mostly non-significant) decreases in both the PRCPTOT and  
19  
20 R10mm precipitation indices over much of the Arabian Peninsula.  
21  
22  
23  
24

25  
26 The number of consecutive dry days (CDD, Fig. 5c), as a measure for absence of  
27  
28 precipitation, also shows trends towards drier conditions. This result is particularly  
29  
30 evident in the eastern part of the Arab region; most of the stations in Egypt, Djibouti and  
31  
32 on the Arabian Peninsula show upward trends (however mostly non-significant). Given  
33  
34 the nature of rainfall in the region, this result suggests that the dry (summer) season is  
35  
36 extending in length. Across the rest of the region, i.e. most parts of North Africa, there is  
37  
38 no clear pattern in the CDD changes.  
39  
40  
41  
42

#### 43 44 **4. The effect of ENSO- and NAO-related variability on the climate extremes**

45  
46 Some of the stations show significant responses to prominent patterns of internal  
47  
48 variability of the climate system. Here, we investigate relationships of the calculated  
49  
50 climate extremes indices with indices representing the state of ENSO and NAO. Both  
51  
52 variability patterns are most pronounced during the boreal winter, which also leads to  
53  
54 the strongest correlation values during this season. Here, we present the results for the  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 seasonal averages during December, January, February (DJF) of both climate extremes  
5  
6 and variability indices.  
7  
8

9  
10  
11 The relationships between ENSO and the climate extremes indices are generally  
12  
13 stronger for the temperature indices than for the precipitation indices. In the west of the  
14  
15 Arab region, the teleconnection between ENSO and the climate extremes appears to be  
16  
17 weak. Although, several stations in Mauritania show significant negative correlations  
18  
19 between SOI and mean minimum temperature. In the east of the region there are some  
20  
21 stations with significant correlations. The strongest correlation signal is observed with  
22  
23 the diurnal temperature range (DTR) index, which combines both daily minimum and  
24  
25 maximum temperatures. The rank correlation coefficients between SOI and DTR show  
26  
27 significant positive correlations at several sites in the Arabian Peninsula and northeast  
28  
29 Africa (Fig. 6a). In La Niña seasons, the DTR tends to be greater than in El Niño  
30  
31 seasons. While mostly positive correlations are found for maximum temperatures, the  
32  
33 signal is mixed and generally weaker for minimum temperatures. The physical  
34  
35 mechanisms behind these teleconnections are not studied here but require further  
36  
37 investigation.  
38  
39  
40  
41  
42

43  
44 The relationships between NAO and the climate extremes are found to be stronger than  
45  
46 those between ENSO and extremes, particularly in the west of the region. This is to be  
47  
48 expected considering the relative proximity of the areas in which these two modes of  
49  
50 variability act. NAO negative periods are related to more southerly storm tracks which  
51  
52 sometimes affect the north of Africa. The results show that NAO seems to have more of  
53  
54 an influence on temperature extremes than precipitation extremes, and that the  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 relationships tend to be stronger with warm extremes than cool extremes. The stations  
5  
6 most strongly influenced by the NAO are largely in the west of Africa or on the  
7  
8 Mediterranean coastline. The NAOI is negatively correlated with the maximum  
9  
10 temperature for the DJF season (TXx) and in northern and western areas of Africa these  
11  
12 correlations are mostly significant (Fig. 6b). This suggests that NAO negative periods  
13  
14 are associated with higher maximum temperatures. Area-averages of each index were  
15  
16 taken for stations in the western part of the investigation area (i.e., Algeria, Libya,  
17  
18 Mauritania, Morocco and Tunisia) and plotted against the NAOI for each season from  
19  
20 1961-2010.  
21  
22  
23  
24

25  
26 The plots of area-averaged TXx and the percentage of time when maximum temperature  
27  
28 is above the 90<sup>th</sup> percentile (TX90P) show that there is an asymmetric relationship  
29  
30 between NAO and the extremes indices (Figs. 6c and 6d): The magnitudes of negative  
31  
32 NAO seasons exert a greater influence on these extreme indices than the strength of  
33  
34 positive NAO seasons do; the slopes of lines of best fit in both plots are significantly  
35  
36 different from zero only for negative NAO seasons. During NAO positive periods, the  
37  
38 Atlantic storm track is located more towards the north; therefore, it is unsurprising that  
39  
40 the magnitude of positive NAO events has little impact on climate extremes in north-  
41  
42 western Africa. Again, further study is required to examine the NAO-extremes  
43  
44 teleconnection in this region, but our results would suggest that NAO provides a degree  
45  
46 of predictability in temperature extremes in north-western Africa.  
47  
48  
49  
50  
51  
52

### 53 **5. Summary, Discussion and Conclusions**

54  
55 We present an analysis of climate extremes in the Arab region, and their changes since  
56  
57  
58  
59  
60

1  
2  
3  
4 the middle of the 20<sup>th</sup> Century. Daily observational data from weather stations across the  
5  
6 Arab region were brought together and subjected to careful assessment for quality and  
7  
8 homogeneity, before calculating climate indices representative of different aspects of  
9  
10 extreme climatic events. The results give evidence for significant changes in the  
11  
12 occurrence of climate extremes during the past five decades. There are consistent  
13  
14 warming trends across the region, most significantly seen in increasing frequencies of  
15  
16 warm days and warm nights, and fewer cool days and cool nights. Significant warming  
17  
18 trends are also found in the absolute temperature values. The changes in the  
19  
20 precipitation-based indices are generally less significant and spatially inconsistent.  
21  
22 Regional time series indicate relatively wet conditions in the 1960s and a shift towards  
23  
24 drier conditions in the early 1970s. Therefore, area-average long-term trends since 1960  
25  
26 show a tendency towards drier conditions, but little change is found since the 1970s.  
27  
28 Locally, in the western part of the region there seems to be a consistent tendency  
29  
30 towards wetter conditions during the past 30 years, whereas in the eastern part there are  
31  
32 some consistent drying trends. These observational results are in qualitative agreement  
33  
34 with climate modelling studies (e.g. Sillmann et al., 2013a).  
35  
36  
37  
38  
39  
40  
41

42 We also find relationships between climate extremes in the Arab region and NAO and  
43  
44 ENSO. While ENSO has a stronger effect on extremes in the eastern part of the Arab  
45  
46 region, NAO has a stronger influence in the western part. For NAO we show that the  
47  
48 relationship with climate extremes in much of North Africa is asymmetric in nature:  
49  
50 correlations are particularly strong for negative NAO seasons, whereas they are largely  
51  
52 non-significant for positive NAO seasons.  
53  
54

55 The relationship of the climate indices with NAO and ENSO may explain some  
56  
57  
58  
59  
60

1  
2  
3  
4 variability, however the strong consistent warming trends across the region are the  
5  
6 dominant characteristic of change.  
7  
8  
9

10 On average, the frequency of temperatures below the 10<sup>th</sup> percentile seems to decrease  
11  
12 faster than temperatures above the 90<sup>th</sup> percentile are becoming more frequent. This  
13  
14 points to a narrowing of the temperature distributions. Indeed, Donat and Alexander  
15  
16 (2012) documented how (in addition to significant shifts towards warmer conditions)  
17  
18 the variance of the temperature distributions has become smaller in the northern  
19  
20 hemisphere extra-tropics during the past 60 years, whereas variance has increased in  
21  
22 low latitudes.  
23  
24  
25  
26  
27

28 While the focus of the ETCCDI indices is primarily on climate extremes, some of them  
29  
30 do not look very far out into the extreme tails of the distribution. A case could be made  
31  
32 that the most environmentally and societally relevant extremes are those major events  
33  
34 that have return periods in excess of 20 years. However, if one only has 50 years of  
35  
36 data available to analyse, one could not make a reliable assessment of how the  
37  
38 frequency of a ~20-year return period extreme were changing as there would only be a  
39  
40 few data points for each station. As the focus of the ETCCDI is on climate change  
41  
42 detection, the indices calculated are those with return periods in the order of once every  
43  
44 10 days (for temperature). This provides enough data points to support robust  
45  
46 assessments of changes. Furthermore, as shown in Peterson et al. (2008), the statistical  
47  
48 behaviour (e.g. trend) of these extremes reflects with great accuracy the behaviour of  
49  
50 extremes that are four times as rare.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 This study fills important gaps in the global picture of how these types of climate  
5 extremes are changing. Despite the coordinated international efforts of filling in data  
6 sparse areas (see Peterson and Manton, 2008), there are still wide gaps over much of  
7 Africa and South America (e.g. Alexander et al., 2006). The data collated for this study  
8 may potentially also extend the production of new global data sets of climate extremes  
9 (Donat et al., 2012). It also shows the value of regional analysis, cross-border  
10 verification of climate change signals, and international collaboration in providing  
11 sound climate change information of value to public and private planners at all levels.  
12 Indeed, meteorological services in all 14 Arab countries participating in this analysis  
13 can use this information as part of their contributions to the Global Framework for  
14 Climate Services and as a start to future analyses.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

### 33 **Acknowledgements**

34  
35 This workshop is part of the Regional Initiative for the assessment of the impact of  
36 Climate Change on water resources and socio-economic vulnerability in the Arab  
37 Region (RICCAR) coordinated by the United Nations Economic and Social  
38 Commission for Western Asia (ESCWA). It was funded through a multi-agency project  
39 under the framework of RICCAR, by the Swedish International Development  
40 cooperation Agency (SIDA), with technical support by the World Meteorological  
41 Organization (WMO) and its Commission for Climatology (CCI) and logistic support  
42 by ESCWA and the Moroccan weather service, the Direction de la Météorologie  
43 Nationale (DMN). In particular, we acknowledge Carol Chouchani Cherfane and Tarek  
44 Sadek from ESCWA and Abdallah Mokssit, Director of DMN, Permanent  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Representative of Morocco with WMO and Third Vice-President of WMO, without  
5  
6 whose excellent organisation the workshop would not have been possible. Markus  
7  
8 Donat was also supported by Australian Research Council grant LP100200690, and  
9  
10 Andrew King is supported by Australian Research Council grant CE110001028. Manola  
11  
12 Brunet acknowledges the support of the European Community's Seventh Framework  
13  
14 Programme (FP7/2007–2013) under Grant Agreement 242093 (EURO4M: European  
15  
16 Reanalysis and Observations for Monitoring). We are grateful to Lisa Alexander for  
17  
18 helpful discussions. We also thank the two anonymous reviewers for their constructive  
19  
20 comments.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

Alexander LV, Zhang X, Peterson TC, Caesar J, Gleason B, Klein Tank AMG, Haylock M, Collins D, Trewin B, Rahim F, Tagipour A, Kumar Kolli R, Revadekar JV, Griffiths G, Vincent L, Stephenson DB, Burn J, Aguilar E, Brunet M, Taylor M, New M, Zhai P, Rusticucci M, Vazquez Aguirre JL. 2006. Global observed changes in daily climate extremes of temperature and precipitation. *Journal of Geophysical Research-Atmospheres*, **111**: D05109, DOI:10.1029/2005JD006290.

Alkolibi, FM. 2002. Possible effects of global warming on agriculture and water resources in Saudi Arabia: Impacts and responses. *Climatic Change* **54**, 225-245.

Almazroui M. 2011. Sensitivity of a regional climate model on the simulation of high intensity rainfall events over the Arabian Peninsula and around Jeddah (Saudi Arabia). *Theoretical and Applied Climatology* 104: 261–276, DOI: 10.1007/s00704-010-0387-3.

Almazroui M, Islam MN, Athar H, Jones PD, Rahman MA. 2012. Recent climate change in the Arabian Peninsula: annual rainfall and temperature analysis of Saudi Arabia for 1978–2009. *International Journal of Climatology* **32**: 953–966, DOI: 10.1002/joc.3446.

Al-Qinna, M. I., Hammouri, N. A., Obeidat, M. M., Ahmad, F. Y. 2011. Drought analysis in Jordan under current and future climates. *Climatic Change* **106**: 421-440.

AlSarmi S, Washington R. 2011. Recent observed climate change over the Arabian

1  
2  
3  
4 Peninsula, *Journal of Geophysical Research* **116**: D11109, DOI:  
5  
6 10.1029/2010JD015459.  
7  
8

9  
10 Dahech S, Beltrando G. 2012. Observed temperature evolution in the City of Sfax  
11 (Middle Eastern Tunisia) for the period 1950–2007. *Climatic Change* DOI:  
12  
13 10.1007/s10584-012-0420-x.  
14  
15

16  
17  
18 Domonkos P. 2011. Homogenising time series: beliefs, dogmas and facts. *Advances in*  
19  
20 *Science and Research* **6**: 167–172, DOI:10.5194/asr-6-167-2011.  
21  
22

23  
24  
25 Donat MG, Alexander LV. 2012. The shifting probability distribution of global daytime  
26  
27 and night-time temperatures, *Geophysical Research Letters*, **39**, L14707,  
28  
29 DOI:10.1029/2012GL052459.  
30  
31

32  
33  
34 Donat MG, Alexander LV, Yang H, Durre I, Vose R, Dunn RJH, Willett KM *et al.* 2012.  
35  
36 Updated analyses of temperature and precipitation extreme indices since the beginning  
37  
38 of the twentieth century: The HadEX2 dataset, *Journal of Geophysical Research:*  
39  
40 *Atmospheres*, 118, doi:10.1002/jgrd.50150  
41  
42  
43

44  
45  
46 Driouech F, Mahé G, Déqué M, Dieulin C, El Heirech T, Milano M, Benabdelfadel A,  
47  
48 Rouche N. 2010. Evaluation d'impacts potentiels de changements climatiques sur  
49  
50 l'hydrologie du bassin versant de la Moulouya au Maroc. *Global Change: Facing Risks*  
51  
52 *and Threats to Water Resources* (Proc. of the Sixth World FRIEND Conference, Fez,  
53  
54 Morocco, October 2010). IAHS Publ. 340, 2010.  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6 El Kenawy A, López-Moreno J, Vicente-Serrano S, Mekld M. 2009. Temperature  
7 trends in Libya over the second half of the 20th century. *Theoretical and Applied*  
8 *Climatology*, **98**, 1-8.  
9  
10

11  
12  
13  
14  
15 El Fadli, Khalid Ibrahim, 2012: Climate change over Libya and impacts on agriculture.  
16 Thesis submitted for the degree of MSc in Meteorology. Cairo University, Cairo. p. 107.  
17  
18

19  
20  
21 El Tantawi, 2005: Climate Change in Libya and Desertification of Jifara Plain. Ph D.  
22 Thesis, Johannes Gutenberg-Universität in Mainz.  
23  
24

25  
26  
27  
28 Field CB, Barros V, Stocker TF, Qin D, Dokken DJ, Ebi KL, Mastrandrea MD, Mach  
29 KJ, Plattner G-K, Allen SK, et al., eds. *Intergovernmental Panel on Climate Change:*  
30 *Managing the Risks of Extreme Events and Disasters to Advance Climate Change*  
31 *Adaptation*. Cambridge: Cambridge University Press; 2012, 1075.  
32  
33  
34  
35  
36

37  
38  
39 Frich P, Alexander LV, Della-Marta P, Gleason B, Haylock M, Klein Tank AMG,  
40 Peterson T. 2002. Observed coherent changes in climatic extremes during the second  
41 half of the 20<sup>th</sup> Century, *Climate Research* **19**: 193-212.  
42  
43  
44  
45  
46

47  
48 Giorgi F, Lionello P. 2008. Climate change projections for the Mediterranean region.  
49 *Global and Planetary Change*, **63**: 90-104, DOI: 10.1016/j.gloplacha.2007.09.005.  
50  
51  
52

53  
54  
55 Hegerl GC, Zwiers FW, Stott PA, Kharin VV. 2004. Detectability of Anthropogenic  
56  
57  
58  
59  
60

1  
2  
3  
4 Changes in Annual Temperature and Precipitation Extremes. *J. Climate*, 17: 3683–3700.  
5  
6  
7

8  
9 Kaniewski, D., Van Campo, E., Weiss, H. 2012. Drought is a recurring challenge in the  
10 Middle East. *Proceedings of the National Academy of Sciences*, **109**: 3862-3867.  
11  
12  
13

14  
15 Klein Tank, A.M.G., Zwiers F.W., Zhang X. 2009. Guidelines on Analysis of extremes  
16 in a changing climate in support of informed decisions for adaptation. WMO-TD No.  
17 1500, 56 pp. (WCDMP-72, WMO-TD/No.1500)  
18  
19  
20  
21

22  
23  
24 Klok EJ., Klein Tank AMG. 2009. Updated and extended European dataset of daily  
25 climate observations. *International Journal of Climatology*. **29 (8)**, 1182-1191 DOI:  
26 10.1002/joc.1779  
27  
28  
29  
30

31  
32  
33 Menne MJ, Durre I, Gleason BG, Houston TG, Vose RS. 2012. An overview of the  
34 Global Historical Climatology Network-Daily database. *Journal of Atmospheric and*  
35 *Oceanic Technology* **29**, 897-910.  
36  
37  
38  
39

40  
41  
42 NCAR- Climate Analysis Section provided NAO Index Data. Accessed 3<sup>rd</sup> July 2012.  
43  
44  
45

46  
47 Ozer P, Mahamoud A. 2012. Recent precipitation and temperature changes in Djibouti  
48 City. *Thirty First Greater Horn of Africa Climate Outlook Forum (GHACOF31):*  
49 *Building resilience to climate related disasters in the Greater Horn of Africa through*  
50 *regional climate forums*. [available at <http://hdl.handle.net/2268/123996>]  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Parmesan C, Root TL, Willig MR. 2000. Impacts of extreme weather and climate on  
5 terrestrial biota. *Bulletin of the American Meteorological Society* **81**, 443-50.  
6  
7

8  
9  
10 Peterson TC. 2005. Climate Change Indices. *WMO Bulletin* **54**, 83-86.  
11  
12

13  
14  
15 Peterson TC, Manton MJ. 2008. Monitoring changes in climate extremes: a tale of  
16 international collaboration. *Bulletin of the American Meteorological Society* **89**, 1266–  
17 1271, DOI: 10.1175/2008BAMS2501.1  
18  
19  
20

21  
22  
23  
24 Peterson TC, Zhang X, Brunet-India M, Vázquez-Aguirre JL. 2008. Changes in North  
25 American extremes derived from daily weather data. *Journal of Geophysical Research –*  
26 *Atmospheres*, **113**, D07113, DOI: 10.1029/2007JD009453.  
27  
28  
29

30  
31  
32  
33 Peterson TC, Stott PA, Herring S. Editors. 2012. Explaining extreme events of 2011  
34 from a climate perspective. *Bulletin of the American Meteorology Society*  
35 10.1175/BAMS-D-12-00021.1.  
36  
37  
38

39  
40  
41 Sanhuri AE. 2011. Natural Vegetation as an indicator of climate change in the white  
42 Nile State(Algitena)area. M.Sc. thesis, Sudan Academy of Science, Advanced  
43 Technology & Environmental Council.  
44  
45  
46  
47

48  
49  
50 Sillmann J, Kharin VV, Zwiers FW, Zhang X, Bronaugh D. 2013a. Climate extremes  
51 indices in the CMIP5 multi-model ensemble. Part 1: Model evaluation in the present  
52 climate. *Journal of Geophysical Research – Atmospheres*, DOI: 10.1002/jgrd.50203, in  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 press.  
5  
6  
7

8 Sillmann J, Kharin VV, Zwiers FW, Zhang X, Bronaugh D. 2013b. Climate extremes  
9 indices in the CMIP5 multi-model ensemble. Part 2: Future projections. *Journal of*  
10 *Geophysical Research – Atmospheres*, DOI: 10.1002/jgrd.50188, in press.  
11  
12  
13  
14

15  
16  
17 Trenberth K. 1984. Signal versus Noise in the Southern Oscillation. *Monthly Weather*  
18 *Review* **112**, 326-332.  
19  
20  
21

22  
23  
24 Wang, X. L. 2008a. Accounting for autocorrelation in detecting mean-shifts in climate  
25 data series using the penalized maximal t or F test. *J. Appl. Meteorol. Climatol.*, 47,  
26 2423–2444, doi:10.1175/2008JAMC1741.1.  
27  
28  
29

30  
31  
32 Wang, X. L. 2008b. Penalized maximal F test for detecting undocumented mean-shifts  
33 without trend-change. *J. Atmos. Oceanic Technol.* 25(3), 368–384,  
34 doi:10.1175/2007JTECHA982.1.  
35  
36  
37

38  
39  
40  
41 Wang, X. L., and Feng Y. 2009. RHtestsV3 user manual, report, 26 pp., Clim. Res. Div.,  
42 Atmos. Sci. and Technol. Dir., Sci. and Technol. Branch, Environ. Canada, Gatineau,  
43 Quebec, Canada. (Available at <http://cccma.seos.uvic.ca/ETCCDI/software.shtml>.)  
44  
45  
46  
47  
48

49  
50  
51 Zhang XB, Aguilar E, Sensoy S, Melkonyan H, Tagiyeva U, Ahmed N, Kutaladze N,  
52 Rahimzadeh F, Taghipour A, Hantosh TH, Albert P, Semawi M, Ali MK, Al-Shabibi  
53 MHS, Al-Oulan Z, Zatarı T, Khelet IA, Hamoud S, Sagir R, Demircan M, Eken M,  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 Adiguzel M, Alexander L, Peterson TC, Wallis T. 2005. Trends in Middle East climate  
5 extreme indices from 1950 to 2003. *Journal of Geophysical Research—Atmospheres*  
6  
7  
8 **110**, D22104. DOI: 10.1029/2005jd006181.  
9

10  
11  
12 Zhang X, Alexander LV, Hegerl GC, Jones P, Klein Tank A, Peterson TC, Trewin B,  
13  
14 Zwiers FW. 2011. Indices for monitoring changes in extremes based on daily  
15 temperature and precipitation data, *WIREs Climate Change* DOI:10.1002/wcc.147.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Figure Captions

**Figure 1:** Locations of all stations from which at least 30 years of homogeneous data were available to be included in this study.

**Figure 2:** Changes in mean annual TMAX and mean annual TMIN (unit: °C/10years). Left: linear trends after 1966, middle: linear trends after 1981, right: time series of regional averages of anomalies from the 1980-2000 period. Upward pointing triangles show increasing trends, downward pointing triangles represent decreasing trends. Significant changes ( $p \leq 0.05$ ) are indicated by filled symbols. Red colour coding indicates warming, blue indicates cooling trends.

**Figure 3:** As Figure 2, but for frequency of cool nights (TN10p), cool days (TX10p), warm nights (TN90p), and warm days (TX90p). Upward pointing triangles show increasing trends, downward pointing triangles represent decreasing trends. Significant changes ( $p \leq 0.05$ ) are indicated by filled symbols. Red colour coding indicates warming, blue indicates cooling trends (unit: % of days/10years).

**Figure 4:** As Figure 2, but for annual maximum temperature (TXx, unit: °C/10years), warm spell duration index (WSDI, unit: days/10years) and cold spell duration index (CSDI, unit: days/10years). Upward pointing triangles show increasing trends, downward pointing triangles represent decreasing trends. Significant changes ( $p \leq 0.05$ ) are indicated by filled symbols. Red colour coding indicates warming, blue indicates cooling trends.

1  
2  
3  
4 **Figure 5:** As Figure 2, but for precipitation indices total annual precipitation on wet  
5 days (PRCPTOT, unit: mm/10years), heavy precipitation days (R10mm, unit:  
6 days/10years) and consecutive dry days (CDD, unit: days/10years). Upward pointing  
7 triangles show increasing trends, downward pointing triangles represent decreasing  
8 trends. Significant changes ( $p \leq 0.05$ ) are indicated by filled symbols. Red colour coding  
9 indicates drying trends, blue indicates trends towards wetter conditions.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19

20 **Figure 6:** Relationship between chosen extreme indices and ENSO or NAO during  
21 boreal winter (DJF). (a) Spearman rank correlation between DTR and the SOI index, (b)  
22 correlation between TXx and NAO. + (-) indicate positive (negative) correlations.  
23 Significant correlations ( $p \leq 0.05$ ) are marked with a circle. Correlations are calculated  
24 for as long as each station provides homogeneous data.  
25  
26  
27  
28  
29  
30  
31 (c) scatter plot for de-trended area-average TXx anomalies during 1961-2010 in the  
32 western part of the investigation area (Mauritania, Morocco, Algeria, Tunisia, Libya)  
33 and NAO, (d) as (c) but for TX90p.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1:** List of stations included in this study. We only use stations with at least 30 years of homogeneous data. <sup>T</sup> indicates stations of which only temperature data were used, <sup>P</sup> indicates stations of which only precipitation data were used.

Country	Station	Latitude	Longitude	Period of homogeneous data
Algeria	Alger Dar-El-Beida	36.41	3.13	1971-2009
Algeria	Annaba	36.5	7.48	1970-2009
Algeria	Ghardaia	32.24	3.48	1964-2009
Algeria	Oran Sennia	35.38	0.36	1971-2009
Algeria	Tamanrasset	22.48	5.26	1950-2009
Bahrain	Bahrain Airport	26.26	50.6	1950-2008
Djibouti	Djibouti	11.57	43.15	1980-2009
Egypt	Cairo Airport	30.13	31.4	1976-2007
Egypt	Alexandria	31.2	29.88	1976-2005
Jordan	Mafraq	32.36	36.25	1964-2011
Jordan	Rwashed	32.5	38.2	1974-2011
Kuwait	Kuwait Intl Airport	29.22	47.96	1981-2011
Libya	Derna	32.4	22.72	1956-2010
Libya	Kufra	24.2	23.3	1979-2010
Libya	Misurata	32.42	15.05	1979-2010
Libya	Tazerbo	25.8	21.13	1963-2002
Libya	Tripoli AP	32.66	13.15	1980-2010
Libya	Zuara	32.88	12.08	1956-2010
Mauritania	Atar	20.51	-13.05	1965-2010
Mauritania	Bir Moghrein	25.23	-11.58	1978-2006
Mauritania	Boutilimit	17.54	-14.7	1969-2010
Mauritania	Ching	20.46	-12.36	1965-2010
Mauritania	Kaedi	16.15	-13.5	1969-2010
Mauritania	Kiffa <sup>T</sup>	16.62	-11.4	1977-2010
Mauritania	Nema <sup>T</sup>	16.61	-7.26	1974-2010
Mauritania	Nouakchott	18.08	-16.0	1965- 2010
Mauritania	Rosso	16.5	-15.77	1969-2007
Mauritania	Tidjkja	18.55	-11.43	1977-2010
Morocco	Laayoune <sup>T</sup>	27.17	-13.22	1976-2011
Morocco	Dakhla <sup>P</sup>	23.72	-15.93	1980-2011
Morocco	Kenitra <sup>P</sup>	34.3	-6.6	1960-2011
Morocco	Rabat <sup>T</sup>	34.05	-6.77	1978-2011
Morocco	Fes	33.97	-4.98	1961-2011

Morocco	Safi	32.28	-9.23	1975-2011
Morocco	Midelt	32.68	-4.73	1977-2011
Morocco	Marrakech <sup>1</sup>	31.62	-8.03	1971-2011
Morocco	Tanger <sup>1</sup>	35.72	-5.9	1972-2011
Saudi Arabia	Bisha	20	42.64	1977-2010
Saudi Arabia	Dharan	26.27	50.15	1970-2004
Saudi Arabia	Hail	27.45	41.7	1978-2011
Saudi Arabia	Jeddah	21.7	39.2	1981-2011
Saudi Arabia	Madinah	24.55	39.7	1970-2011
Saudi Arabia	Riyadh	24.7	46.74	1970-2010
Saudi Arabia	Taif	21.5	40.55	1977-2010
Sudan	Abudamed	19.32	33.2	1943-2011
Sudan	El Fasher	13.37	25.2	1940-2009
Sudan	Gadaref	14.02	35.24	1943-2009
Sudan	Juba	4.52	31.36	1950-2009
Sudan	Khartoum	15.36	32.33	1945-2009
Sudan	Senar	13.33	33.37	1950-2009
Syria	Safita	34.82	36.12	1965-2003
Syria	Kamishli	37.03	41.22	1968-2003
Tunisia	Bizerte <sup>1</sup>	37.27	9.87	1972-2009
Tunisia	Jendouba <sup>1</sup>	36.5	8.17	1973-2009
Tunisia	Mednine <sup>1</sup>	33.35	10.48	1978-2009
Tunisia	Monastir <sup>1</sup>	35.75	10.91	1969-2009
Tunisia	Tozeur <sup>1</sup>	33.93	8.13	1966-2009
UAE	Abu Dhabi	24.43	54.65	1982-2011
UAE	Dubai	25.25	55.33	1975-2011
UAE	Ras Al Khimah	25.62	55.93	1977-2011
UAE	Sharjah	25.33	55.52	1977-2011

**Table 2:** List of the ETCCDI climate indices. All indices are calculated annually, \* denotes indices which are also calculated monthly.

Index	Definition	Unit
<b>A. Temperature Intensity</b>		
TXn*	Min Tmax	Coldest daily maximum temperature °C
TNn*	Min Tmin	Coldest daily minimum temperature °C
TXx*	Max Tmax	Warmest daily maximum temperature °C
TNx*	Max Tmin	Warmest daily minimum temperature °C
DTR*	Diurnal temperature range	Mean difference between daily maximum and daily minimum temperature °C
<b>Duration</b>		
GSL	Growing season length	Annual number of days between the first occurrence of 6 consecutive days with Tmean > 5°C and first occurrence of consecutive 6 days with Tmean < 5°C. For the Northern Hemisphere this is calculated from 1 January to 31 December while for the Southern Hemisphere it is calculated from 1 July to 30 June. Days
CSDI	Cold spell duration indicator	Annual number of days with at least 6 consecutive days when Tmin < 10 <sup>th</sup> percentile Days
WSDI	Warm spell duration indicator	Annual number of days with at least 6 consecutive days when Tmax > 90 <sup>th</sup> percentile Days
<b>Frequency</b>		
TX10p*	Cool days	Share of days when Tmax < 10 <sup>th</sup> percentile % of days
TN10p*	Cool nights	Share of days when Tmin < 10 <sup>th</sup> percentile % of days
TX90p*	Warm days	Share of days when Tmax > 90 <sup>th</sup> percentile % of days
TN90p*	Warm nights	Share of days when Tmin > 90 <sup>th</sup> percentile % of days
FD	Frost days	Annual number of days when Tmin < 0°C Days
ID	Icing days	Annual number of days when Tmax < 0°C Days
SU	Summer days	Annual number of days when Tmax > 25°C Days
TR	Tropical nights	Annual number of days when Tmin > 20°C Days
<b>B. Precipitation Intensity</b>		
Rx1day*	Max 1-day precipitation	Maximum 1-day precipitation total mm
Rx5day*	Max 5-day precipitation	Maximum 5-day precipitation total mm
SDII	Simple daily intensity index	Annual total precipitation divided by the number of wet days (i.e. when precipitation ≥ 1.0 mm) mm/day

Index		Definition	Unit
R95p	Annual contribution from very wet days	Annual Sum of daily precipitation > 95 <sup>th</sup> percentile	mm
R99p	Annual contribution from extremely wet days	Annual sum of daily precipitation > 99 <sup>th</sup> percentile	mm
PRCPTOT	Annual contribution from wet days <i>Duration</i>	Annual sum of daily precipitation $\geq 1$ mm	mm
CWD	Consecutive wet days	Maximum annual number of consecutive wet days (i.e. when precipitation $\geq 1.0$ mm)	Days
CDD	Consecutive dry days	Maximum annual number of consecutive dry days (i.e. when precipitation < 1.0 mm)	Days
	<i>Frequency</i>		
R10mm	Heavy precipitation days	Annual number of days when precipitation $\geq 10$ mm	Days
R20mm	Very heavy precipitation days	Annual number of days when precipitation $\geq 20$ mm	Days
Rnnmm	Precipitation above a user-defined threshold	Annual number of days when precipitation $\geq nn$ mm (nn: user-defined threshold)	Days

## Replies to the Reviewers' comments

*We are grateful to the reviewers for providing constructive comments which have helped to clarify and improve our manuscript. Both reviewers had some useful and valuable suggestions. We have addressed all their comments, and we hope that the reviewers and the Editor will agree that it has produced a much clearer and more coherent paper. We give a point-by-point response to each of the reviewers comments below – all of our responses are in italics.*

*Since the first submission of the paper in July 2012, we updated the station data from Jordan to cover the most recent decade until 2011 as well. Concerns have been raised that Israel is not part of the “Arab Region” and as such did not participate in the workshop. In order to avoid misunderstandings, we have removed the two Israeli stations which were included (data were obtained from the GHCN-D archive) in the initial submission of the paper. These modifications did not change any of our conclusions, which also confirms the robustness of the identified trends.*

### Referee: 1

Comments to the Author

The paper is very good. I recommend that the paper be accepted to the Journal after a few minor revisions. Please see attached document.

This paper presents the changes in extreme temperature and precipitation in the Arab region. The data were prepared during and following a workshop on climate change indices held in Casablanca in March 2012. There are very few published papers about climate changes in this region. The paper is well organized and very well written. I recommend that the paper be accepted to the Journal after a few minor revisions.

1. Page 9, line 24. Is it Tantawi 2005 or El Tantawi 2005?

*We corrected the reference: El Tantawi, 2005.*

2. Page 12, line 4. Was it necessary to correct many daily temperature and precipitation after the application of the quality control procedure? Maybe a sentence can be added about the QC results.

*The quality of data and number of entries which needed correction varied strongly between the data sources. For some stations/countries almost no QC issues were detected, whereas for others it was a list of up to 100 daily entries which needed to be revisited. However, not all of them turned out to be erroneous. As the amount of revisited/corrected data is very different among the data, we feel that adding a sentence can not reflect this step, and may be confusing to the reader. However, we added some text regarding QC following the suggestions by reviewer 2.*

3. Page 12, line 28. Was any metadata available for these stations? Was it possible to retrieve the cause of any change points? Maybe a sentence can be added to provide more information regarding the results of homogeneity testing.

*For a large number of stations metadata were available, so that changes to station location/instruments could be identified as the most common cause of change points. We added some details on the homogeneity testing method, and also extended the following paragraph (at the bottom of former page 12) with some additional information regarding the outcomes of homogeneity testing.*

1 4. Page 14, line 22. Is it possible to provide a reference or a link for the Hurrell NAO index?  
2  
3

4 *We have added a reference to the Climate Data Analysis group at NCAR.*  
5

6 5. It might be helpful to provide a table showing the trend for each index for both periods 1966-  
7 2010 and 1981-2010.  
8

9 *We like the idea to provide an overview of changes in all indices, and worked on such a table.*  
10 *However, inconsistencies arise from the fact that the number of stations changes strongly with*  
11 *time. This is obvious when looking at the maps and time series in Figs 2 to 5, but we have*  
12 *concerns that this information may get lost when showing an independent table. To make sure to*  
13 *keep the paper clear and comprehensible, we decided to eventually omit this table.*  
14  
15

16 6. The triangles in Figure 2 (and other figures) are very small and it is difficult to see the trends  
17 which are not significant. Maybe the triangles should be a little bigger.  
18

19 *We optimised Figures 2, 3, 4, and 5 for map and panel sizes, as well as size of the symbols. We*  
20 *are confident that details can be seen more clearly now.*  
21  
22

23 7. It is also difficult to read the units of Figure 6 c and d.  
24

25 *We modified Figure 6 to ensure that all information is clearly readable.*  
26  
27  
28  
29  
30

31 Referee: 2

32 Comments to the Author

33 Review of "Changes in extreme temperature and precipitation in the Arab region: long-term  
34 trends and variability related to ENSO and NAO", by Donat et al.  
35  
36

37 Overview:

38 This paper describes the outcomes of a workshop held in Morocco during 2012, at which  
39 participants from Met Services from the Arab region participated in processing and analyzing  
40 daily temperature and precipitation data for their countries. These data were used to calculate  
41 indices representing aspects of changing climate extremes, and have been combined to produce a  
42 regional study. In addition, some preliminary analysis has been carried out to investigate the  
43 relationships between climate extremes over the region and potential large-scale driving  
44 mechanisms such as ENSO and the NAO.  
45  
46

47 The main novel aspects of paper are that this is the first consistent assessment of changing  
48 climate extremes over the entire North Africa and Middle East region. It updates analyses carried  
49 out previously over some parts of the region (e.g. the Middle East, Zhang et al., 2005), but also  
50 provides some useful new insights into links with large-scale modes of climate variability.  
51  
52

53 This is a well-structured and clearly written paper, of an appropriate length and with suitable  
54 references. If anything can usefully be added to the discussion, it may be worth adding text to  
55 place the results in context to both the impacts of climate change over the region, and also future  
56 projections of climate change over the region. For example, what do multi-model ensembles (e.g.  
57 CMIP3/5) say about climate change projections over the region (or sub-regions)? - is there any  
58 consensus between the models in terms of their projections, and how does this relate to the  
59 observed trends found in this study?  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*We have extended the Introduction to this study by including a paragraph on findings from climate modelling studies. We have also added a statement on the agreement of our observational results with climate model results in the Discussion.*

The description of the methods comes across as somewhat brief although I am aware that the techniques and application have been described in other extremes indices workshop papers. Perhaps a few additional citations of methods/applications could be included in this section, to direct the reader to further information?

Specific Comments:

P4, L24: ENSO and NAO are introduced here without being defined. I am not sure of the policy in terms of introducing acronyms in the abstract.

*We revised to introduce El Niño–Southern Oscillation (ENSO) and North Atlantic Oscillation (NAO) here, as abbreviations are repeatedly used after.*

P6, L28: The phrase "occurred just in 2011 are likely not to be unusual" sounds somewhat awkward.

*This sentence has been altered and we hope it is clearer now.*

P7, L4: Place brackets, or commas, around the phrase "once selected and extracted from their national databases"

*Thank you, we placed brackets around this phrase.*

P7, L20: This is the first point that the Arab region is mentioned in the main text, but it is not really defined until line 30 of page 8.

*In fact, the Arab region is described already in the next but one paragraph, just 15 lines below. The paragraph here describes the ETCCDI workshop itself which was held for the League of Arab States countries - a purely political/organisational description. We feel that it is reasonable if the geographical description of the region under investigation follows immediately after the description of the ETCCDI workshops and programs.*

P8, L30 - P9, L34: This section may benefit from more background information of regional climate change (if available). Given the proximity to the Mediterranean, and the greater availability of climate studies for the that region, it might be appropriate to comment on that here, or add something on this to the discussion section.

*We have added some references to modelling studies of relevance to this region.*

P8, L30 - P9, L34: Similarly, are there are any notable climate related events of recent decades that could be noted here? Particularly if they were associated with socio-economic impacts. This would provide enhanced motivation to justify the study.

*We included a paragraph discussing some recent extreme events.*

P9, L4: Are "more" and "less" intended to refer to frequency here, or are they intended more generally to convey changes in magnitude and frequency? It reads a little vague. e.g. should

1 "less" be "fewer" or "cold" be "cool"?

2  
3 *We do not see what this comment refers to at the indicated part of the text. However, we checked*  
4 *the use of these words in the remainder of the text.*

5  
6 P9, L24: Reference should be "El Tantawi"? Consistency with Reference section.

7  
8  
9 *Thank you. We corrected accordingly.*

10  
11 P11, L54: Perhaps be a bit more specific about what "unreasonable values" are.

12  
13 *We inserted examples for specification.*

14  
15  
16 P12, L42-56: Can any references be provided that discuss climate data quality? One example  
17 might be:

18  
19 *We included the suggested reference which describes the quality control procedure.*

20  
21 P12, L9-40: Any references that could be cited to discuss theory of and/or practical application of  
22 this type of homogeneity testing?

23  
24  
25 *We have added some details on the applied method for homogeneity testing, and also included*  
26 *some relevant references.*

27  
28  
29 P13, L18-22: Sentence here is a little unclear.

30  
31 *We rephrased this sentence.*

32  
33 P13, L29: "Station" should be plural.

34  
35  
36 *Thank you, we corrected accordingly.*

37  
38 P13, L53: Is this the first point at which the 1980-2000 common reference period is mentioned?  
39 Maybe a little more clarity is needed to note that this is different from the 1981-2010 base period  
40 for calculating percentile values.

41  
42  
43 *We included a sentence to clarify that the 31-year reference period differs slightly from the 30-*  
44 *year base period used for calculating the percentile values.*

45  
46 P14, L22: Is there a reference or weblink for the Hurrell NAO Index?

47  
48  
49 *We added a reference to NCAR.*

50  
51 P15, L17: "more significant" - clarify.

52  
53  
54 *We removed these words as the sentence already says "...warming trends are generally*  
55 *stronger..."*

56  
57 P16, L42: Mention that this difference between temp and precip has been found in other studies  
58 for many other regions around the world, and provide references.

59  
60 *We included a sentence to mention that a lower signal-to-noise ratio is common finding for*  
<http://mc.manuscriptcentral.com/joc>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*precipitation.*

P18, L6-29: Why has DTR been chosen here? Does it show a stronger relationship than for other indices? DTR is perhaps less "interesting" than other possible options, such as TXx or TN90, so would be useful to say why this is shown.

*We chose DTR as it has the strongest relationship. We agree it is perhaps not the most interesting of indices, so we discuss in more detail a few of the other extreme temperature indices.*

P19, L20-22: "magnitudes of negative NAO seasons exerts" - this seems to read oddly. Should "exerts" be "exert"? Please check.

*Thank you. We corrected to "exert".*

P20, L13: "Regional" time series?

*We reworded as suggested to "regional time series".*

P20, L31: Whereby instead of thereby? Or an alternative?

*This sentence has been separated into two.*

P20, L40: Non-significant instead of insignificant?

*We reworded as suggested.*

P21, L35: Does "behaviour" here refer to statistical behaviour, as opposed to impacts?

*We inserted a word for clarification.*

References:

Almazouri et al. (2012b) appears to be missing from text?

*This entry was still in the text from an earlier version of the text. We are grateful about the attentiveness of this reviewer, and removed this entry.*

Require a comma after "AlSarmi S".

*Thank you. We corrected this entry.*

Peterson et al. (2008a) appears to be missing from text?

*We removed this entry from the reference list.*

Zwiers (2011) appears to be missing from text?

*There is no Zwiers (2011) in the reference list. We assume that the reviewer has mistaken the last 2 lines to be an independent entry due to a page change? In fact, F.W. Zwiers is co-author of the Zhang et al. (2011) reference.*

Figure 1: It would be helpful to show the country boundaries on this map if possible, and if it does not affect the image clarity too much.

1 *As political borders are disputed in several cases in this region, we prefer not to show borders.*

2  
3  
4 Figures 2-6: It is not clear what the units of the trends are on these figures. Also, they are quite  
5 small and it is not easy to see if some symbols are filled or not.  
6

7 *We have added the units to the figure captions. We have also optimised map sizes, panel sizes*  
8 *and size of the symbols, which improves the readability of the figures.*  
9

10  
11 Figure 3: The order in which the indices are presented does not match the figure caption.  
12 Preferable to rewrite caption as "frequency of (a) cool nights (TN10p), (b) cool days (TX10p),  
13 (c) warm nights (TN90p) and (d) warm days (TX90p)." Also, perhaps the caption could re-  
14 emphasize that the red/blue colouring relates to warming/cooling, as opposed to the direction of  
15 the trend (positive/negative).  
16

17  
18 *We corrected the figure caption accordingly, and also re-emphasise the meaning of the colour*  
19 *coding.*  
20

21  
22 Figure 6: Text on (c) and (d) panels is a bit small and unclear.  
23

24 *We modified Figure 6 to ensure that all information is clearly readable.*  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





Figure 1: Locations of all stations from which at least 30 years of homogeneous data were available to be included in this study.

127x79mm (300 x 300 DPI)



24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 2: Changes in mean annual TMAX and mean annual TMIN (unit: °C/10years). Left: linear trends after 1966, middle: linear trends after 1981, right: time series of regional averages of anomalies from the 1980-2000 period. Upward pointing triangles show increasing trends, downward pointing triangles represent decreasing trends. Significant changes ( $p \leq 0.05$ ) are indicated by filled symbols. Red colour coding indicates warming, blue indicates cooling trends.

139x69mm (300 x 300 DPI)




Figure 3: As Figure 2, but for frequency of cool nights (TN10p), cool days (TX10p), warm nights (TN90p), and warm days (TX90p). Upward pointing triangles show increasing trends, downward pointing triangles represent decreasing trends. Significant changes ( $p \leq 0.05$ ) are indicated by filled symbols. Red colour coding indicates warming, blue indicates cooling trends (unit: % of days/10years).

279x279mm (300 x 300 DPI)




Figure 4: As Figure 2, but for annual maximum temperature (TXx, unit: °C/10years), warm spell duration index (WSDI, unit: days/10years) and cold spell duration index (CSDI, unit: days/10years). Upward pointing triangles show increasing trends, downward pointing triangles represent decreasing trends. Significant changes ( $p \leq 0.05$ ) are indicated by filled symbols. Red colour coding indicates warming, blue indicates cooling trends.

209x157mm (300 x 300 DPI)




Figure 5: As Figure 2, but for precipitation indices total annual precipitation on wet days (PRCPTOT, unit: mm/10years), heavy precipitation days (R10mm, unit: days/10years) and consecutive dry days (CDD, unit: days/10years). Upward pointing triangles show increasing trends, downward pointing triangles represent decreasing trends. Significant changes ( $p \leq 0.05$ ) are indicated by filled symbols. Red colour coding indicates drying trends, blue indicates trends towards wetter conditions.  
209x157mm (300 x 300 DPI)




Figure 6: Relationship between chosen extreme indices and ENSO or NAO during boreal winter (DJF). (a) Spearman rank correlation between DTR and the SOI index, (b) correlation between TXx and NAO. + (-) indicate positive (negative) correlations. Significant correlations ( $p \leq 0.05$ ) are marked with a circle.

Correlations are calculated for as long as each station provides homogeneous data.

(c) scatter plot for de-trended area-average TXx anomalies during 1961-2010 in the western part of the investigation area (Mauritania, Morocco, Algeria, Tunisia, Libya) and NAO, (d) as (c) but for TX90p.

203x147mm (300 x 300 DPI)