


Article

# On the Secure Total Domination Number of Graphs

Abel Cabrera Martínez <sup>1</sup>, Luis P. Montejano <sup>2,\*</sup> and Juan A. Rodríguez-Velázquez <sup>1</sup> 

<sup>1</sup> Departament d'Enginyeria Informàtica i Matemàtiques, Universitat Rovira i Virgili, Av. Països Catalans 26, 43007 Tarragona, Spain; abel.cabrera@urv.cat (A.C.M.); juanalberto.rodriguez@urv.cat (J.A.R.-V.)

<sup>2</sup> Euncet University Business School, Universitat Politècnica de Catalunya, 08225 Terrassa, Spain

\* Correspondence: luis.pedro.montejano@euncet.es

Received: 14 August 2019; Accepted: 12 September 2019; Published: 15 September 2019

**Abstract:** A total dominating set  $D$  of a graph  $G$  is said to be a secure total dominating set if for every vertex  $u \in V(G) \setminus D$ , there exists a vertex  $v \in D$ , which is adjacent to  $u$ , such that  $(D \setminus \{v\}) \cup \{u\}$  is a total dominating set as well. The secure total domination number of  $G$  is the minimum cardinality among all secure total dominating sets of  $G$ . In this article, we obtain new relationships between the secure total domination number and other graph parameters: namely the independence number, the matching number and other domination parameters. Some of our results are tight bounds that improve some well-known results.

**Keywords:** secure total domination; secure domination; independence number; matching number; domination

**MSC:** 05C69; 05C70

## 1. Introduction

The following approach to the protection of a graph was proposed by Cockayne et al. [1]. Suppose that one or more entities are stationed at some of the vertices of a graph  $G$  and that an entity at a vertex can deal with a problem at any vertex in its closed neighbourhood. In general, an entity could consist of an observer, a robot, a guard, a legion, and so on. Informally, we say that  $G$  is protected under a given placement of entities if there exists at least one entity available to handle a problem at any vertex. The simplest cases of graph protection are those in which you can locate at most one entity per vertex. In such a case, the set of vertices containing the entities is said to be a dominating set.

In a graph  $G = (V(G), E(G))$ , a vertex dominates itself and its neighbours. A subset  $S \subseteq V(G)$  is said to be a *dominating set* of  $G$  if  $S$  dominates every vertex of  $G$ , while  $S$  is said to be a *total dominating set* if every vertex  $v \in V(G)$  is dominated by at least one vertex in  $S \setminus \{v\}$ . As usual, the neighbourhood of a vertex  $v \in V(G)$  will be denoted by  $N(v)$ . Now, a set  $S \subseteq V(G)$  is said to be a *secure (total) dominating set* if  $S$  is a (total) dominating set and for every  $v \in V(G) \setminus S$  there exists  $u \in N(v) \cap S$  such that  $(S \cup \{v\}) \setminus \{u\}$  is a (total) dominating set. In the case of secure (total) domination, the graph is deemed protected by a (total) dominating set and when an entity moves (to deal with a problem) to a neighbour not included in the (total) dominating set, the new set of entities obtained from the movement of the entity is a (total) dominating set which protects the graph as well.

The minimum cardinality among all dominating sets of  $G$  is the *domination number* of  $G$ , denoted by  $\gamma(G)$ . The *total domination number*, the *secure domination number* and the *secure total domination number* of  $G$  are defined by analogy, and are denoted by  $\gamma_t(G)$ ,  $\gamma_s(G)$  and  $\gamma_{st}(G)$ , respectively.

The domination number and the total domination number have been extensively studied. For instance, we cite the following books [2–4]. The secure domination number, which has been less studied, was introduced by Cockayne et al. in [1] and studied further in several works including [5–10], while the secure total domination number was introduced by Benecke et al. in [11] and studied further in [9,12–14].

In this work we study the relationships between the secure total domination number and other graph parameters. The article is organized as follows. In Section 2 we define key terms and additional notation. In Section 3 we show that  $\gamma_{st}(G) \leq \alpha(G) + \gamma(G)$ , where  $\alpha(G)$  denotes the independence number of  $G$ . Since  $\gamma(G) \leq \alpha(G)$ , this result improves the bound  $\gamma_{st}(G) \leq 2\alpha(G)$  obtained in [14]. Section 4 is devoted to the study of relationships between the secure total domination number and other domination parameters. In particular, we outline some known results that become tools to derive new ones. Finally, in Section 5 we obtain several bounds on the secure total domination number in terms of the matching number and other graph parameters.

## 2. Some Additional Concepts and Notation

All graphs considered in this paper are finite and undirected, without loops or multiple edges. The minimum degree of a graph  $G$  will be denoted by  $\delta(G)$  and the maximum degree by  $\Delta(G)$ . As usual, the *closed neighbourhood* of a vertex  $v \in V(G)$  is denoted by  $N[v] = N(v) \cup \{v\}$ . We say that a vertex  $v \in V(G)$  is a *universal vertex* of  $G$  if  $N[v] = V(G)$ . By analogy with the notation used for vertices, for a set  $S \subseteq V(G)$ , its *open neighbourhood* is the set  $N(S) = \cup_{v \in S} N(v)$ , and its *closed neighbourhood* is the set  $N[S] = N(S) \cup S$ . We also define the following sets associated with  $v \in V(G)$ .

- The *internal private neighbourhood* of  $v$  relative to  $S$  is defined by

$$ipn(v, S) = \{u \in S : N(u) \cap S = \{v\}\}.$$

- The *external private neighbourhood* of  $v$  relative to  $S$  is defined by

$$epn(v, S) = \{u \in V(G) \setminus S : N(u) \cap S = \{v\}\}.$$

- The *private neighbourhood* of  $v$  relative to  $S$  is defined by

$$pn(v, S) = ipn(v, S) \cup epn(v, S) = \{u \in V(G) : N(u) \cap S = \{v\}\}.$$

The subgraph induced by  $S \subseteq V(G)$  will be denoted by  $\langle S \rangle$ , while the graph obtained from  $G$  by removing all the vertices in  $S \subseteq V(G)$  (and all the edges incident with a vertex in  $S$ ) will be denoted by  $G - S$ . If  $H$  is a graph, then we say that a graph  $G$  is  $H$ -free if  $G$  does not contain any copy of  $H$  as an induced subgraph.

We denote the set of leaves of a graph  $G$  by  $L(G)$ , and the set of support vertices (vertices adjacent to leaves) by  $S(G)$ . The set of isolated vertices of  $\langle V(G) \setminus (S(G) \cup L(G)) \rangle$  will be denoted by  $I_G$ .

We will use the notation  $C_n$ ,  $N_n$  and  $P_n$  for cycle graphs, empty graphs and path graphs of order  $n$ , respectively.

Let  $f : V(G) \rightarrow \{0, 1, 2\}$  be a function. For any  $i \in \{0, 1, 2\}$  we define the subsets of vertices  $V_i = \{v \in V(G) : f(v) = i\}$  and we identify  $f$  with the three subsets of  $V(G)$  induced by  $f$ . Thus, in order to emphasize the notation of these sets, we denote the function by  $f(V_0, V_1, V_2)$ . Given a set  $X \subseteq V(G)$ , we define  $f(X) = \sum_{v \in X} f(v)$ , and the *weight* of  $f$  is defined to be  $\omega(f) = f(V(G)) = |V_1| + 2|V_2|$ .

A (total) *weak Roman dominating function* is a function  $f(V_0, V_1, V_2)$  satisfying that  $V_1 \cup V_2$  is (total) dominating set and for every vertex  $v \in V_0$  there exists  $u \in N(v) \cap (V_1 \cup V_2)$  such that the function  $f'(V'_0, V'_1, V'_2)$ , defined by  $f'(v) = 1$ ,  $f'(u) = f(u) - 1$  and  $f'(x) = f(x)$  whenever  $x \in V(G) \setminus \{u, v\}$ ,

satisfies that  $V_1' \cup V_2'$  is (total) dominating set. Notice that  $S \subseteq V(G)$  is a secure (total) dominating set if and only if there exists a (total) weak Roman dominating function  $f(V_0, V_1, V_2)$  such that  $V_2 = \emptyset$  and  $V_1 = S$ .

The *weak Roman domination number*, denoted by  $\gamma_r(G)$ , is the minimum weight among all weak Roman dominating functions on  $G$ . By analogy we define the *total weak Roman domination number*, which is denoted by  $\gamma_{tr}(G)$ . The weak Roman domination number was introduced by Henning and Hedetniemi [15] and studied further in several works including [7,8,10,16,17], while the total weak Roman domination number was recently introduced in [12].

A dominating set of cardinality  $\gamma(G)$  will be called a  $\gamma(G)$ -set. A similar agreement will be assumed when referring to optimal sets associated with other parameters used in the article. As usual, we will use the acronyms TDS and STDS to refer to total dominating sets and secure total dominating sets, respectively.

A TDS  $X$  is said to be a *total outer-connected dominating set* if the subgraph induced by  $V(G) \setminus X$  is connected. The *total outer-connected domination number* of  $G$ , denoted by  $\gamma_{toc}(G)$ , is the minimum cardinality among all total outer-connected dominating sets of  $G$ . This parameter was introduced by Cyman in [18] and studied further in [19–21].

An *independent set* of a graph  $G$  is a subset of vertices such that no two vertices in the subset represent an edge of  $G$ . The maximum cardinality among all independent sets is the *independence number* of  $G$ , denoted by  $\alpha(G)$ . Analogously, two edges in a graph  $G$  are independent if they are not adjacent in  $G$ . A set of pairwise independent edges of  $G$  is called a *matching* of  $G$ . The *matching number*  $\alpha'(G)$ , sometimes known as the *edge independence number*, is the maximum cardinality among all matchings of  $G$ .

For the remainder of the paper, definitions will be introduced whenever a concept is needed.

### 3. Secure Total Domination & Independence

Klostermeyer and Mynhardt [9] in 2008, established the following upper bound.

**Theorem 1.** [9] For any graph  $G$  with no isolated vertex,

$$\gamma_{st}(G) \leq 3\alpha(G) - 1.$$

In 2017 Duginov [14] answered the following open question posed by Klostermeyer and Mynhardt [9] p. 282: Is there a graph  $G$  such that  $\gamma_{st}(G) = 3\alpha(G) - 1$ , where  $\alpha(G) \geq 2$ ? Duginov provided a negative answer to this question by confirming the suspicions of Klostermeyer and Mynhardt that  $\gamma_{st}(G) \leq 2\alpha(G)$ .

**Theorem 2.** [14] For any graph  $G$  with no isolated vertex,

$$\gamma_{st}(G) \leq 2\alpha(G).$$

We now proceed to improve the bound above.

**Lemma 1.** For any graph  $G$  and any set  $D \subseteq V(G)$ , there exists an  $\alpha(G)$ -set  $I$  such that for any  $x \in I$ ,  $ipn(x, D \cup I) = \emptyset$ .

**Proof.** Let  $I$  be an  $\alpha(G)$ -set,  $D \subseteq V(G)$  and  $D_I = \{x \in I : ipn(x, D \cup I) \neq \emptyset\}$ . We can assume that we have taken  $I$  in such a way that  $|D_I|$  is minimum among all  $\alpha(G)$ -sets. Suppose that there exists  $u \in D_I$ , and consequently, let  $v \in ipn(u, D \cup I)$ . Observe that  $v \in D \setminus I$  and  $I' = (I \cup \{v\}) \setminus \{u\}$  is an  $\alpha(G)$ -set. Since  $ipn(v, D \cup I') = \emptyset$  and  $ipn(x, D \cup I') = ipn(x, D \cup I)$  for every  $x \in I' \setminus \{v\}$ , we can conclude that  $I'$

is an  $\alpha(G)$ -set satisfying that  $|D_{I'}| < |D_I|$ , which is a contradiction. Therefore,  $D_I = \emptyset$ , which completes the proof.  $\square$

Since  $\gamma(G) \leq \alpha(G)$ , the following result improves Theorem 2.

**Theorem 3.** For any graph  $G$  with no isolated vertex,

$$\gamma_{st}(G) \leq \alpha(G) + \gamma(G).$$

**Proof.** Let  $D$  be a  $\gamma(G)$ -set. By Lemma 1 there exists an  $\alpha(G)$ -set  $I$  such that  $ipn(x, D \cup I) = \emptyset$  for every  $x \in I$ . We define the set  $S \subseteq V(G)$  as follows.

- (a)  $D \cup I \subseteq S$ .
- (b) For every vertex  $x \in D \cap I$ ,
  - if  $epn(x, D \cup I) \neq \emptyset$ , then choose one vertex  $u \in epn(x, D \cup I)$  and set  $u \in S$ .
  - if  $epn(x, D \cup I) = \emptyset$ , then choose one vertex  $u \in N(x) \setminus (D \cup I)$  (if any) and set  $u \in S$ .

It is readily seen that  $S$  is a TDS. Now, let  $v \in V(G) \setminus S$ . Since  $I \subseteq S$  is also a dominating set, there exists a vertex  $u \in N(v) \cap I \subseteq N(v) \cap S$ . To conclude that  $S$  is STDS, we only need to prove that  $S' = (S \setminus \{u\}) \cup \{v\}$  is a TDS. To this end, we differentiate two cases for any  $w \in N(u)$ .

Case 1.  $u \in I \setminus D$ . If  $w \notin D$ , then there exists some vertex in  $D \subseteq S'$  which dominates  $w$ , as  $D$  is a dominating set. If  $w \in D$ , then by Lemma 1 we have that  $w \notin ipn(u, D \cup I)$ . Hence, there exists some vertex in  $(D \cup I) \setminus \{u\} \subseteq S'$  which dominates  $w$ .

Case 2.  $u \in I \cap D$ . If  $w \in D$ , then by Lemma 1 we have that  $w \notin ipn(u, D \cup I)$ , and so there exists some vertex in  $(D \cup I) \setminus \{u\} \subseteq S'$  which dominates  $w$ . From now on, suppose that  $w \notin D$ . If  $w \notin epn(u, D \cup I)$ , then  $w$  is dominated by some vertex in  $(D \cup I) \setminus \{u\} \subseteq S'$ . If  $w \in epn(u, D \cup I)$  then, as all vertices in  $epn(u, D \cup I)$  form a clique and by (b) in the definition of  $S$ ,  $w$  is dominated by some vertex in  $S \setminus \{u\} \subseteq S'$ .

Now, since  $S$  is a TDS, we have that every vertex in  $V(G) \setminus N(u)$  is dominated by some vertex in  $S'$  and, according to the two cases above, we can conclude that  $S'$  is a TDS. Therefore,  $S$  is a STDS and so  $\gamma_{st}(G) \leq |S| \leq \alpha(G) + \gamma(G)$ , which completes the proof.  $\square$

The bound above is tight. For instance, it is achieved for any corona product graph  $G = H_1 \odot H_2$ , where  $H_1$  is an arbitrary graph and  $H_2$  is the disjoint union of  $k$  complete nontrivial graphs. Notice that  $\alpha(G) = k|V(H_1)|$ ,  $\gamma(G) = |V(H_1)|$  and  $\gamma_{st}(G) = (k+1)|V(H_1)| = \alpha(G) + \gamma(G)$ . Another example is the graph  $G$  shown in Figure 2, where  $\gamma_{st}(G) = 8$ ,  $\alpha(G) = 6$  and  $\gamma(G) = 2$ .

#### 4. Secure Total Domination & Other Kinds of Domination

For any graph  $G$  with no isolated vertex,  $V(G)$  is a secure total dominating set, which implies that  $\gamma_{st}(G) \leq |V(G)|$ . All graphs achieving this trivial bound were characterized by Benecke et al. as follows.

**Theorem 4.** [11] Let  $G$  be a graph of order  $n$ . Then  $\gamma_{st}(G) = n$  if and only if  $V(G) \setminus (L(G) \cup S(G))$  is an independent set.

Since every secure total dominating set is a total dominating set, it is clear that  $\gamma_t(G) \leq \gamma_{st}(G)$ . All graphs satisfying the equality were characterized by Klostermeyer and Mynhardt in [9].

**Theorem 5.** [9] If  $G$  is a connected graph, then the following statements are equivalent.

- $\gamma_{st}(G) = \gamma_t(G)$ .
- $\gamma_{st}(G) = 2$ .
- $G$  has two universal vertices.


The result above is an important tool to characterize all graphs with  $\gamma_{st}(G) = 3$ . To begin with, we need to state the following basic tool.

**Proposition 1.** *If  $H$  is a spanning subgraph (with no isolated vertex) of a graph  $G$ , then*

$$\gamma_{st}(G) \leq \gamma_{st}(H).$$

**Proof.** Let  $E^- = \{e_1, \dots, e_k\}$  be the set of all edges of  $G$  not belonging to the edge set of  $H$ . Let  $H_0 = G$  and, for every  $i \in \{1, \dots, k\}$ , let  $X_i = \{e_1, \dots, e_i\}$  and  $H_i = G - X_i$ . Since any STDS of  $H_i$  is a STDS of  $H_{i-1}$ , we can conclude that  $\gamma_{st}(H_{i-1}) \leq \gamma_{st}(H_i)$ . Hence,  $\gamma_{st}(G) = \gamma_{st}(H_0) \leq \gamma_{st}(H_1) \leq \dots \leq \gamma_{st}(H_k) = \gamma_{st}(H)$ .  $\square$

Let  $\mathcal{G}$  be the family of graphs  $H$  of order  $n \geq 3$  such that the subgraph induced by three vertices of  $H$  contains a path  $P_3$  and the remaining  $n - 3$  vertices have degree two and they form an independent set. Figure 1 shows a graph belonging to  $\mathcal{G}$ .



**Figure 1.** A graph  $H$  belonging to  $\mathcal{G}$ . The set of black-coloured vertices forms a  $\gamma_{st}(H)$ -set

**Theorem 6.** *Given a graph  $G$ , the following statements are equivalent.*

- $\gamma_{st}(G) = 3$ .
- $G$  has at most one universal vertex and there exists  $H \in \mathcal{G}$  which is a spanning subgraph of  $G$ .

**Proof.** Let  $D$  be a  $\gamma_{st}(G)$ -set and assume that  $|D| = 3$ . By Theorem 5,  $G$  has at most one universal vertex. Let  $D = \{u, v, w\}$  and notice that  $\langle D \rangle$  contains a path  $P_3$ , as  $D$  is a total dominating set of  $G$ . Since  $D$  is a STDS of  $G$ , we observe that  $|N(z) \cap D| \geq 2$  for every  $z \in V(G) \setminus D$ . Hence, in this case,  $G$  contains a spanning subgraph belonging to  $\mathcal{G}$ .


Conversely, since  $G$  has at most one universal vertex, by Theorem 5 we have that  $\gamma_{st}(G) \geq 3$ . Moreover, it is readily seen that  $\gamma_{st}(H) \leq 3$  for any  $H \in \mathcal{G}$ . Hence, if  $H \in \mathcal{G}$  is a spanning subgraph of  $G$ , by Proposition 1 it follows that  $\gamma_{st}(G) \leq 3$ . Therefore,  $\gamma_{st}(G) = 3$ .  $\square$

We now consider the relationship between  $\gamma_s(G)$  and  $\gamma_{st}(G)$ .

**Theorem 7.** [9] *Let  $G$  be a graph with no isolated vertex.*

- (i) *If  $\delta(G) = 1$ , then  $\gamma_s(G) + 1 \leq \gamma_{st}(G)$ .*
- (ii) *If  $\delta(G) \geq 2$ , then  $\gamma_s(G) \leq \gamma_{st}(G) \leq 2\gamma_s(G)$ .*

A natural question is if the bound  $\gamma_{st}(G) \leq 2\gamma_s(G)$ , due to Klostermeyer and Mynhardt, can be improved with  $\gamma_{st}(G) \leq \gamma_s(G) + \gamma(G)$ . The example given in Figure 2 shows that, in general, this inequality does not hold.



**Figure 2.** A graph  $G$  with  $\gamma_{st}(G) = 8$ ,  $\gamma_s(G) = 5$  and  $\gamma(G) = 2$ . The set of black-coloured vertices forms a  $\gamma_{st}(G)$ -set.

In Theorem 10 we will show some cases in which  $\gamma_{st}(G) \leq \gamma_s(G) + \gamma(G)$ . To this end, we need to outline the following two known results.

**Theorem 8.** [12] *The following inequalities hold for any graph  $G$  with no isolated vertex.*

- (i)  $\gamma_t(G) \leq \gamma_{tr}(G) \leq \gamma_{st}(G)$ .
- (ii)  $\gamma_{tr}(G) \leq \min\{\gamma_t(G), \gamma_r(G)\} + \gamma(G)$ .

Although the problem of characterizing all graphs with  $\gamma_{tr}(G) = \gamma_{st}(G)$  remains open, some particular cases were described in [12].

**Theorem 9.** [12]

- (i)  $\gamma_{st}(G) = \gamma(G) + 1$  if and only if  $\gamma_{tr}(G) = \gamma(G) + 1$ .
- (ii) For any  $\{K_{1,3}, K_{1,3} + e\}$ -free graph  $G$  with no isolated vertex,  $\gamma_{st}(G) = \gamma_{tr}(G)$ .
- (iii) For any graph  $G$  with no isolated vertex and maximum degree  $\Delta(G) \leq 2$ ,  $\gamma_{st}(G) = \gamma_{tr}(G)$ .

From Theorems 8 and 9 (ii), and using the fact that  $\gamma_r(G) \leq \gamma_s(G)$ , we can show that the bound  $\gamma_{st}(G) \leq 2\gamma_s(G)$  established in Theorem 7 can be improved for any  $\{K_{1,3}, K_{1,3} + e\}$ -free graph.

**Theorem 10.** For any  $\{K_{1,3}, K_{1,3} + e\}$ -free graph  $G$  with no isolated vertex,

$$\gamma_{st}(G) \leq \min\{\gamma_t(G), \gamma_r(G)\} + \gamma(G) \leq \gamma_s(G) + \gamma(G).$$

The previous bounds are tight. They are achieved, for instance, for the wheel graph  $G \cong N_1 + C_4$  and for  $G \cong N_2 + P_3$ , which is the join of  $N_2$  and  $P_3$ . For these two graphs we have that  $\gamma_{st}(G) = 3$ ,  $\gamma_s(G) = \gamma_r(G) = \gamma_t(G) = 2$  and  $\gamma(G) = 1$ .

To derive a consequence of Theorem 10 we need to state the following result due to Burger et al. [6].

**Theorem 11.** [6] For any connected graph  $G \not\cong C_5$  of order  $n$  and  $\delta(G) \geq 2$ ,

$$\gamma_s(G) \leq \left\lfloor \frac{n}{2} \right\rfloor.$$

Notice that  $\gamma_{st}(C_5) = 4 = \left\lfloor \frac{5}{2} \right\rfloor + \gamma(C_5)$ . Hence, from Theorems 10 and 11 we immediately have the next result.

**Theorem 12.** For any connected  $\{K_{1,3}, K_{1,3} + e\}$ -free graph  $G$  of order  $n$  and  $\delta(G) \geq 2$ ,

$$\gamma_{st}(G) \leq \left\lfloor \frac{n}{2} \right\rfloor + \gamma(G).$$

The bound above is tight. It is achieved for  $G \cong N_1 + C_4$ ,  $G \cong C_5$  and  $G \cong C_6$ , where  $\gamma_{st}(G)$  equals 3, 4 and 5, respectively.

The following result shows us a relationship between the secure total domination number and the total outer-connected domination number.

**Theorem 13.** *Let  $G$  be a graph of order  $n$ . If  $\gamma_{toc}(G) \leq n - 2$ , then*

$$\gamma_{st}(G) \leq \left\lfloor \frac{\gamma_{toc}(G) + n}{2} \right\rfloor.$$

**Proof.** We assume that  $\gamma_{toc}(G) \leq n - 2$ . Let  $D$  be a  $\gamma_{toc}(G)$ -set and  $S$  a  $\gamma(\langle V(G) \setminus D \rangle)$ -set. Since  $D$  is a TDS of  $G$ ,  $D \cup S$  is a TDS as well. Furthermore, every vertex  $u \in V(G) \setminus (D \cup S)$  is dominated by some vertex  $v \in S$ , and  $D \subseteq (D \cup S \cup \{u\}) \setminus \{v\}$  is a TDS of  $G$ . Hence,  $D \cup S$  is a STDS of  $G$ , which implies that  $\gamma_{st}(G) \leq |D \cup S| = |D| + |S|$ . Now, since  $\langle V(G) \setminus D \rangle$  is a connected nontrivial graph, we have that  $|S| = \gamma(\langle V(G) \setminus D \rangle) \leq \frac{|V(G) \setminus D|}{2} = \frac{n - \gamma_{toc}(G)}{2}$ . Therefore,  $\gamma_{st}(G) \leq \lfloor \frac{\gamma_{toc}(G) + n}{2} \rfloor$ , which completes the proof.  $\square$

The bound above is tight. For instance, it is achieved for the wheel graph  $G \cong N_1 + C_4$  and for  $G \cong N_2 + P_3$ . In both cases  $\gamma_{st}(G) = 3$  and  $\gamma_{toc}(G) = 2$ .

The following result was obtained by Favaron et al. in [20].

**Theorem 14.** [20] *For any graph  $G$  of order  $n$ , diameter  $diam(G) \leq 2$  and minimum degree  $\delta(G) \geq 3$ ,*

$$\gamma_{toc}(G) \leq \left\lfloor \frac{2n - 2}{3} \right\rfloor.$$

The following result is a direct consequence of combining the result above and Theorem 13.

**Theorem 15.** *For any graph  $G$  of order  $n$ , diameter two and minimum degree  $\delta(G) \geq 3$ ,*

$$\gamma_{st}(G) \leq \left\lfloor \frac{5n - 2}{6} \right\rfloor.$$

The bound above is achieved for the wheel graph  $G \cong N_1 + C_4$  and for  $G \cong N_2 + P_3$ . As we already know, in both cases  $\gamma_{st}(G) = 3$ .

### 5. Secure Total Domination & Matching

To begin this section, we proceed to introduce new definitions and terminology. Given a matching  $\mathcal{M}$  of a graph  $G$ , let  $V_{\mathcal{M}}$  be the set formed by the end-vertices of edges belonging to  $\mathcal{M}$ . Given a vertex  $v \in V_{\mathcal{M}}$ , we say that  $v' \in V_{\mathcal{M}}$  is the *partner* of  $v$  if  $vv' \in \mathcal{M}$ . Observe that if  $v'$  is the partner of  $v$ , then  $v$  is the partner of  $v'$ .

A *maximum matching* is a matching of cardinality  $\alpha'(G)$ . The following lemmas show some properties of maximum matchings.

**Lemma 2.** *Let  $\mathcal{M}$  be a maximum matching of a graph  $G$ . The following statements hold.*

- (i)  $N(u) \subseteq V_{\mathcal{M}}$  for every  $u \in V(G) \setminus V_{\mathcal{M}}$ .
- (ii) If  $u \in V(G) \setminus V_{\mathcal{M}}$  is adjacent to  $v \in V_{\mathcal{M}}$ , then  $N(v') \subseteq V_{\mathcal{M}} \cup \{u\}$ , where  $v'$  is the partner of  $v$ .

**Proof.** Let  $u \in V(G) \setminus V_{\mathcal{M}}$ . If there exists a vertex  $w \in N(u) \cap (V(G) \setminus V_{\mathcal{M}})$ , then the set  $\mathcal{M} \cup \{uw\}$  is a matching of  $G$  of cardinality greater than  $|\mathcal{M}|$ , which is a contradiction. Hence,  $N(u) \subseteq V_{\mathcal{M}}$  and (i) follows.

Now, we suppose that there exists  $u \in V(G) \setminus V_{\mathcal{M}}$  and a vertex  $v \in N(u) \cap V_{\mathcal{M}}$ . Let  $v'$  be the partner of  $v$ . If there exists a vertex  $w \in N(v') \cap (V(G) \setminus (V_{\mathcal{M}} \cup \{u\}))$ , then the set  $\mathcal{M} \setminus \{vv'\} \cup \{uv, v'w\}$  is a matching of  $G$  of cardinality greater than  $|\mathcal{M}|$ , which is a contradiction. Hence,  $N(v') \subseteq V_{\mathcal{M}} \cup \{u\}$  and (ii) follows.  $\square$

**Lemma 3.** For any graph  $G$  with  $L(G) \neq \emptyset$ , there exists a maximum matching  $\mathcal{M}$  such that for each vertex  $x \in S(G)$  there exists  $y \in L(G)$  such that  $xy \in \mathcal{M}$ .

**Proof.** Let  $\mathcal{M}$  be a maximum matching of  $G$  such that  $|V_{\mathcal{M}} \cap L(G)|$  is maximum. It is easy to see that the maximality of  $\mathcal{M}$  leads to  $S(G) \subseteq V_{\mathcal{M}}$ . Suppose that there exists a support vertex  $x$  such that  $xx' \in \mathcal{M}$  and  $x' \notin L(G)$ . Let  $y \in N(x) \cap L(G)$ . Notice that the set  $\mathcal{M}' = \mathcal{M} \setminus \{xx'\} \cup \{xy\}$  is a maximum matching of  $G$  and  $|V_{\mathcal{M}'} \cap L(G)| > |V_{\mathcal{M}} \cap L(G)|$ , which is a contradiction. Therefore, the result follows.  $\square$

The next result provides a relationship between the secure total domination number, the matching number and some special vertices of a graph.

**Theorem 16.** For any graph  $G$  with minimum degree  $\delta(G) = 1$ ,

$$\gamma_{st}(G) \leq 2\alpha'(G) + |L(G)| - |S(G)| + |I_G|.$$

**Proof.** Let  $\mathcal{M}$  be a maximum matching satisfying Lemma 3. Let  $S = V_{\mathcal{M}} \cup L(G) \cup I_G$ . Notice that  $V_{\mathcal{M}} \cap I_G = \emptyset$  and  $S(G) \subseteq V_{\mathcal{M}}$ . Hence,  $|S| = 2\alpha'(G) + |L(G)| - |S(G)| + |I_G|$ .

Notice that  $S$  is a TDS of  $G$ . We shall show that  $S$  is a STDS of  $G$ . Now, let  $v \in V(G) \setminus S$ . Since  $v \notin I_G$  and  $V_{\mathcal{M}}$  is a dominating set of  $G$ , there exists a vertex  $u \in V_{\mathcal{M}} \setminus S(G)$  which is adjacent to  $v$ . Let  $S' = (S \setminus \{u\}) \cup \{v\}$ . We will see that  $S'$  is a TDS of  $G$  as well. Since  $S$  is a TDS of  $G$ , every vertex  $w \in V(G) \setminus N(u)$  is adjacent to some vertex belonging to  $S'$ . Let  $w \in N(u)$  and observe that  $|N(w)| \geq 2$  as  $u \notin S(G)$ .

If  $w \in V(G) \setminus V_{\mathcal{M}}$ , then by Lemma 2 (i) we have that  $N(w) \subseteq V_{\mathcal{M}}$ . Hence there exists a vertex in  $V_{\mathcal{M}} \setminus \{u\} \subseteq S'$  which is adjacent to  $w$ , as  $|N(w)| \geq 2$ . Now, if  $w \in V_{\mathcal{M}} \setminus \{u'\}$ , where  $u'$  is the partner of  $u$ , then  $w$  is adjacent to its partner, which belongs to  $S'$ . Finally, if  $w = u'$ , then by Lemma 2 (ii) we have that  $N(w) \subseteq V_{\mathcal{M}} \cup \{v\}$  and since  $|N(w)| \geq 2$  it follows that  $N(w) \subseteq (V_{\mathcal{M}} \setminus \{u\}) \cup \{v\} \subseteq S'$ .

Thus,  $S'$  is a TDS of  $G$ , as desired. Therefore,  $S$  is a STDS and so  $\gamma_{st}(G) \leq |S| = 2\alpha'(G) + |L(G)| - |S(G)| + |I_G|$ .  $\square$

The bound above is tight. For instance, it is achieved for the graph shown in Figure 3. In this case,  $\gamma_{st}(G) = 22$ ,  $\alpha'(G) = 7$ ,  $|L(G)| = 12$ ,  $|S(G)| = 6$  and  $|I_G| = 2$ .




Figure 3. The set of black-coloured vertices forms a  $\gamma_{st}(G)$ -set.

From now on we consider the case of graphs with minimum degree  $\delta(G) \geq 2$ .



**Definition 1.** Given a maximum matching  $\mathcal{M}$  of a graph  $G$  with  $\delta(G) \geq 2$ , we construct a set  $D_{\mathcal{M}} \subseteq V_{\mathcal{M}}$  as follows.

- (i)  $|D_{\mathcal{M}}| = \alpha'(G)$ .
- (ii)  $xy \notin \mathcal{M}$  for all  $x, y \in D_{\mathcal{M}}$ .
- (iii)  $|N(x) \cap (V(G) \setminus V_{\mathcal{M}})| \geq |N(x') \cap (V(G) \setminus V_{\mathcal{M}})|$  for all  $x \in D_{\mathcal{M}}$ , where  $x'$  is the partner of  $x$ .

We proceed to show some properties of  $D_{\mathcal{M}} \subseteq V_{\mathcal{M}}$ .

**Lemma 4.** Let  $\mathcal{M}$  be a maximum matching of a graph  $G$  with  $\delta(G) \geq 2$ . The following statements hold.

- (a) If  $u \in V(G) \setminus V_{\mathcal{M}}$  is adjacent to  $v' \in V_{\mathcal{M}} \setminus D_{\mathcal{M}}$ , then  $u$  is adjacent to  $v \in D_{\mathcal{M}}$ , where  $v$  is the partner of  $v'$ .
- (b)  $D_{\mathcal{M}}$  is a dominating set of  $G$ .
- (c) If  $v \in D_{\mathcal{M}}$ , then its partner  $v' \in V_{\mathcal{M}} \setminus D_{\mathcal{M}}$  satisfies that  $|N(v') \cap V_{\mathcal{M}}| \geq \delta(G) - 1$ .

**Proof.** Let  $u \in V(G) \setminus V_{\mathcal{M}}$ . By Lemma 2 (i) we have that  $N(u) \subseteq V_{\mathcal{M}}$ . If there exists a vertex  $v' \in V_{\mathcal{M}} \setminus D_{\mathcal{M}}$ , then by Lemma 2 (ii) we have that  $N(v) \subseteq V_{\mathcal{M}} \cup \{u\}$  (where  $v \in D_{\mathcal{M}}$  is the partner of  $v'$ ). By item (iii) in the definition of  $D_{\mathcal{M}}$  it follows that  $u \in N(v)$  and (a) holds.

From item (a) we deduce that  $N(u) \cap D_{\mathcal{M}} \neq \emptyset$ . Now, by definition of  $D_{\mathcal{M}}$ , every vertex in  $V_{\mathcal{M}} \setminus D_{\mathcal{M}}$  is dominated by its partner, which belongs to  $D_{\mathcal{M}}$ . Therefore,  $D_{\mathcal{M}}$  is a dominating set of  $G$  and so (b) follows.

Now, let  $z \in D_{\mathcal{M}}$  and  $z'$  its partner. If  $|N(z') \cap V_{\mathcal{M}}| \leq \delta(G) - 2$ , then there exist two vertices  $x, y \in N(z') \cap (V(G) \setminus V_{\mathcal{M}})$ . By Lemma 4 (a) we have that  $x, y \in N(z)$ , which is a contradiction by Lemma 2 (ii). Therefore,  $|N(z') \cap V_{\mathcal{M}}| \geq \delta(G) - 1$  and (c) follows, which completes the proof.  $\square$

**Theorem 17.** For any graph  $G$  with minimum degree  $\delta(G) \geq 2$ ,

$$\gamma_{st}(G) \leq 2\alpha'(G) - \delta(G) + 2.$$

**Proof.** Let  $n$  be the order of  $G$ . Let  $v \in V(G)$  be a vertex of degree  $\delta(G)$  and  $u \in N(v)$ . It is readily seen that the set  $S = (V(G) \setminus N(v)) \cup \{u\}$  is a STDS of  $G$  and, as a consequence,  $\gamma_{st}(G) \leq n - \delta(G) + 1$ . Thus, the inequality holds for  $2\alpha'(G) \in \{n - 1, n\}$ .

From now on we suppose that  $2\alpha'(G) \leq n - 2$ . Let  $\mathcal{M}$  be a maximum matching of  $G$ . Since  $|V_{\mathcal{M}}| = 2\alpha'(G) \leq n - 2$ , there exist two vertices  $x, y \in V(G) \setminus V_{\mathcal{M}}$ . By Lemma 4 (b) we have that  $D_{\mathcal{M}}$  is a dominating set of  $G$ , which implies that there exists a vertex  $v_x \in N(x) \cap D_{\mathcal{M}}$ . Since  $\delta(G) \geq 2$ , by Lemmas 2 and 4 (a), there exists a vertex  $v_y \in N(y) \cap (D_{\mathcal{M}} \setminus \{v_x\})$  and also we deduce that  $N(x) \cup N(y) \subseteq V_{\mathcal{M}}$  and  $N(x) \cap N(y) \subseteq D_{\mathcal{M}}$ . Let  $R = (N(x) \cup N(y)) \cap D_{\mathcal{M}}$ . Hence  $|R| = |N(x) \cap N(y)| + |(N(x) \setminus N(y)) \cap D_{\mathcal{M}}| + |(N(y) \setminus N(x)) \cap D_{\mathcal{M}}| \geq (|N(x)| + |N(y)|)/2 \geq \delta(G)$ . Let  $Z \subseteq R \setminus \{v_x, v_y\}$  such that  $|Z| = \delta(G) - 2$  and let  $Z'$  be the set of partners of the vertices in  $Z$ .

Let  $\mathcal{M}' = (\mathcal{M} \setminus \{v_x v'_x, v_y v'_y\}) \cup \{xv_x, yv_y\}$ , where  $v'_x$  and  $v'_y$  are the partners of  $v_x$  and  $v_y$  respectively. Notice that  $\mathcal{M}'$  is a maximum matching of  $G$  and the set  $D_{\mathcal{M}'} = D_{\mathcal{M}} \subseteq V_{\mathcal{M}'}$  satisfies the conditions given in Definition 1.

We will prove that  $S = V_{\mathcal{M}'} \setminus Z'$  is a STDS of  $G$ . By Lemma 4 (b) we have that  $D_{\mathcal{M}}$  is a dominating set of  $G$ , which implies that every vertex in  $V(G) \setminus S$  is dominated by some vertex in  $D_{\mathcal{M}} \subseteq S$ . Also, every vertex in  $Z$  is dominated by either  $x$  or  $y$ , which belong to  $S$ , and every vertex in  $S \setminus Z$  satisfies that its partner belongs to  $S$  as well. Hence  $S$  is a TDS of  $G$ .

Let  $v \in V(G) \setminus S$  and let  $S' = (S \setminus \{v^*\}) \cup \{v\}$ , where either  $v^* = v'$  is the partner of  $v$  if  $v \in Z'$ , or  $v^*$  is a vertex belonging to  $N(v) \cap D_{\mathcal{M}}$  if  $v \in V(G) \setminus V_{\mathcal{M}'}$  (notice that in this case,  $v^*$  exists since  $D_{\mathcal{M}}$

is a dominating set). We only need to prove that  $S'$  is a TDS of  $G$ . Since  $S$  is a TDS of  $G$ , every vertex in  $V(G) \setminus N(v^*)$  has at least one neighbour in  $S'$ . Now, let  $u \in N(v^*)$  and consider the following two cases.

Case 1.  $u \in V(G) \setminus V_{\mathcal{M}'}$ . Since  $|V_{\mathcal{M}'} \cap (V(G) \setminus S')| = \delta(G) - 1$ , by Lemma 2 (i) we deduce that there exists some vertex in  $N(u) \cap S'$ .

Case 2.  $u \in V_{\mathcal{M}'}$ . In this case, we analyse three subcases. If  $u \in Z$ , then  $u$  is dominated by either  $x$  or  $y$ , which belong to  $S'$ . If  $u = v$ , then as  $u \in V_{\mathcal{M}'} \setminus D_{\mathcal{M}'}$ , by Lemma 4 (c) it follows that  $|N(u) \cap V_{\mathcal{M}'}| \geq \delta(G) - 1$ . As in this case  $|V_{\mathcal{M}'} \cap (V(G) \setminus S')| = \delta(G) - 2$ , we deduce that  $N(u) \cap S' \neq \emptyset$ . Finally, if  $u \in V_{\mathcal{M}'} \setminus (Z \cup \{v\})$ , then its partner belongs to  $S'$ .

Hence,  $S'$  is a TDS of  $G$ , as desired. Therefore,  $S$  is a STDS of  $G$  and  $\gamma_{st}(G) \leq |S| = |V_{\mathcal{M}'} \setminus Z'| = 2\alpha'(G) - \delta(G) + 2$ , which completes the proof.  $\square$

The bound above is tight. For instance, it is achieved for the graphs  $G \cong N_2 + P_3$  and  $G \cong N_1 + C_4$ . In both cases  $\gamma_{st}(G) = 3$ ,  $\alpha'(G) = 2$  and  $\delta(G) = 3$ .

Cockayne et al. in [8] obtained the following bound on the secure domination number in terms of the order and the matching number.

**Theorem 18.** [8] *If a graph  $G$  of order  $n$  does not have isolated vertices, then*

$$\gamma_s(G) \leq n - \alpha'(G).$$

Therefore, by Theorems 10 and 18 we deduce the following result.

**Theorem 19.** *For any  $\{K_{1,3}, K_{1,3} + e\}$ -free graph  $G$  with minimum degree  $\delta(G) \geq 1$  and order  $n$ ,*

$$\gamma_{st}(G) \leq n - \alpha'(G) + \gamma(G).$$

The bound above is tight. For instance, it is achieved for the graphs  $G \cong C_6$  and  $G \cong P_6$ , as for these graphs we have  $\gamma_{st}(G) = 5$ ,  $\alpha'(G) = 3$  and  $\gamma(G) = 2$ .

The  $k$ -domination number of  $G$ , denoted by  $\gamma_k(G)$ , is another well-known parameter [3]. The following theorem is a contribution of DeLaViña et al. in [22].

**Theorem 20.** [22] *Let  $k$  be a positive integer. For any graph  $G$  with minimum degree  $\delta(G) \geq 2k - 1$ ,*

$$\gamma_k(G) \leq \alpha'(G).$$

Since every  $\gamma_2(G)$ -set is a secure dominating set of  $G$ , it is immediate that  $\gamma_s(G) \leq \gamma_2(G)$ , and so Theorems 10 and 20 lead to the following result.

**Theorem 21.** *For any  $\{K_{1,3}, K_{1,3} + e\}$ -free graph  $G$  with minimum degree  $\delta(G) \geq 3$ ,*

$$\gamma_{st}(G) \leq \alpha'(G) + \gamma(G).$$

The bound above is tight. For instance, it is achieved for the wheel graph  $G \cong N_1 + C_4$  and for  $G \cong N_2 + P_3$ , as in both cases  $\gamma_{st}(G) = 3$ ,  $\alpha'(G) = 2$  and  $\gamma(G) = 1$ .

## 6. Conclusions

This article is a contribution to the theory of protection of graphs. In particular, it is devoted to the study of the secure total domination number of a graph. We study the properties of this parameter in order to obtain its exact value or general bounds. Among our main contributions we highlight the following.

- We show that  $\gamma_{st}(G) \leq \alpha(G) + \gamma(G)$ . Since  $\gamma(G) \leq \alpha(G)$ , this result improves the bound  $\gamma_{st}(G) \leq 2\alpha(G)$  obtained in [14].
- We characterize the graphs with  $\gamma_{st}(G) = 3$ .
- We show that if  $G$  is a  $\{K_{1,3}, K_{1,3} + e\}$ -free graph  $G$  with no isolated vertex, then  $\gamma_{st}(G) \leq \min\{\gamma_t(G), \gamma_r(G)\} + \gamma(G) \leq \gamma_s(G) + \gamma(G)$ .
- We study the relationship that exists between the secure total domination number and the matching number of a graph. In particular, we obtain the following results.
  - (a)  $\gamma_{st}(G) \leq 2\alpha'(G) + |L(G)| - |S(G)| + |I_G|$  for any graph  $G$  of minimum degree one.
  - (b)  $\gamma_{st}(G) \leq 2\alpha'(G) - \delta(G) + 2$  for every graph  $G$  of minimum degree  $\delta(G) \geq 2$ .
  - (c)  $\gamma_{st}(G) \leq \alpha'(G) + \gamma(G)$  for every  $\{K_{1,3}, K_{1,3} + e\}$ -free graph  $G$  of minimum degree  $\delta(G) \geq 3$ .

All bounds obtained in the paper are tight.

**Author Contributions:** The results presented in this paper were obtained as a result of collective work sessions involving all authors. The process was organized and led by J.A.R.-V.

**Funding:** This work has been partially supported by the Spanish Ministry of Economy, Industry and Competitiveness, under the grants MTM2016-78227-C2-1-P and MTM2017-90584-REDT.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Cockayne, E.J.; Grobler, P.J.P.; Gründlingh, W.R.; Munganga, J.; van Vuuren, J.H. Protection of a graph. *Util. Math.* **2005**, *67*, 19–32.
2. Haynes, T.W.; Hedetniemi, S.T.; Slater, P.J. *Domination in Graphs: Volume 2: Advanced Topics*; Chapman & Hall/CRC Pure and Applied Mathematics; Taylor & Francis Group: Abingdon, UK, 1998.
3. Haynes, T.W.; Hedetniemi, S.T.; Slater, P.J. *Fundamentals of Domination in Graphs*; Chapman and Hall/CRC Pure and Applied Mathematics Series; Marcel Dekker, Inc.: New York, NY, USA, 1998.
4. Henning, M.A.; Yeo, A. *Total Domination in Graphs, Springer Monographs in Mathematics*; Springer: New York, NY, USA, 2013.
5. Boumediene Merouane, H.; Chellali, M. On secure domination in graphs. *Inform. Process. Lett.* **2015**, *115*, 786–790. [[CrossRef](#)]
6. Burger, A.P.; Henning, M.A.; van Vuuren, J.H. Vertex covers and secure domination in graphs. *Quaest. Math.* **2008**, *31*, 163–171. [[CrossRef](#)]
7. Chellali, M.; Haynes, T.W.; Hedetniemi, S.T. Bounds on weak Roman and 2-rainbow domination numbers. *Discrete Appl. Math.* **2014**, *178*, 27–32. [[CrossRef](#)]
8. Cockayne, E.J.; Favaron, O.; Mynhardt, C.M. Secure domination, weak Roman domination and forbidden subgraphs. *Bull. Inst. Combin. Appl.* **2003**, *39*, 87–100.
9. Klostermeyer, W.F.; Mynhardt, C.M. Secure domination and secure total domination in graphs. *Discuss. Math. Graph Theory* **2008**, *28*, 267–284. [[CrossRef](#)]
10. Valveny, M.; Rodríguez-Velázquez, J.A. Protection of graphs with emphasis on Cartesian product graphs. *Filomat* **2019**, *33*, 319–333.
11. Benecke, S.; Cockayne, E.J.; Mynhardt, C.M. Secure total domination in graphs. *Util. Math.* **2007**, *74*, 247–259.
12. Cabrera Martínez, A.; Montejano, L.P.; Rodríguez-Velázquez, J.A. Total weak Roman domination in graphs. *Symmetry* **2019**, *11*, 831. [[CrossRef](#)]

13. Kulli, V.R.; Chaluvvaraju, B.; Kumara, M. Graphs with equal secure total domination and inverse secure total domination numbers. *J. Inf. Optim. Sci.* **2008**, *39*, 467–473. [[CrossRef](#)]
14. Duginov, O. Secure total domination in graphs: bounds and complexity. *Discrete Appl. Math.* **2017**, *222*, 97–108. [[CrossRef](#)]
15. Henning, M.A.; Hedetniemi, S.T. Defending the Roman Empire—A new strategy. *Discrete Math.* **2003**, *266*, 239–251. [[CrossRef](#)]
16. Pushpam, P.R.L.; Malini Mai, T.N.M. Weak Roman domination in graphs. *Discuss. Math. Graph Theory* **2011**, *31*, 115–128.
17. Valveny, M.; Pérez-Rosés, H.; Rodríguez-Velázquez, J.A. On the weak Roman domination number of lexicographic product graphs. *Discrete Appl. Math.* **2019**, *263*, 257–270. [[CrossRef](#)]
18. Cyman, J. Total outer-connected domination numbers in trees. *Discuss. Math. Graph Theory* **2010**, *30*, 377–383. [[CrossRef](#)]
19. Cyman, J.; Raczek, J. Total outer-connected domination numbers of trees. *Discrete Appl. Math.* **2009**, *157*, 3198–3202. [[CrossRef](#)]
20. Favaron, O.; Karami, H.; Sheikholeslami, S.M. On the total outer-connected domination in graphs. *J. Comb. Optim.* **2014**, *27*, 451–461. [[CrossRef](#)]
21. Hattingh, J.; Joubert, E. A note on total outer-connected domination numbers of a tree. *AKCE J. Graphs Comb.* **2010**, *7*, 223–227.
22. DeLaViña, E.; Goddard, W.; Henning, M.A.; Pepper, R.; Vaughan, E.R. Bounds on the  $k$ -domination number of a graph. *Appl. Math. Lett.* **2011**, *24*, 996–998. [[CrossRef](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).