

Development of Predicted Environmental Concentrations to prioritize the occurrence of pharmaceuticals in rivers from Catalonia.

Cristian Gómez-Canela^{1,2*}, Víctor Pueyo², Carlos Barata², Sílvia Lacorte², Rosa Maria Marcé¹

¹Department of Analytical Chemistry and Organic Chemistry, Campus Sescelades, Faculty of Chemistry, Universitat Rovira i Virgili, Marcel·lí Domingo s/n, Tarragona, Catalonia, Spain.

²Department of Environmental Chemistry, IDAEA-CSIC. Jordi Girona 18-26, 08034 Barcelona, Catalonia, Spain.

*Corresponding author:

Email: cristian.gomez@urv.cat (C. Gómez-Canela)

Tel: +34 93 400 61 00 / Fax: +34 93 204 59 04

Abstract

The main objective of the present study is to prioritize those pharmaceuticals that have higher chances to be detected in water due to incomplete removal in Wastewater Treatment Plants (WWTPs). To do so, the total consumption of pharmaceuticals in Catalonia (NE Spain) were compiled to calculate the predicted environmental concentrations (PECs) in wastewater effluents and in river water. PECs were estimated using publicly available consumption data in the period of 2013-2016 for a suite of 165 compounds. The selected compounds were based on generic pharmaceuticals with emphasis on drugs consumed by people aged 65 or over as they represent the age group with the highest consumption of pharmaceuticals. The mean total consumption of pharmaceuticals in the period studied was of 623 ± 3 t per year. Paracetamol, metformin and ibuprofen were the most administered drugs although the highest PEC values corresponded to metformin, amoxicillin and metamizole. Finally, predicted environmental levels together with acute and chronic toxicological data allowed estimating the risks of these compounds. Amoxicillin is expected to pose adverse effects for cyanobacteria, whereas metformin and ibuprofen pose a small potential for adverse effects to invertebrates and fish, respectively.

Keywords: Pharmaceuticals; prioritization; PECs; rivers; risk assessment.

1. Introduction

Nowadays, water pollution represents one of the most serious ecological threats we face. Hence, quality of water has to be preserved to protect the environment and human health from compounds capable of exerting an effect at low levels of concentration. Among other chemicals, pharmaceuticals represent nowadays a relevant class of contaminants because they are continuously released into the aquatic environment and are considered as 'pseudo-persistent' pollutants (Daughton, 2003; Kümmerer, 2009). The first findings of pharmaceuticals in the aquatic environment were reported in the seventies, where the presence of drugs and drug metabolites were detected in sewage water effluent (Hignite and Azarnoff, 1977). Since then, the recurrent and global occurrence of pharmaceuticals in wastewater treatment plants (WWTPs) and surface waters (river, seas, lakes) is reported in the range of ng L^{-1} and $\mu\text{g L}^{-1}$ (Hernando et al., 2006; Kümmerer, 2001; Navarro-Ortega et al., 2012a; Navarro-Ortega et al., 2012b; Ternes et al., 2001). Human consumption, animal use, waste disposal and/or manufacturing are the main sources of pharmaceuticals to the environment (Daouk et al., 2015).

Regarding human consumption, the drugbank has 11901 drugs in its database (Wishart et al., 2018) and the European Medical Agency (EMA) compiles around 2500 active ingredients. Consumption of pharmaceuticals by the global population vary among countries and its use is expected to grow as the population ages and upon polimedication. Virtually every country in the world is experiencing growth in the number and proportion of the elder population. In 2017, 962 million people aged 60 or over were estimated at a global scale, comprising 13 per cent of the global population, and raising to 25% in Europe. This age group is growing at a rate of about 3 per cent per year (United Nations (UN), 2015).. Elder population in developed countries have a high consumption of pharmaceuticals (Valderrama Gama et al., 1998), Typical consumption of 5-10 pills/patient/day in residents of senior residences which is translated in a total consumption of pharmaceuticals of hundreds of milligrams per day/person (Lacorte et al., 2017).

Because monitoring of all pharmaceuticals consumed is practically impossible in terms of time and cost, prioritization tools are needed for compounds that deem to pose negative effects in the environment. Predicted environmental concentrations (PECs) is a practical way (Carballa et al., 2008; Franquet-Griell et al., 2015), to estimate the concentrations of drugs expected to be found in the environment based on consumption data. Its use has been reported before for cytostatic (Franquet-Griell et al., 2015; Franquet-Griell et al., 2017) and other drugs (Burns et al., 2018; Guo et al., 2016; Verlicchi et al., 2014) in wastewater and surface waters.

On the other hand, many pharmaceuticals have been reported to be acutely and chronically toxic to aquatic biota with effects varying largely across taxa and among chemical groups (Crane et al., 2006; Fent et al., 2006a; Overturf et al., 2015; Owen et al., 2007; Santos et al., 2010). Acute toxicity values (immobilization or mortality, algae growth within 48-96 h and the half-maximum effective concentration or lethal concentration, EC_{50}/LC_{50} , value) of pharmaceuticals to algae, aquatic plants, *Daphnia magna* and fish species have been generally reported in the order of $mg\ L^{-1}$. Moreover, in terms of chronic toxicity, most individual pharmaceuticals (such as antibiotics, antibacterial drugs, anti-inflammatory drugs, non-steroidal anti-inflammatory drugs) induce reproductive toxicity in fish and crustaceans at low or environmentally relevant concentrations ($ng\ L^{-1}$ - $\mu g\ L^{-1}$), (Crane et al., 2006; Fent et al., 2006a; Overturf et al., 2015; Owen et al., 2007; Santos et al., 2010). In addition, the combination of different drugs sharing a common mechanism of action could produce additive effects that would be enough to enhance toxicity (Cleuvers, 2004; Fent et al., 2006b).

Taking into account the large amount and diversity of pharmaceuticals consumed and discharged to the environment, the aim of this study was to prioritize and evaluate their environmental risk by calculating PECs according to the consumption in hospitals and pharmacies from 2013 to 2016. The procedure has been optimized considering Catalonia, a region of 7.5 million inhabitants, with an elderly population (≥ 65 years) of 24.5%. This region suffers from water scarcity and thus, the levels and potential toxic effects of pharmaceuticals can be high. PECs were calculated in 11 river basins according to specific population

distribution and dilution factor (DF). In addition, the potential risk for the aquatic environment was studied considering acute and chronic toxicity of different aquatic taxa for selected pharmaceuticals with highest PEC_{river} .

2. Material and methods

2.1. Consumption data

Consumption data, being the term “consumption” the dispensation or sales of pharmaceuticals, were obtained from Catalan Health Service (CatSalut) over the period 2013–2016 in pharmacies and hospitals. Compound selection was based on medical prescriptions for the elderly (greater than 65 years) in Catalonia in the last years (<http://observatorisalut.gencat.cat>) (Generalitat de Catalunya - Catalan Health System Observatory, 2018). A total of 165 drugs belonging to 13 different Anatomical Therapeutic Classification (ATC) groups (A, B, C, D, G, H, J, L, M, N, R, S and V) were included. More than 60% of the pharmaceuticals presented in this study are included in the top 200 worldwide sales and prescriptions realized by Njardarson et al. (McGrath et al., 2010; Njardarson, 2010). Table SI1 (Supplementary Information) shows the ATC and the most relevant physico-chemical properties of target compounds. This information does not include consumption through mutual insurance companies, as this data was not available for the public health system. Hospital data was given as the number of pills, capsules or any formulation of each active pharmaceutical ingredient dispensed (called activities). From the composition and the number of activities of each pharmaceutical, the total consumption (in $kg\ year^{-1}$) was calculated (Table SI2). The consumptions of the pharmaceuticals in pharmacies were directly provided in $mg\ yr^{-1}$ by the “Pharmaceutical and Medication Management” of CatSalut.

2.2. PEC calculations (effluent, river and basins)

To calculate the predicted environmental concentrations in WWTP effluents (PEC_{eff}) and surface waters (PEC_{river}) in $\mu g\ L^{-1}$, the following equations adapted from Besse et al. (Besse et al., 2008) were used:

$$PEC_{eff} \left(\frac{\mu g}{L} \right) = \frac{consumption \times F_{exc} \times (1 - F_{wwtp})}{W \times inhab} \times 10^9 \quad (1)$$

$$PEC_{river} \left(\frac{\mu g}{L} \right) = \frac{consumption \times F_{exc} \times (1 - F_{wwtp})}{W \times inhab \times DF} \times 10^9 \quad (2)$$

Consumption (g day⁻¹) is the quantity of each pharmaceutical consumed in Catalonia; *F_{exc}* is the excreted fraction of the unchanged drug, considering both urine and feces. Values were extracted from the Drugbank database (Wishart et al., 2018). For those drugs whose values could not be found, a default value of 0.5 was applied, considering that a pharmaceutical will not be totally excreted as parental compound; *F_{wwtp}* is the removal fraction in WWTP. Here, when no data was obtained from the bibliography, a default value of 0.5 was used. Then, *1 - F_{wwtp}*, is the fraction of pharmaceutical's emission from WWTPs to surface waters; *W* (L inhab⁻¹ d⁻¹) is the mean water consumption per person per day (about 130.9 L inhab⁻¹ d⁻¹ in Catalonia). This value was calculated from the mean total urban water consumption in Catalonia in the period studied and the number of inhabitants (IDESCAT, 2017). *Inhab* is the number of inhabitants in a defined zone (in Catalonia, a mean value of 7,446,487 during the period of 2013-2016). In the calculation of PECs, there are several uncertainties which are related to the input parameters of the PEC formula. These uncertainties can produce a bias in the PEC calculation and can alter the results obtained, and thus, their applicability. Among all parameters used in the PEC calculation, the two most significant in the variability of the results are: *F_{exc}* and *F_{wwtp}*. In the case of excretion rate, several publications are available on the metabolism of pharmaceuticals and different excretion factors are reported for each drug. The observed differences are probably explained by genomically distinct metabolizing capacities, as well as differences in the routes of administration, sex, age, and health status of the studied subjects (Wishart et al., 2018). On the other hand, removal rates of pharmaceuticals in WWTPs are largely dependent on hydrophobicity and persistence characteristics of the substances. Persistent hydrophilic substances will be present to a greater extent in the treated liquid effluent, hydrophobic ones in the sludge. Expectable removals in WWTPs can thus be inferred to some extent from the substances degradation rates and *K_{ow}* or *K_{oc}* values (Lindim et al., 2016). Removal efficiency values can vary between

10-90% and thus, it is important to use the most accurate value in a given site (Estrada-Arriaga et al., 2016; Gros et al., 2017; Gros et al., 2010; Kasprzyk-Hordern et al., 2009; Lajeunesse et al., 2012; Lin et al., 2009; Rosal et al., 2010). The values can be different in WWTP locations according to the served population, capacity, the configuration and type of treatment, in operating parameters and in hydraulic and solid retention times. Other factors such as meteorological conditions, sampling procedure (grab, composite or flow-proportional (Ort et al., 2010) and sampling period (seasonality) can also affect the empirical F_{wwtp} . In the present study, WWTP removals have been selected following previous published papers, and in the case that no removal values were founded, a 50% of removal was used. Removal efficiencies can range from 9-100%

Finally, DF is the dilution factor used from WWTP effluents to surface waters. Changes in this value can vary the results more than 100-fold (Franquet-Griell et al., 2017). In 2006, EMA recommended the use of a default value of 10 (EMA, 2006). However, a more accurate value has been proposed considering the dilution factor of each country, and a value of 25 as the median of the DF s for Spain (Keller et al., 2014). Initially, PEC_{river} values in the present study were calculated using this DF .

However, to obtain more accurate PEC s, DF s for the main hydrographic basins in Catalonia were calculated to obtain a better representation of the contamination levels, taking into account the specific characteristics of flow and population. Studied basins were Muga and Fluvià, Ter, Besòs, Llobregat, Francolí, Tordera, Ebro, Segre (a tributary of Ebro), and Noguera Pallaresa and Ribagorçana (tributaries of Segre), which cover most of the area. Table 1 shows geographical information of each river studied, including the population, the river basin area, the length, the water use (L/inhab/day) and the mean, minimum and maximum DF calculated. These new DF s were calculated adapting the formula from Keller et al. (2014) (Keller et al., 2014):

$$DF = \frac{Q_r \times 31536000}{inhab\ basin \times W_{basin}} \quad (3)$$

where,

Q_r (m^3/s) is the river flow of each river. The flow data were collected from the hydrographic confederations of each basin and considered geographic and seasonal variability along the basin. Maximum, minimum and mean river flows were used to better estimate PEC variability. The flow data reflect the withdrawal of water in each area. $Inhab_{basin}$ is the population in the basin area (inhabitants). W_{basin} is the water use per capita in the basin, including domestic and industrial use ($\text{m}^3 \text{ inhab}^{-1} \text{ year}^{-1}$). If data was not available, water consumption at national scale ($130 \text{ m}^3 \text{ inhab}^{-1} \text{ year}^{-1}$) was used. 31,536,000 are seconds per year used to convert units.

To calculate PEC_{river} for each basin, the DF derived from high, mean and low flows in each river basin were applied to equation (1) and consumption of pharmaceuticals was considered to be proportional to the population in the studied area. Finally, the $\sum PEC_{river}$ considering each individual drug represented the global occurrence of these compounds in each basin.

2.4. Acute toxicity and Environmental risk assessment (ERA)

The acute toxicity of the prioritize pharmaceuticals was reported from the bibliography. However, in the case of metformin, the acute toxicity was determined in the crustacean *Daphnia magna* following standardized protocols (Organisation of Economic Co-operation and Development (OECD) [1981](#)) according to Gómez-Canela et al. (2014). Exposure concentrations were prepared in American Society for Testing Materials (ASTM) water at 20 °C using acetone as a carrier (0.1 mL L^{-1}). Negative controls (no acetone) and acetone controls gave no response. Aging of ASTM water was conducted in the laboratory at 20 °C in the darkness to prevent photolysis. Assays were conducted in 50 mL of test medium with 10 animals, per duplicate, and were started at <24-h-old neonates and ended at 48 h. Lethal median concentration effects were estimated fitting immobility concentration responses to the Hill regression model.

After calculating PECs in surface water (PEC_{river}), risk assessment was performed to determine if these predicted concentrations could cause hazard in the aquatic environment. The guidelines recommends performing a risk

assessment when PEC_{river} are higher than $0.01 \mu g L^{-1}$ (EMA, 2006). Herein, ERA has been calculated for pharmaceuticals with PEC_{river} values higher than $0.2 \mu g L^{-1}$ in 2016 because of the low toxicity of pharmaceuticals with PEC_{river} values between 0.01 and $0.2 \mu g L^{-1}$. The risk quotient (RQ) was calculated using the following equation (eq. 4), depending on the available data (Gómez-Canela et al., 2014):

$$RQ = \frac{PEC}{PNEC} \approx \frac{PEC}{NOEC/f_1} \approx \frac{PEC}{LC_{50}/f_2} \quad (4)$$

where, PEC is the predicted concentration for a specific basin only in 2016 (described in the previous section) and PNEC is the predicted no-effect concentration. PNEC was estimated using the chronic toxicity NOEC (no-observed effect concentration) and a security factor f_1 of 10 (OECD, 2002). If NOEC was not available, we took LOEC values as a NOEC proxy. When chronic toxicity NOEC/LOEC were not available, PNEC were estimated using $E(L)C_{50}$, and a security factor f_2 of 1000 (OECD, 2002). For metformin, experimental acute toxicity results in *Daphnia magna* and zebrafish embryos were obtained following existing guidelines OECD 202 (OECD, 2004) and OECD 236 (OECD, 2013) used to estimate PNEC. For data interpretation, the maximum probable risk for ecological effects from contaminated water was followed as recommended by Wentsel et al. (Wentsel et al., 1996):

RQ < 1.0 indicates no significant risk;

$1.0 \leq RQ < 10$ indicates a small potential for adverse effects;

$10 \leq RQ < 100$ indicates significant potential for adverse effects;

$RQ \geq 100$ indicates that potential adverse effects should be expected.

3. Results and discussion

3.1. Consumption of pharmaceuticals: the case study of Catalonia.

The mean total consumption of 165 pharmaceuticals in the four years studied (2013-2016) was 623 ± 3 t per year (Table SI2). In 2013, the total consumption

of these pharmaceuticals was 599 t year⁻¹, increasing up to 638 t year⁻¹ in 2016, showing a 6.5% increase. ATC groups N (nervous system), A (alimentary tract and metabolism) and M (musculo-skeletal system) showed the highest consumptions with values between 47 and 318 t year⁻¹ respectively. Figure 1 displays the consumption trends for all pharmaceuticals compiled ordered by their ATC codes from high to low values. Analyzing the period 2013-2016, the pharmaceuticals belonging to ATC groups N, A, J, H and G have increased their consumption in 2016 with respect to 2013. However, the pharmaceuticals belonging to ATC groups M, S, R and L decreased their consumption in 2016 regard to 2013 (see Fig. 1).

A- Alimentary tract and metabolism: The 24 pharmaceuticals for which consumption data was requested had a mean total annual consumption of 182 t year⁻¹ (Table SI2). Metformin, the first-line medication for the treatment of type 2 diabetes, is the most consumed drug in this family with levels between 156 and 164 t year⁻¹, increasing its consumption along the years. The next most consumed drugs were lactulose, omeprazole and ranitidine with levels from 2.3 to 11 t year⁻¹ (Table SI2). The other compounds were consumed at levels < 2 t year⁻¹. Many of these drugs are consumed by the elderly population.

B- Blood and blood forming organs: In this group, the 11 drugs showed a slight decreasing consumption from 18.8 to 17.4 t year⁻¹ was observed (Fig. 1). Acetylsalicylic acid (also known as aspirin) is a medication used to treat pain, fever, or inflammation and was consumed up to 15.9 t year⁻¹ (Table SI2). It was followed by clopidogrel, tranexamic acid and ferrous glycine sulfate. The consumptions of these drugs varied from 0.6 (in 2013) to 1.5 t (in 2016). The remaining seven drugs were below 0.19 t in 2016 (Table SI2).

C- Cardiovascular system: Thirty different drugs administered in Catalonia had a mean total annual consumption of 28.6 t. The highest consumptions in this ATC group corresponded to pentoxifylline, valsartan, simvastatin, enalapril, hydrochlorothiazide, furosemide, atorvastatin, losartan and atenolol with levels between 1.5 to 6.2 t year⁻¹ in the period studied (see Table SI2). The consumptions remained constant all the period, except for pentoxifylline that decreased from 6.2 t (2013) to 4.5 t (2016), while valsartan and losartan

increased their consumptions from 3.2 t (2013) to 5.5 (2016) and from 1.8 t (2013) to 2.2 t (2016), respectively. Herein, the case of furosemide is curious because its values remained slightly constant in the years 2013 (1.8 t), 2014 (1.9 t) and 2016 (2.1 t), but in 2015 its consumption decreased considerably to 0.4 t (Table SI2).

D- Dermatologicals: Among 7 drugs for which consumption data was requested, ketoconazole and clotrimazole were the most consumed pharmaceuticals. Levels in this group were low compared to other ATC groups varied from 0.002 (gentamicin) to 0.32 t year⁻¹ (ketoconazole).

G- Genito urinary system and sex hormones: Only ciproterone, tamusoline and finasteride were the pharmaceuticals compiled for this ATC group. Consumptions increasing along the period studied with values between 0.011 t year⁻¹ and 0.035 t year⁻¹, attributed mainly to finasteride (used for hair loss treatments) which its consumption increased from 27 kg in 2013 up to 36 kg in 2016 (Table SI2). However, overall consumptions are low.

H- Systemic hormonal preparations, excl. sex hormones and insulins: Five different drugs of this ATC group (dexamethasone, methylprednisolone, prednisone, triamcinolone and glucagon) were consumed in Catalonia. The mean total annual consumption of these drugs was 0.3 t in the period studied. The highest consumption was for prednisone, a synthetic corticosteroid used to certain inflammatory diseases and some types of cancer, ranging from 0.24 t in 2013 and 0.29 t in 2016 (see Table SI2).

J- Antiinfectives for systemic use: Twelve drugs belong to this group were dispensed, with consumptions between 30.3 t (in 2013) and 33.5 t (in 2015) (Fig. 1). The slight decreasing between 2015 and 2016 (32.9 t) could be due to the awareness campaign against the inappropriate use of antibiotics promoted by the World Health Organization (WHO) (WHO, 2018). Amoxicillin, used for the treatment of a number of bacterial infections, was the most consumed pharmaceutical accounting for more than the 87% of the total consumption in this ATC group.

L- Antineoplastic and immune modulating agents: Antineoplastic drugs, also called cytostatic drugs, are pharmaceutical used in the cancer treatments. In this study, only megestrol was requested as data for other cytostatics has been previously published (Franquet-Griell et al., 2015; Franquet-Griell et al., 2017). Megestrol, a progestin with antiandrogen activity, is mainly used as an appetite stimulant and sometimes in the treatment of breast and endometrial cancers. Levels of megestrol have been decreasing along the years and its consumption has changed from 0.16 t year⁻¹ in 2013 to 0.11 t year⁻¹ in 2016 (see Table SI2).

M- Musculo-skeletal system: Among 7 drugs for which information was requested, ibuprofen was the pharmaceutical with the highest values followed by allopurinol and diclofenac. The total levels of these 8 drugs were in the range of 64.5 t year⁻¹ (in 2013) decreasing to 47.8 t year⁻¹ (in 2016). In this case, ibuprofen represented approximately the 88% of the total consumption in this ATC group (Table SI2), followed by diclofenac, whose levels decreased considerably along the last years (Table SI2). This lower consumption for ibuprofen in 2016 could be related to the fact that EMA (2014) started to review the cardiovascular risks with systemic ibuprofen medicines (such as those taken by mouth but not topical medicines like creams and gels). The cardiovascular risks being evaluated concern high-dose ibuprofen (2,400 mg per day) taken regularly for long periods (EMA, 2014). As in the case of ibuprofen, in 2013, EMA advised the additional risk to suffer heart attacks using diclofenac at high doses (Baigent et al., 2013; EMA, 2013).

N- Nervous system: This ATC group was the one for which more compounds information was requested, as represent the pharmaceuticals administered mostly to the elder population. Total consumptions of 50 compounds increased from 270 t in 2013 up to 319 t in 2016 (Fig. 1). The most consumed pharmaceuticals were paracetamol (201-235 t year⁻¹), metamizole (13.6-20.9 t year⁻¹), gabapentin (10.9-12.1 t year⁻¹), valproic acid (6.7-6.9 t year⁻¹) and levetiracetam (6.4-8.4 t year⁻¹). The other pharmaceuticals studied in this group had levels between 0.09 and 34.47 t year⁻¹.

R- Respiratory system: In this ATC group, the 8 drugs included had consumptions ranging between 0.48 t in 2013 and 0.34 t in 2016 (Table SI2). Theophylline, a methylxanthine drug used in therapy for respiratory diseases such as asthma, was the most consumed drug with a mean value of 0.29 t year⁻¹. In general, lower levels were reported in comparison to other ATC groups.

S- Sensory organs: Chloramphenicol, norfloxacin, dorzolamide, timolol and latanoprost were the most consumed pharmaceuticals. Among those, norfloxacin, a synthetic chemotherapeutic antibacterial agent, was the most consumed with ranges between 0.57 t year⁻¹ (in 2013) and 0.46 t year⁻¹ (in 2016).

V- Various: This group comprises many different types of drugs. In the present study, naloxone and flumazenil have been included. These two drugs were the least consumed group (Table SI2).

3.2. PECs of pharmaceuticals in wastewaters and rivers

PECs were calculated for the target pharmaceuticals administered mainly in the elder population from Catalonia. Two different PECs (in $\mu\text{g L}^{-1}$) were calculated; PEC_{eff}, which represent the estimated levels in WWTP effluents and PEC_{river}, which takes into account the DF from WWTP to surface water and represents the estimated concentration in the rivers. In fact, the value of PEC calculation relies in having precise information on demography, geographical data and water management issues. Using a DF of 25 according to Keller et al. (Keller et al., 2014), Table SI3 shows PEC values for 165 pharmaceuticals consumed in Catalonia. Out of 165 drugs, 46 had PEC_{river} > 0.01 $\mu\text{g L}^{-1}$, which is the EMA threshold value for risk assessment. Low PECs were attributed to compounds with low consumption, poor excretion or high degradability in the WWTPs.

Considering $\sum\text{PECs}$, values ranged between 148 and 162 $\mu\text{g L}^{-1}$ (PEC_{eff}) and between 5.7 and 6.2 $\mu\text{g L}^{-1}$ (PEC_{river}), observing a slightly increase over the study period. Table 2 shows those drugs with highest PECs obtained (both PEC_{eff} and PEC_{river}), including their consumption during the four years (kg day⁻¹), excretion rate (F_{exc}) and WWTP removal (F_{wwtp}).

Values of predicted concentrations for each drug are explained below, described from higher to lower PEC and according to temporal trends for each compound. In the period analyzed (2013-2016), the pharmaceuticals with highest estimated concentrations in the WWTP effluents and consequently in rivers were metformin, amoxicillin, metamizole, levetiracetam, pentoxifylline and gabapentin, with levels between 5.1 and 46 $\mu\text{g L}^{-1}$ (PEC_{eff}) and between 0.20 and 1.79 $\mu\text{g L}^{-1}$ ($\text{PEC}_{\text{river}}$) in 2016 (see Table 2).

Figure 2 displays the maximum, minimum, the 75% quartile and the 25% quartile of the PEC_{eff} ($\mu\text{g L}^{-1}$) of all pharmaceuticals evaluated. Only the data of 2016 has been represented because the profiles of the other years are similar. The 6 drugs with the highest PECs and which should be prioritized in monitoring studies are described below.

Metformin, a first-line medication for the treatment of type 2 diabetes, was a highly consumed drug with the highest PEC values. The PEC_{eff} and $\text{PEC}_{\text{river}}$ were constant among the period 2013-16, ranging from 43.6 to 46.2 $\mu\text{g L}^{-1}$ and from 1.68 to 1.78 $\mu\text{g L}^{-1}$, respectively (Table 2). It has a high excretion rate of 100% (Wishart et al., 2018) and WWTP removal of 94% (Estrada-Arriaga et al., 2016). Little information is available on metformin in waters. Carmona et al. detected metformin in influent, effluent and river water from Turia River (Spain) at 5.927, 1.252 and 0.013 $\mu\text{g L}^{-1}$, respectively (Carmona et al., 2017).

Second, amoxicillin had a consumption slightly increasing in the period 2013-15, but suffered a decrease in 2016. Its maximum PEC_{eff} and $\text{PEC}_{\text{river}}$ were 37.8 $\mu\text{g L}^{-1}$ and 1.46 $\mu\text{g L}^{-1}$, respectively in 2016 (Table 2). These values are in the order of the levels of amoxicillin in effluents from Girona WWTP that ranged between 0.216 and 0.258 $\mu\text{g L}^{-1}$ (Gros et al., 2013).

The following drug with higher values of PEC was metamizole. This pharmaceutical increased its PEC_{eff} and $\text{PEC}_{\text{river}}$ values from 9.5 and 0.36 $\mu\text{g L}^{-1}$ in 2013 up to 14.8 and 0.57 $\mu\text{g L}^{-1}$ in 2016, respectively (Table 2). Metamizole, a pharmaceutical used as a painkiller, spasm and fever reliever, is excreted partially (Wishart et al., 2018) and thus, it is expected to be found in aquatic environments. Guedes-Alonso et al. (2011) reported levels of 13 pharmaceuticals

in WWTPs from Gran Canaria Island (Spain), detecting concentrations of metamizole between 0.24 and 8.25 $\mu\text{g L}^{-1}$ (Guedes-Alonso et al., 2013). However, not many studies have published the presence of metamizole in waste and river waters from Spain because its rapid degradation to the metabolite 4-acetamidoantipyrine (Huntscha et al., 2012).

Levetiracetam, a medication used to treat epilepsy, is excreted via urine around a 66% of the total drug consumed (Wishart et al., 2018). Its PEC_{eff} and $\text{PEC}_{\text{river}}$ slightly increased from 5.8 to 7.8 $\mu\text{g L}^{-1}$ and from 0.23 to 0.30 $\mu\text{g L}^{-1}$, respectively (Table 2 and Table SI3). The presence of levetiracetam in Spanish waters has never been studied. However, this compound was detected in influents and effluents from a WWTP from Dresden (Germany) with a concentrations higher than 1 $\mu\text{g L}^{-1}$ (Gurke et al., 2015).

Gabapentin, a medication used to treat partial seizures, neuropathic pain, hot flashes, and restless legs syndrome, was estimated in WWTP effluents at concentrations between 4.6 $\mu\text{g L}^{-1}$ (2013) and 5.1 $\mu\text{g L}^{-1}$ (2016), and in catalan rivers between 0.17 and 0.20 $\mu\text{g L}^{-1}$ in the period studied. Gabapentin has also been detected in European rivers at levels of low $\mu\text{g L}^{-1}$ (Klančar et al., 2018).

Finally, pentoxifylline, is a xanthine derivative used as a drug to treat muscle pain in people with peripheral artery disease and is the 6th drug with the highest PECs. PECs ranged between 5.1 and 6.9 $\mu\text{g L}^{-1}$ (PEC_{eff}), and between 0.20 and 0.27 $\mu\text{g L}^{-1}$ ($\text{PEC}_{\text{river}}$), see Table 2 and Table SI3. Pentoxifylline has not been detected in Spanish waters but Vanderford et al. detected this drug at 0.0022 $\mu\text{g L}^{-1}$ in waters from Las Vegas (United States) (Vanderford et al., 2003).

Other highly consumed drugs had low PEC values because of their high removal efficiency in the WWTPs. This is the case of paracetamol, ibuprofen and acetylsalicylic acid. Paracetamol was the pharmaceutical with highest consumption (Table SI3), however its PEC values were low. Paracetamol can be excreted via urine in a 90% (Wishart et al., 2018) and is completely eliminated in WWTP. Its PEC_{eff} ranged from 0.17 and 0.2 $\mu\text{g L}^{-1}$ in the 2013-2016 period and the $\text{PEC}_{\text{river}}$ from 0.0065 and 0.0077 $\mu\text{g L}^{-1}$, following consumption patterns (Table SI3). Many studies have reported the presence of paracetamol in rivers of around the world (Carmona et al., 2017). Paracetamol was detected at mean

concentration of $0.98 \mu\text{g L}^{-1}$ and with 45.5% detection frequency in a monitoring studied carried out in 2014 along the Guadalquivir River (Jaén, South Spain) (Robles-Molina et al., 2014). In another study, López-Roldán et al. detected paracetamol at $0.034 \mu\text{g L}^{-1}$ in the Llobregat River (Catalonia, Spain) (López-Roldán et al., 2010), highly approaching the PECs calculated in this study. Although paracetamol is predicted to be present at very low levels, its high consumption and the high frequency of detection in waters justifies its inclusion in monitoring studies related to water quality. Ibuprofen, a well-known antiinflammatory drug, has a very high consumption (mean= 47.8 t year^{-1}) but taking into account that only 10% is excreted (Wishart et al., 2018), PEC values were low (see Table SI3). Ibuprofen has been recurrently monitored in waste and surface waters (Hedgespeth et al., 2012; Lindberg et al., 2014). In Catalonia, ibuprofen was detected in influent waters ($0.172\text{--}4.21 \mu\text{g L}^{-1}$) and effluent waters ($0.03\text{--}0.95 \mu\text{g L}^{-1}$) from sewage treatment plants (Pedrouzo et al., 2011). Recently, ibuprofen was detected at concentrations up to $2.66 \mu\text{g L}^{-1}$ in sites impacted by raw wastewater in Ebro river (Mandaric et al., 2018). In another study, López-Serna et al. studied the occurrence of 95 pharmaceuticals and transformation products in the metropolis of Barcelona, including Besòs River. Authors reported concentrations of ibuprofen at $0.061 \mu\text{g L}^{-1}$ (López-Serna et al., 2013). Ibuprofen was also detected in Llobregat river at $0.152 \mu\text{g L}^{-1}$ (López-Roldán et al., 2010).

Finally, acetylsalicylic acid had constant PEC_{eff} (from 0.89 to $0.82 \mu\text{g L}^{-1}$) and $\text{PEC}_{\text{river}}$ (from 0.034 to $0.032 \mu\text{g L}^{-1}$) over the studied period. Acetylsalicylic acid has been detected in water as salicylic acid due to its rapid degradation (Skibinski and Komsta, 2016). In the North Sea, the Scheldt estuary and in Belgian harbours salicylic acid was detected at levels ranging from 0.011 to $0.855 \mu\text{g L}^{-1}$ between the period May 2007 and June 2009 (Wille et al., 2010). In another study, salicylic acid was detected in influents and effluents of two WWTPs in Portugal at levels between 1.17 and $61.26 \mu\text{g L}^{-1}$ (influent) and between 0.11 and $0.30 \mu\text{g L}^{-1}$ (effluent) and in the receiving water of the Lis River at a concentration range of 0.025 to $0.29 \mu\text{g L}^{-1}$ (Paíga et al., 2016).

477

3.3. Application of PECs in hydrographic basins

478 To obtain more accurate information and due to the important role of DF in
 479 the calculation of PECs (Franquet-Griell et al., 2017), DFs were recalculated for
 480 the most important hydrographic basin in Catalonia according to the Eq. 3.
 481 Mediterranean rivers, are characterized by high flow fluctuations because of
 482 seasonal variations linked with the weather and precipitations. Thus, it is
 483 important to determine PECs in each river basin considering this flow variability.
 484 For that reason, mean flow values from the middle course of the river, and
 485 maximum and minimum flows were compiled for each river during 2016 to
 486 represent the flow variability (Table 1). Table 3 displays the PEC_{river} ranges in
 487 each river basin for these main compounds detected in each basin. To determine
 488 the global impact of pharmaceuticals in the different basins, total PEC_{river}
 489 ($\sum PEC_{river}$) were calculated considering all compounds using the mean DF of
 490 each river basin in 2016. Figure 3 shows the hydrographic basins studied colored
 491 according to their $\sum PEC_{river}$. Among the eleven main Catalan river basins studied,
 492 Noguera Pallaresa and Noguera Ribagorçana, Fluvià, Segre and Ebro River
 493 (only the part of the river in Catalonia) were the basins with lowest $\sum PEC_{river}$
 494 (between 0.2 and 3.3 $\mu g L^{-1}$), due to either high DF values (Table 1) and/or low
 495 population. Contrarily, Francolí and Besòs were the basins with the highest
 496 $\sum PEC_{river}$ with 50.3 and 66.6 $\mu g L^{-1}$, respectively, as these two rivers have the
 497 lowest DF of all of them (see Table 1) and, in the case of Besòs river, it is
 498 characterized by highly urbanized area. The high variability in the Llobregat river
 499 flow (see Table 1) report high $\sum PEC_{river}$ in the time with low flows and lower
 500 $\sum PEC_{river}$ values in the time with highest flows (Table 3).

501 Table SI4 displays all PEC_{river} values for all pharmaceuticals in each
 502 hydrographic basin in Catalonia considering the mean DF, and minimum and
 503 maximum DFs. In Besòs, Francolí and Llobregat rivers, more than 90% of the
 504 pharmaceuticals compiled had PEC_{river} values higher 0.010 $\mu g L^{-1}$, which is the
 505 EMA proposed threshold (EMA, 2006). However, in the rest of rivers studied, less
 506 than 47% of the pharmaceuticals had PEC_{river} values higher 0.010 $\mu g L^{-1}$. Even
 507 so, the levels of PECs reported in the present study are higher in comparison to

the predicted environmental concentrations of pharmaceuticals in different studies (Burns et al., 2018; Franquet-Griell et al., 2015; Franquet-Griell et al., 2017; Verlicchi et al., 2014). For instance, in Besòs and Francolí, the most impacted river basins in Catalonia, metformin and amoxicillin had a PEC_{river} of 26.5 and 21.3 $\mu\text{g L}^{-1}$ in Besòs basin, and 20.0 and 16.1 $\mu\text{g L}^{-1}$ in Francolí basin (in 2016). Following, metamizole had 8.5 and 6.4 $\mu\text{g L}^{-1}$, levetiracetam 4.5 and 3.4 $\mu\text{g L}^{-1}$, gabapentin 2.9 and 2.2 $\mu\text{g L}^{-1}$ and pentoxifylline acid 2.9 and 2.2 $\mu\text{g L}^{-1}$ respectively for the two basins (Table 3). On the other hand, Nogueres and Ebro, which accounted for the lowest impacted basins, PEC_{river} for metformin was of 0.8-0.12 and 0.08 $\mu\text{g L}^{-1}$ respectively, much lower than values predicted in the other basins. Recently, Lindim et al. evaluated the emissions and concentrations of 54 pharmaceuticals in Swedish river basins, reporting that metformin, gabapentin and atenolol were the pharmaceuticals with the highest emissions (Lindim et al., 2016). The prioritized pharmaceuticals for monitoring (metformin, amoxicillin, metamizole, levetiracetam, gabapentin and pentoxifylline) would always be detected in low river flow conditions (low values of DF). In this scenario, Llobregat would also be impacted by high concentrations of pharmaceuticals whereas in Muga, Fluvià, Ter, Tordera, Ebro, Segre and Nogueres the PEC_{river} values would be lower (Table 3 and Table SI4). Thus, because the studied rivers have high variable flow, the use of specific DFs which consider the seasonal variability of water flows can provide a better characterization of the presence of pharmaceuticals in a specific river basin and thus, a more accurate risk assessment.

3.4. Environmental risk assessment (ERA) of pharmaceuticals: the case study of Catalonia.

For the six pharmaceuticals having the highest PECs, we calculated the environmental risks using the worst river scenario (Besòs river), which has the lowest dilution factor of the studied rivers and hence the highest PECs. We also included paracetamol, ibuprofen and acetylsalicylic acid due to their highly

prescription rate and usage in Spain. Reported concentration effects from acute and chronic toxicity studies for at least three different aquatic phyla were considered (i.e. algae or plant, vertebrate, invertebrate) to estimate predicted non-effect concentrations (PNECs). Assessment factors of 10 and 1000 were used to determine PNECs from chronic and acute concentration effects, respectively. Toxicity data was obtained from toxicological databases (DrugBank, EPA), chemical suppliers (Sigma-Aldrich) and selected publications and experimental test for metformin. When more than one concentration effect was reported within a given toxicity source, chemical and species, its median was considered. Environmental risks were obtained using the risk quotient approach depicted in eq. 4. Results reported in Table 4 reflect the reported known scarcity for chronic toxicity data on pharmaceuticals, which in most cases was limited to a single species (Fent et al., 2006a). For three chemicals, levetiracetam, gabapentin and pentoxifylline there was no toxicological information and for metamizole, toxicological information was limited to a single study. Table 4 also evidenced the great disparity of toxicological information within and among species, which in several cases expanded several orders of magnitude (Santos et al., 2010). For the antibiotic amoxicilin cyanobacteria species were about three orders of magnitude more sensitive than the rest of species, which is in line with previous studies (Andreozzi et al., 2004). Paracetamol acute toxicity data was the most variable within a given species (over 4 orders of magnitude for *Danio rerio* LC₅₀) and ibuprofen showed the greatest differences between acute and chronic toxicity responses (350 fold in *Danio rerio*) and among algae species (85 fold). Disparity of results within species for metformin and acetylsalicylic acid were within an order of magnitude. When outlier values from Table 4 were not considered, *Daphnia* and algae species were about 10 fold more sensitive than fish to paracetamol and acetylsalicylic acid and fish were up to 300 fold more sensitive to metformin and ibuprofen than *Daphnia* and algae. The above mentioned lack of consistence in species sensitivity across the studied pharmaceuticals agrees with previous reported toxicological information (Santos et al., 2010) but partly disagrees with the argument claiming that pharmaceuticals should affect to greater extent fish than algae or invertebrates due to the presence of more biological targets in fish (Fent et al., 2006a; Gunnarsson et al., 2008).

Estimated risks reported in Table 4 showed the same high variability within and among species and chemicals than PNEC. Thus to properly prioritize environmental risks of the studied compounds we excluded out the extreme values for ibuprofen (RQ= 4.71), paracetamol (RQ= 11.43), metformin (RQ= 6.22) and amoxicillin (RQ= 5325). Except for metamizole and gabapentin, whose high risk quotients (170 and 0.29, respectively) were based on a single study, the compound having the greatest risks to aquatic biota was amoxicillin (average $\pm$ SE, RQ= 1.47 ± 1.40) followed by decreasing order by metformin (0.66 ± 0.21), ibuprofen (0.26 ± 0.23), acetylsalicylic acid (0.004 ± 0.001) and paracetamol (0.002 ± 0.002). For amoxicillin, metformin and ibuprofen predicted risks exceeded for some species. Therefore, environmental risk assessment data depicted in Table 4 allowed identifying key features among the prioritized 5 pharmaceuticals with at least three RQ. Amoxicillin is expected to pose adverse effects for cyanobacteria, and metformin and ibuprofen pose a small potential for adverse effects to invertebrates and fish, and fish, respectively. Alternatively, paracetamol and acetylsalicylic acid posed no risk to aquatic biota.

As a final remark it is important to take into account that in this study we based our prioritization on PECs and not on toxicity. For example, among the 43 compounds depicted in Table 2 disulfiram followed by fluoxetine, lithium and diclofenac were quite toxic having EC_{50} for zebrafish embryonic development or *D. magna* immobilization of (Mean $\pm$ SE): $0.027 \pm 0.013 \text{ mg L}^{-1}$ for disulfiram, $0.399 \pm 0.104 \text{ mg L}^{-1}$ for fluoxetine, $3.340 \pm 1.274 \text{ mg L}^{-1}$ for lithium and $4.752 \pm 1.248 \text{ mg L}^{-1}$ for diclofenac (data obtained from E.P.A. (EPA), ECOTOX Knowledgebase). These values are quite low when compared with those reported in Table 4, which means that the above reported compound can also be considered potential toxic to aquatic biota. Future research, thus, should assess if there is a need to prioritize compounds by their toxicity rather than by their PECs.

4. Conclusions

This study reveals the importance of consumption data and temporal patterns for estimating the occurrence and risk of pharmaceuticals consumed by elderly

people in surface waters. An extensive data compilation on pharmaceuticals consumption in Catalonia was performed. The mean total consumption of these pharmaceuticals in the period studied (2013-2016) was 623 ± 3 t per year. ATC groups N (nervous system), A (alimentary tract and metabolism) and M (musculo-skeletal system) showed the highest consumptions, being paracetamol, metformin and ibuprofen the top consumed. However, metformin, amoxicillin, metamizole, were the pharmaceuticals with highest PEC values ($> 0.01 \mu\text{g L}^{-1}$). In addition, recalculation of PECs according to specific river dilution factors permits to refine the levels likely to be detected at river basin scale. It is clear from this study that PEC calculation permits to better prioritize compounds, which have high probability to be detected in the environment. Finally, predicted environmental levels together with acute and chronic toxicological data allowed to estimate the risks of these compounds. Amoxicillin is expected to pose adverse effects for cyanobacteria, and metformin and ibuprofen pose a small potential for adverse effects to invertebrates and fish, and fish, respectively. Alternatively, paracetamol and acetylsalicylic acid posed no risk to aquatic biota.

Acknowledgments

The authors wish to thank CatSalut, and especially Pere Carbonell and Montserrat Bosch, for kindly providing the consumption data of pharmaceuticals in Catalonia. The authors also gratefully acknowledge Teresa Sala-Comorera for helping in the initial calculations of PECs. This study has been financed by the SUDOE program with the project Innovec'Eau (2016-2019) / Project SOE1/P1/F0173 Interreg SUDOE funded by FEDER: 1.177.875,64 €; <http://innovec-eau.univ-perp.fr>.

638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672

References

Andreozi R, Caprio V, Ciniglia C, De Champdoré M, Lo Giudice R, Marotta R, et al. Antibiotics in the environment: Occurrence in Italian STPs, fate, and preliminary assessment on algal toxicity of amoxicillin. *Environmental Science and Technology* 2004; 38: 6832-6838.

Baigent C, Bhala N, Emberson J, Merhi A, Abramson S, Arber N, et al. Vascular and upper gastrointestinal effects of non-steroidal anti-inflammatory drugs: Meta-analyses of individual participant data from randomised trials. *The Lancet* 2013; 382: 769-779.

Besse JP, Kausch-Barreto C, Garric J. Exposure assessment of pharmaceuticals and their metabolites in the aquatic environment: Application to the French situation and preliminary prioritization. *Human and Ecological Risk Assessment* 2008; 14: 665-695.

Burns EE, Carter LJ, Kolpin DW, Thomas-Oates J, Boxall ABA. Temporal and spatial variation in pharmaceutical concentrations in an urban river system. *Water Research* 2018; 137: 72-85.

Carballa M, Omil F, Lema JM. Comparison of predicted and measured concentrations of selected pharmaceuticals, fragrances and hormones in Spanish sewage. *Chemosphere* 2008; 72: 1118-1123.

Carmona E, Andreu V, Picó Y. Multi-residue determination of 47 organic compounds in water, soil, sediment and fish—Turia River as case study. *Journal of Pharmaceutical and Biomedical Analysis* 2017; 146: 117-125.

Cleuvers M. Mixture toxicity of the anti-inflammatory drugs diclofenac, ibuprofen, naproxen, and acetylsalicylic acid. *Ecotoxicology and Environmental Safety* 2004; 59: 309-315.

Crane M, Watts C, Boucard T. Chronic aquatic environmental risks from exposure to human pharmaceuticals. *Science of the Total Environment* 2006; 367: 23-41.

- Daouk S, Chèvre N, Vernaz N, Bonnabry P, Dayer P, Daali Y, et al. Prioritization methodology for the monitoring of active pharmaceutical ingredients in hospital effluents. *Journal of Environmental Management* 2015; 160: 324-332.
- Daughton CG. Cradle-to-cradle stewardship of drugs for minimizing their environmental disposition while promoting human health. I. Rational for and avenues toward a green pharmacy. *Environmental Health Perspectives* 2003; 111: 757-774.
- EMA. Guideline on the environmental risk assessment of medicinal products for human use. In: Agency EM, editor. Guideline on the environmental risk assessment of medicinal products for human use. EMEA/CHMP/SWP/4447/00, 2006.
- EMA. PRAC recommends the same cardiovascular precautions for diclofenac as for selective COX-2 inhibitors. 2017, 2013.
- EMA. European Medicines Agency starts review of ibuprofen medicines, 2014.
- Estrada-Arriaga EB, Cortés-Muñoz JE, González-Herrera A, Calderón-Mólgora CG, de Lourdes Rivera-Huerta M, Ramírez-Camperos E, et al. Assessment of full-scale biological nutrient removal systems upgraded with physico-chemical processes for the removal of emerging pollutants present in wastewaters from Mexico. *Science of The Total Environment* 2016; 571: 1172-1182.
- Fent K, Weston AA, Caminada D. Ecotoxicology of human pharmaceuticals. *Aquatic Toxicology* 2006a; 76: 122-159.
- Fent K, Weston AA, Caminada D. Erratum to “Ecotoxicology of human pharmaceuticals” [*Aquatic Toxicology* 76 (2006) 122–159]. *Aquatic Toxicology* 2006b; 78: 207.
- Franquet-Griell H, Gómez-Canela C, Ventura F, Lacorte S. Predicting concentrations of cytostatic drugs in sewage effluents and surface waters of Catalonia (NE Spain). *Environmental Research* 2015; 138: 161-172.
- Franquet-Griell H, Gómez-Canela C, Ventura F, Lacorte S. Anticancer drugs: Consumption trends in Spain, prediction of environmental concentrations and potential risks. *Environmental Pollution* 2017; 229: 505-515.
- Generalitat de Catalunya - Catalan Health System Observatory. Pharmacy prescriptions invoiced to Catalan Health Service 2018, 2018.
- Gómez-Canela C, Barata C, Lacorte S. Occurrence, elimination, and risk of anticoagulant rodenticides and drugs during wastewater treatment. *Environmental Science and Pollution Research* 2014; 21: 7194-7203.
- Gros M, Blum KM, Jernstedt H, Renman G, Rodríguez-Mozaz S, Haglund P, et al. Screening and prioritization of micropollutants in wastewaters from on-site sewage treatment facilities. *Journal of Hazardous Materials* 2017; 328: 37-45.
- Gros M, Petrović M, Ginebreda A, Barceló D. Removal of pharmaceuticals during wastewater treatment and environmental risk assessment using hazard indexes. *Environment International* 2010; 36: 15-26.
- Gros M, Rodríguez-Mozaz S, Barceló D. Rapid analysis of multiclass antibiotic residues and some of their metabolites in hospital, urban wastewater and river water by ultra-high-performance liquid chromatography coupled to quadrupole-linear ion trap tandem mass spectrometry. *Journal of Chromatography A* 2013; 1292: 173-188.
- Guedes-Alonso R, Afonso-Olivares C, Montesdeoca-Esponda S, Sosa-Ferrera Z, Santana-Rodríguez JJ. An assessment of the concentrations of pharmaceutical compounds in wastewater treatment plants on the island of Gran Canaria (Spain). *SpringerPlus* 2013; 2: 1-8.
- Gunnarsson L, Jauhiainen A, Kristiansson E, Nerman O, Larsson DGJ. Evolutionary conservation of human drug targets in organisms used for environmental risk assessments. *Environmental Science and Technology* 2008; 42: 5807-5813.
- Guo J, Sinclair CJ, Selby K, Boxall ABA. Toxicological and ecotoxicological risk-based prioritization of pharmaceuticals in the natural environment. *Environmental Toxicology and Chemistry* 2016; 35: 1550-1559.

Gurke R, Rossmann J, Schubert S, Sandmann T, Rößler M, Oertel R, et al. Development of a SPE-HPLC-MS/MS method for the determination of most prescribed pharmaceuticals and related metabolites in urban sewage samples. *Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences* 2015; 990: 23-30.

Hedgspeth ML, Sapozhnikova Y, Pennington P, Clum A, Fairey A, Wirth E. Pharmaceuticals and personal care products (PPCPs) in treated wastewater discharges into Charleston Harbor, South Carolina. *Science of the Total Environment* 2012; 437: 1-9.

Hernando MD, Mezcuá M, Fernández-Alba AR, Barceló D. Environmental risk assessment of pharmaceutical residues in wastewater effluents, surface waters and sediments. *Talanta* 2006; 69: 334-342.

Hignite C, Azarnoff DL. Drugs and drug metabolites as environmental contaminants: Chlorophenoxyisobutyrate and salicylic acid in sewage water effluent. *Life Sciences* 1977; 20: 337-341.

Huntscha S, Singer HP, McArdell CS, Frank CE, Hollender J. Multiresidue analysis of 88 polar organic micropollutants in ground, surface and wastewater using online mixed-bed multilayer solid-phase extraction coupled to high performance liquid chromatography-tandem mass spectrometry. *Journal of Chromatography A* 2012; 1268: 74-83.

IDESCAT. Official Statistics Website of Catalonia. 2017, 2017.

Kasprzyk-Hordern B, Dinsdale RM, Guwy AJ. The removal of pharmaceuticals, personal care products, endocrine disruptors and illicit drugs during wastewater treatment and its impact on the quality of receiving waters. *Water Research* 2009; 43: 363-380.

Keller VDJ, Williams RJ, Lofthouse C, Johnson AC. Worldwide estimation of river concentrations of any chemical originating from sewage-treatment plants using dilution factors. *Environmental Toxicology and Chemistry* 2014; 33: 447-452.

Klančar A, Trontelj J, Roškar R. Development of a Multi-Residue Method for Monitoring 44 Pharmaceuticals in Slovene Surface Water by SPE-LC-MS/MS. *Water, Air, and Soil Pollution* 2018; 229.

Kümmerer K. Drugs in the environment: emission of drugs, diagnostic aids and disinfectants into wastewater by hospitals in relation to other sources – a review. *Chemosphere* 2001; 45: 957-969.

Kümmerer K. The presence of pharmaceuticals in the environment due to human use – present knowledge and future challenges. *Journal of Environmental Management* 2009; 90: 2354-2366.

Lacorte S, Luis S, Gómez-Canela C, Sala-Comorera T, Courtier A, Roig B, et al. Pharmaceuticals released from senior residences: occurrence and risk evaluation. *Environmental Science and Pollution Research* 2017: 1-12.

Lajeunesse A, Smyth SA, Barclay K, Sauvé S, Gagnon C. Distribution of antidepressant residues in wastewater and biosolids following different treatment processes by municipal wastewater treatment plants in Canada. *Water Research* 2012; 46: 5600-5612.

Lin AY-C, Yu T-H, Lateef SK. Removal of pharmaceuticals in secondary wastewater treatment processes in Taiwan. *Journal of Hazardous Materials* 2009; 167: 1163-1169.

Lindberg RH, Östman M, Olofsson U, Grabic R, Fick J. Occurrence and behaviour of 105 active pharmaceutical ingredients in sewage waters of a municipal sewer collection system. *Water Research* 2014; 58: 221-229.

Lindim C, van Gils J, Georgieva D, Mekenyan O, Cousins IT. Evaluation of human pharmaceutical emissions and concentrations in Swedish river basins. *Science of The Total Environment* 2016; 572: 508-519.

López-Roldán R, de Alda ML, Gros M, Petrovic M, Martín-Alonso J, Barceló D. Advanced monitoring of pharmaceuticals and estrogens in the Llobregat River basin (Spain) by liquid chromatography-triple quadrupole-tandem mass spectrometry in combination with ultra performance liquid chromatography-time of flight-mass spectrometry. *Chemosphere* 2010; 80: 1337-1344.

López-Serna R, Jurado A, Vázquez-Suñé E, Carrera J, Petrović M, Barceló D. Occurrence of 95 pharmaceuticals and transformation products in urban groundwaters underlying the metropolis of Barcelona, Spain. *Environmental Pollution* 2013; 174: 305-315.

Mandarić L, Mor J-R, Sabater S, Petrović M. Impact of urban chemical pollution on water quality in small, rural and effluent-dominated Mediterranean streams and rivers. *Science of The Total Environment* 2018; 613-614: 763-772.

McGrath NA, Brichacek M, Njardarson JT. A Graphical Journey of Innovative Organic Architectures That Have Improved Our Lives. *Journal of Chemical Education* 2010; 87: 1348-1349.

Navarro-Ortega A, Acuña V, Batalla RJ, Blasco J, Conde C, Elorza FJ, et al. Assessing and forecasting the impacts of global change on Mediterranean rivers. The SCARCE Consolider project on Iberian basins. *Environmental Science and Pollution Research* 2012a; 19: 918-933.

Navarro-Ortega A, Sabater S, Barceló D. Understanding effects of global change on water quantity and quality in river basins- The SCARCE Project. *Environmental Science and Pollution Research* 2012b; 19: 915-917.

Njardarson JT. Top selling pharmaceuticals. 2018, 2010.

OECD. Report of the OECD Workshop on Environmental Hazard/risk Assessment, 2002.

OECD. Test No. 202: *Daphnia* sp. Acute Immobilisation Test, 2004.

OECD. Test No. 236: Fish Embryo Acute Toxicity (FET) Test, 2013.

Ort C, Lawrence MG, Reungoat J, Mueller JF. Sampling for PPCPs in wastewater systems: Comparison of different sampling modes and optimization strategies. *Environmental Science and Technology* 2010; 44: 6289-6296.

Overturf MD, Anderson JC, Pandelides Z, Beyger L, Holdway DA. Pharmaceuticals and personal care products: A critical review of the impacts on fish reproduction. *Critical reviews in toxicology* 2015; 45: 469-491.

Owen SF, Giltrow E, Huggett DB, Hutchinson TH, Saye J, Winter MJ, et al. Comparative physiology, pharmacology and toxicology of β -blockers: Mammals versus fish. *Aquatic Toxicology* 2007; 82: 145-162.

Paíga P, Santos LHMLM, Ramos S, Jorge S, Silva JG, Delerue-Matos C. Presence of pharmaceuticals in the Lis river (Portugal): Sources, fate and seasonal variation. *Science of the Total Environment* 2016; 573: 164-177.

Pedrouzo M, Borrull F, Pocurull E, Marcé RM. Presence of pharmaceuticals and hormones in waters from sewage treatment plants. *Water, Air, and Soil Pollution* 2011; 217: 267-281.

Robles-Molina J, Gilbert-López B, García-Reyes JF, Molina-Díaz A. Monitoring of selected priority and emerging contaminants in the Guadalquivir River and other related surface waters in the province of Jaén, South East Spain. *Science of The Total Environment* 2014; 479-480: 247-257.

Rosal R, Rodríguez A, Perdígón-Melón JA, Petre A, García-Calvo E, Gómez MJ, et al. Occurrence of emerging pollutants in urban wastewater and their removal through biological treatment followed by ozonation. *Water Research* 2010; 44: 578-588.

Santos LHMLM, Araújo AN, Fachini A, Pena A, Delerue-Matos C, Montenegro MCBSM. Ecotoxicological aspects related to the presence of pharmaceuticals in the aquatic environment. *Journal of Hazardous Materials* 2010; 175: 45-95.

Skibinski R, Komsta L. The stability and degradation kinetics of acetylsalicylic acid in different organic solutions revisited - An UHPLC-ESI-QTOF spectrometry study. *Current Issues in Pharmacy and Medical Sciences* 2016; 29: 39-41.

Ternes T, Bonerz M, Schmidt T. Determination of neutral pharmaceuticals in wastewater and rivers by liquid chromatography–electrospray tandem mass spectrometry. *Journal of Chromatography A* 2001; 938: 175-185.

United Nations (UN). World Population Ageing, 2015.


- Valderrama Gama E, Rodríguez Artalejo F, Palacios Díaz A, Gabarre Orús P, Pérez del Molino Martín J. Consumo de medicamentos en los ancianos: resultados de un estudio poblacional. *Revista Española de Salud Pública* 1998; 72: 209-219.
- Vanderford BJ, Pearson RA, Rexing DJ, Snyder SA. Analysis of Endocrine Disruptors, Pharmaceuticals, and Personal Care Products in Water Using Liquid Chromatography/Tandem Mass Spectrometry. *Analytical Chemistry* 2003; 75: 6265-6274.
- Verlicchi P, Al Aukidy M, Jelic A, Petrović M, Barceló D. Comparison of measured and predicted concentrations of selected pharmaceuticals in wastewater and surface water: A case study of a catchment area in the Po Valley (Italy). *Science of The Total Environment* 2014; 470-471: 844-854.
- Wentzel RS, Thomas W, Point L, Simini M, Checkai RT, Ludwig D. Tri-service procedural guidelines for ecological risk assessments: Defense Technical Information Center, 1996.
- WHO. Antibiotic resistance. 2018, 2018.
- Wille K, Noppe H, Verheyden K, Vanden Bussche J, De Wulf E, Van Caeter P, et al. Validation and application of an LC-MS/MS method for the simultaneous quantification of 13 pharmaceuticals in seawater. *Analytical and Bioanalytical Chemistry* 2010; 397: 1797-1808.
- Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: a major update to the DrugBank database for 2018. *Nucleic Acids Research* 2018; 46: D1074-D1082.

Figure captions

Figure 1. Consumption of pharmaceuticals (in t year⁻¹) ordered by their ATC codes.

Figure 2. Boxplot in logarithmic scale of pharmaceuticals PEC_{eff} levels in 2016 along each ATC family. The rest of the years studied have not been represented because their same profiles. Supplementary information provides complete data for all the years for which consumption data was requested (Table SI3).

Figure 3. $\sum$ PEC_{river} (µg L⁻¹) of pharmaceuticals in the main hydrographic basins of Catalonia in 2016.




862

863

Figure 1. Consumption of pharmaceuticals (in t year⁻¹) ordered by their ATC codes.

864




865

866

867

Figure 2. Boxplot in logarithmic scale of pharmaceuticals PEC_{eff} levels in 2016 for each ATC family. The rest of the years studied have not been represented because they have similar profiles. N indicates the number of pharmaceuticals studied in each family.

868
869
870
871



872
873 **Figure 3.** $\Sigma\text{PEC}_{\text{river}}$ of pharmaceuticals in the main hydrographic basins of
874 Catalonia in 2016.

875