

The International Surface Pressure Databank version 2

Thomas A. Cram^{1,*}, Gilbert P. Compo^{2,3}, Xungang Yin⁴, Robert J. Allan^{5,6}, Chesley McColl^{2,3}, Russell S. Vose⁷, Jeffrey S. Whitaker³, Nobuki Matsui^{2,3}, Linden Ashcroft^{8,†}, Renate Auchmann⁹, Pierre Bessemoulin¹⁰, Theo Brandsma¹¹, Philip Brohan⁶, Manola Brunet^{12,17}, Joseph Comeaux¹, Richard Crouthamel¹³, Byron E. Gleason Jr⁷, Pavel Y. Groisman^{7,14,15}, Hans Hersbach¹⁶, Philip D. Jones^{17,18}, Trausti Jónsson¹⁹, Sylvie Jourdain²⁰, Gail Kelly⁵, Kenneth R. Knapp⁷, Andries Kruger²¹, Hisayuki Kubota²², Gianluca Lentini^{23,‡}, Andrew Lorrey²⁴, Neal Lott⁷, Sandra J. Lubker³, Jürg Luterbacher²⁵, Gareth J. Marshall²⁶, Maurizio Maugeri²³, Cary J. Mock²⁷, Hing Y. Mok²⁸, Øyvind Nordli²⁹, Mark J. Rodwell^{6,§}, Thomas F. Ross⁷, Douglas Schuster¹, Lidija Srnec³⁰, Maria Antónia Valente³¹, Zsuzsanna Vizi^{32,¶}, Xiaolan L. Wang³³, Nancy Westcott³⁴, John S. Woollen^{35,36} and Steven J. Worley¹

¹National Center for Atmospheric Research, Boulder, CO, USA

²University of Colorado, CIRES, Boulder, CO, USA

³NOAA Earth System Research Laboratory, Physical Sciences Division, Boulder, CO, USA

⁴ERT, Inc., Asheville, NC, USA

⁵Atmospheric Circulation Reconstructions over the Earth Initiative, Met Office Hadley Centre, Exeter, UK

⁶Met Office Hadley Centre, Exeter, UK

⁷NOAA National Centers for Environmental Information, Asheville, NC, USA

⁸School of Earth Sciences, University of Melbourne, Melbourne, Vic., Australia

⁹Institute of Geography and Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

¹⁰Météo-France, Toulouse, France

¹¹Royal Netherlands Meteorological Institute (KNMI), De Bilt, The Netherlands

¹²Centre for Climate Change, Universitat Rovira i Virgili, Tarragona, Spain

¹³International Environmental Data Rescue Organization (IEDRO), Deale, MD, USA

¹⁴University Corporation for Atmospheric Research, Boulder, CO, USA

¹⁵P. P. Shirshov Institute of Oceanology, Russian Academy of Sciences, Moscow, Russia

¹⁶European Centre for Medium-Range Weather Forecasts (ECMWF), Reading, UK

¹⁷Climatic Research Unit, University of East Anglia, Norwich, UK

¹⁸Center of Excellence for Climate Change Research/Department of Meteorology, King Abdulaziz University, Jeddah, Saudi Arabia

¹⁹Icelandic Met Office, Reykjavik, Iceland

²⁰Météo-France, Direction de la Climatologie, Toulouse, France

²¹South African Weather Service, Pretoria, South Africa

²²Department of Coupled Ocean–Atmosphere–Land Processes Research, Japan Agency for Marine–Earth Science and Technology, Yokosuka, Japan

²³Dipartimento di Fisica, Università degli Studi di Milano, Milan, Italy

²⁴National Institute of Water and Atmospheric Research (NIWA), Auckland, New Zealand

²⁵Department of Geography, Climatology, Climate Dynamics and Climate Change, Justus Liebig University Giessen, Giessen, Germany

²⁶British Antarctic Survey, Natural Environment Research Council, Cambridge, UK

²⁷Department of Geography, University of South Carolina, Columbia, SC, USA

²⁸Hong Kong Observatory, Hong Kong, China

²⁹Norwegian Meteorological Institute, Oslo, Norway

³⁰Meteorological and Hydrological Service, Zagreb, Croatia

³¹Instituto Dom Luiz, Laboratório Associado, Universidade de Lisboa, Lisbon, Portugal,

³²Nicolaus Copernicus University, Toruń, Poland

³³Climate Research Division, Science and Technology Branch, Environment Canada, Toronto, ON, Canada

³⁴Midwestern Regional Climate Center, Illinois State Water Survey, University of Illinois at Urbana–Champaign, Urbana, IL, USA,

³⁵I.M. Systems Group, Inc., Rockville, MD, USA

³⁶NOAA National Centers for Environmental Prediction, Environmental Modeling Center, Camp Springs, MD, USA

*Correspondence: Thomas Cram, National Center for Atmospheric Research, P.O. Box 3000, Boulder, CO 80307-3000, USA, E-mail: tcram@ucar.edu

†Current address: Centre for Climate Change, Universitat Rovira i Virgili, Tarragona, Spain

‡Current address: Ente Regionale per i Servizi all'Agricoltura e alle Foreste (ERSAF), Milan, Italy

§Current address: European Centre for Medium-Range Weather Forecasts (ECMWF), Reading, UK

¶Current address: Department of Space and Climate Physics, University College London, London, UK

The International Surface Pressure Databank (ISPD) is the world's largest collection of global surface and sea-level pressure observations. It was developed by extracting observations from established international archives, through international cooperation with data recovery facilitated by the Atmospheric Circulation Reconstructions over the Earth (ACRE) initiative, and directly by contributing universities, organizations, and countries. The dataset period is currently 1768–2012 and consists of three data components: observations from land stations, marine observing systems, and tropical cyclone best track pressure reports. Version 2 of the ISPD (ISPDv2) was created to be observational input for the Twentieth Century Reanalysis Project (20CR) and contains the quality control and assimilation feedback metadata from the 20CR. Since then, it has been used for various general climate and weather studies, and an updated version 3 (ISPDv3) has been used in the ERA-20C reanalysis in connection with the European Reanalysis of Global Climate Observations project (ERA-CLIM). The focus of this paper is on the ISPDv2 and the inclusion of the 20CR feedback metadata. The Research Data Archive at the National Center for Atmospheric Research provides data collection and access for the ISPDv2, and will provide access to future versions.

Geosci. Data J. **2**: 31–46 (2015), doi: 10.1002/gdj3.25

Received: 15 May 2014, revised: 3 June 2015, accepted: 8 June 2015

Key words: surface pressure, sea level pressure, observations, stations, ships

Dataset

Identifier: doi:<http://dx.doi.org/10.5065/D6SQ8XDW>

Creators: Compo, G. P., J. S. Whitaker, P. D. Sardeshmukh, N. Matsui, R. J. Allan, X. Yin, B. E. Gleason, R. S. Vose, G. Rutledge, P. Bessemoulin, S. Brönnimann, M. Brunet, R. I. Crouthamel, A. N. Grant, P. Y. Groisman, P. D. Jones, M. C. Kruk, A. C. Kruger, G. J. Marshall, M. Maugeri, H. Y. Mok, O. Nordli, T. F. Ross, R. M. Trigo, X. L. Wang, S. D. Woodruff and S. J. Worley

Title: International Surface Pressure Databank (ISPDv2)

Publisher: Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory, Boulder, Colorado, USA

Publication year: 2010

Resource type: Dataset

Version: 2.0

Introduction

The International Surface Pressure Databank (ISPD) is the world's largest collection of global surface and sea-level pressure observations. Since its inception in 2002, its development has been facilitated by cooperative efforts between the Global Climate Observing System (GCOS)/World Climate Research Program (WCRP) Working Group on Observational Data Sets for Reanalysis [2007 to 2011], and the ongoing efforts of the GCOS Working Group on Surface Pressure and the international Atmospheric Circulation Reconstructions over the Earth initiative (ACRE, Allan *et al.*, 2011). The full observational record is extracted and assembled from established international archives and newly available collections from more than 60 different contributing organizations, shown in Tables 1 and 2.

The ISPD version 2 (ISPDv2) covers the period 1768–2012. It merges data from three input components: observations from land stations, marine observing systems, and tropical cyclone best track pressure reports. The land station component was extracted from many national and international collections of sea-level pressure and surface pressure. The largest contributor to this component was the Integrated Surface Database (Smith *et al.*, 2011), which consists of

global hourly and synoptic surface observations collected from many sources. The land stations were merged following a two-step algorithm to first remove duplicates within a collection and then remove duplicates between collections (Yin *et al.*, 2008).

The marine component consists of sea-level pressure observations extracted from the International Comprehensive Ocean–Atmosphere Data Set (ICOADS, Woodruff *et al.*, 1998; Parker *et al.*, 2004; Woodruff *et al.*, 2005; Worley *et al.*, 2005; Woodruff *et al.*, 2011), a global ocean marine meteorological and surface ocean dataset that is considered the most complete of its kind. It is comprised of measurements and observations extracted from many international data sources, including ship reports, moored and drifting buoys, coastal stations, and other marine platforms. ICOADS release 2.4 was used in the ISPDv2 for the period 1952–2008 and ICOADS release 2.5 (Woodruff *et al.*, 2011) was used for the period 1785–1951 and 2009–2012.

The tropical cyclone component was taken from the International Best Track Archive for Climate Stewardship (IBTrACS; Knapp *et al.*, 2010). The IBTrACS dataset consists of global tropical cyclone best track position and intensity observations, and reports collected from each of the World Meteorological Organization (WMO) Regional Specialized Meteorological

Table 1. Contributing organizations to the International Surface Pressure Databank.

1. All-Russian Research Institute of Hydrometeorological Information World Data Center^a
2. Atmospheric Circulation Reconstructions over the Earth (ACRE)^a
3. Australian Bureau of Meteorology^a
4. Australian Meteorological Association, Todd Project team
5. British Antarctic Survey^a
6. Cook Islands Meteorological Service
7. Danish Meteorological Institute^a
8. Deutscher Wetterdienst (DWD)^a
9. Environment Canada, Climate Research Division^a
10. ETH Zurich, Switzerland^a
11. European and North Atlantic Daily to Multidecadal Climate Variability (EMULATE)^a
12. EUropean Reanalysis and Observations for Monitoring (EURO4M)/The WMO MEditerranean DAta REscue Initiative (MEDARE)
13. European Reanalysis of Global Climate Observations (ERA-CLIM)
14. GCOS Atmospheric Observation and Ocean Observation Panels for Climate Working Group on Surface Pressure^a
15. GCOS/WCRP Working Group on Observational Data Sets for Reanalysis^a
16. Hong Kong Observatory^a
17. Icelandic Meteorological Office (IMO)
18. Instituto Geofísico da Universidade do Porto, Portugal^a
19. International Best Track Archive for Climate Stewardship (IBTrACS)^a
20. International Comprehensive Ocean–Atmosphere Data Set (ICOADS)^a
21. International Environmental Data Rescue Organization (IEDRO)^a
22. Japan Agency for Marine–Earth Science and Technology (JAMSTEC)
23. Japan Meteorological Agency^a
24. Jersey Met Department^a
25. Koninklijk Nederlands Meteorologisch Instituut (KNMI; Royal Netherlands Meteorological Institute)^a
26. Lamont–Doherty Earth Observatory of Columbia University
27. McGill University, Canada
28. Met Office Hadley Centre, UK^a
29. Météo-France^a
30. Météo-France – Division of Climate
31. Meteorological and Hydrological Service, Croatia^a
32. National Center for Atmospheric Research (NCAR), USA^a
33. National Institute for Water and Atmospheric Research (NIWA), New Zealand
34. Nicolaus Copernicus University – Department of Meteorology and Climatology, Poland
35. Niue Meteorological Service
36. NOAA Climate Database Modernization Program (CDMP), USA^a

(continued)

Table 1. (continued)

37. NOAA Earth System Research Laboratory (ESRL), Physical Sciences Division, USA^a
38. NOAA Midwest Regional Climate Center at UIUC, USA^a
39. NOAA National Centers for Environmental Prediction (NCEP), USA^a
40. NOAA National Climatic Data Center (NCDC), USA^a
41. NOAA Northeast Regional Climate Center at Cornell University, USA^a
42. NOAA Pacific Marine Environmental Laboratory, USA
43. Norwegian Meteorological Institute^a
44. Ohio State University – Byrd Polar Research Center, USA^a
45. Oldweather.org
46. Portuguese Institute of Sea and Atmosphere (IPMA), Portugal
47. Proudman Oceanographic Laboratory, UK^a
48. Signatures of environmental change in the observations of the Geophysical Institutes (SIGN)^a
49. South African Weather Service^a
50. South Eastern Australian Recent Climate History (SEARCH) project, The University of Melbourne
51. Tanzania Meteorological Agency
52. University of Aberdeen, Scotland, UK
53. University of Bern, Switzerland
54. University of Colorado – Climate Diagnostics Center (CDC) of the Cooperative Institute for Research in Environmental Sciences (CIRES)^a
55. University of East Anglia – Climatic Research Unit, UK^a
56. University of Giessen – Department of Geography, Germany
57. University of Lisbon – Instituto Dom Luiz, Portugal^a
58. University of Milan – Department of Physics, Italy^a
59. University of Porto-Instituto Geofísico, Portugal^a
60. University Rovira i Virgili – Centre for Climate Change (C3), Spain^a
61. University of South Carolina, USA
62. University of Toronto – Department of Physics, Canada
63. University of Washington, USA
64. World Meteorological Organization – MEditerranean climate DAta REscue (MEDARE)
65. Zentralanstalt für Meteorologie und Geodynamik (ZAMG; Austrian Weather Service)^a

^aContributors to version 2 (cf. Compo *et al.*, 2011, Table II).

Centers, Tropical Cyclone Warning Centers, and other national agencies. The IBTrACS Beta version was used for the years 1952–2006, version v01r01 for 1871–1883 and 1886–1951, version v02r01 for 1884–1885 and 2007–2008, version v03r02 for 2009–2010, and version v03r05 for 2011–2012.

The inclusion of various versions of both ICOADS and IBTrACS in ISPDv2 was the result of using the most up-to-date data that were available during the assimilation into the Twentieth Century Reanalysis (20CR; Compo *et al.*, 2011). This rationale is elaborated upon in Section 2.1.

Table 2. ISPDv2 dataset collection identifiers and names.

ISPD ID	Name	Description	Period	No. stations/records	NCDC reference	NCAR reference	Reference publication
0100	ICODS Release 2.1	Global Marine Surface Observations	1784–2005	–/–185 million	DSI-1170	ds540.0, ds540.1	Parker <i>et al.</i> (2004), Woodruff <i>et al.</i> (1998, 2005), Worley <i>et al.</i> (2005)
0104	ICODS Release 2.4	Global Marine Surface Observations	1784–2007	–/–238 million	DSI-1170	ds540.0, ds540.1	Parker <i>et al.</i> (2004), Woodruff <i>et al.</i> (1998, 2005), Worley <i>et al.</i> (2005)
0105	ICODS Release 2.5	Global Marine Surface Observations	1784–2007	–/–261 million	DSI-1173	ds540.0, ds540.1	
0200	ICODS Auxiliary Kobe	Global Marine Surface Observations	1889–1943	–/3135		ds530.0	
0300	ICODS Auxiliary Whaling	Global Marine Surface Observations	1950–1984	–/20		ds530.0	
0400	ICODS Auxiliary Russian	Global Marine Surface Observations	1950–2000	–/1789		ds530.0	
0500	ICODS Auxiliary Russian	Global Marine Surface Observations	1889–2000	–/7873		ds530.0	
0700	ICODS Auxiliary Challenger	Global Marine Surface Observations	1872–1876	(n/a)		ds530.0	
1000	Federal Climate Complex Integrated Surface Database	Global Land Surface Observations	1901–2008	23 363/–1.2 billion	DSI-3505	ds463.3	Smith <i>et al.</i> (2011)
1002	CDMP SAO/1001 Forms	US Land Surface Observations	1928–1948	27/236 886	DSI-3851		Dupigny-Giroux <i>et al.</i> (2007)
1003	Russian Empire Stations	Russian Land Surface Observations	1849–2000	1860/87 795 180	DSI-9290c		
1004	Air Weather Service TD13	Global Land Surface Observations	1901–1973	32/55 748		ds467.0	
1005	Hadley Centre	Individual Stations from Hadley Centre	1833–present	11/520 594			Allan <i>et al.</i> (2011)
1006	CDMP-International	Chile, Mexico, Uruguay	1800s–1980	14/326 175			
1007	READER Antarctic & Southern Hemisphere	20 Stations via British Antarctic Survey	1947–2007	18/562 784			Dupigny-Giroux <i>et al.</i> (2007)
1010	U.S. Air Force DATSAV	US Air Force Compilation	1976–1980		DSI-9685	ds463.0	
1011	Royal Netherlands Meteorological Institute (KNMI)	KNMI stations	1911–2006	3/1 045 250			
1012	CDMP Forts	US Army Signal Service and other 19 th Century Voluntary Observations	1841–1893	49/629 491	DSI-3297		Dupigny-Giroux <i>et al.</i> (2007)
2000	NCEP-NCAR BUFR Archive	Global Observations	1948–2003	6625/26 225 199		ds090.0	Kalnay <i>et al.</i> (1996), Kistler <i>et al.</i> (2001)
2001	NCEP Operational BUFR Archive	Global Observations	1928–1948, 2003–2005	1053/2 940 516		ds090.0	
3002	WASA Stations Observations	Northern Europe, Greenland	1871–1996	20/2 181 628			Schmith <i>et al.</i> (1997, 1998)
3004	Sea Level Pressure Environmental Canada Pressure Observations	Canadian Stations	1842–2004	746/25 277 856			
3005	West African Synoptic observations	West African Land Surface	1850–1980	123/3 022 702			
3006	The Australian Bureau of Meteorology Station Pressure Dataset	Australian Land stations	1900–1956	63/1 418 093			

(continued)

Table 2. (continued)

ISPD ID	Name	Description	Period	No. stations/records	NCDC reference	NCAR reference	Reference publication
3007	Northern Italian Pressure Observations		1878–1940	1/62 253			Maugeri <i>et al.</i> (2004, 2008)
3008	Brazil Surface Observations		1951–1980	8/78			
3009	Spanish Hourly Pressure Observations from EMULATE	Hourly Spanish Land Stations	1850–2003	5/190 899	ds486.0		Ansell <i>et al.</i> (2006)
3010	Germany	Deutscher Wetterdienst (DWD) Web Archive	1876–2000	9/405 167			
3011	Austria	Emulate Stations	1872–2002	9/467 124			Ansell <i>et al.</i> (2006)
3012	Switzerland	Emulate Stations	1900–1973	8/815 363			Ansell <i>et al.</i> (2006)
3013	South Africa	South African Weather Service Stations	1850–2003	187/2 636 479			
3014	Norway	Land Stations	1863–2007	16/225 687			
3015	Croatia Meteorological and Hydrological Service Land Stations		1858–2005	4/444 398			
3016	Portugal	Portuguese SIGN stations	1860–2006	2/151 989			Valente <i>et al.</i> (2008)
4000	Hong Kong Hourly Pressure Observations	Hong Kong Observatory	1885–1939	1/430 920			
4001	Jakarta/Batavia Pressure Observations	Dutch Royal Observatory	1866–1944	1/604 518	ds490.0		
4002	Liverpool	Proudman Ocean. Lab stations	1768–1793	1/9094			
4003	Jersey Channel Island Pressure Observations	Channel Island Stations	1864–1913	7/159 781			
4004	CMDP-USNO	US Naval Observatory at Washington	1841–1913	1/83 261			Dupigny-Giroux <i>et al.</i> (2007)
5000	Antarctic Expedition keyed by Hadley Centre		1899–1941	11/119 991			Allan <i>et al.</i> (2011)
5002	Byrd Antarctic Expeditions Observations	Monthly Weather Review Supplemental No. 41	1929–1930				
8000	Atlantic/North Eastern Pacific Hurricane Reanalysis	US Hurricane Re-Analysis Dataset	1848–present				
8001	International Best Track Archive for Climate Stewardship (IBTrACS)	National Climatic Data Center	1848–present		DS1-9637		Knapp <i>et al.</i> (2010)
010000–019999	NCAR Upper Air Stations	National Center for Atmospheric Research	1943–1998	1081/1 275 272	ds370.1		

The columns denote, respectively, the ISPDv2 collection ID, collection name, description, yearly period of record, total number of stations and observational records in the collection, reference number for the corresponding National Climatic Data Center (NCDC) dataset, reference number for the corresponding dataset archived in the Research Data Archive (RDA) at the National Center for Atmospheric Research (NCAR), and reference publication.

*The Air Weather Service TD13 observational records originating from collection ID 1004 are incorrectly labeled with collection ID 2001 in the native HDF5 tables in both ISPDv2 and ISPD version 3.

1. Dataset content and coverage

The ISPDv2 dataset is stored in the Hierarchical Data Format version 5 (HDF5: <http://www.hdfgroup.org/>), which allows for efficient archiving of large, complex data structures and a diverse set of metadata. Each HDF5 file contains 13 tables and 15 directory subgroups that define how to decode the individual records, and each observation record includes metadata defining the data collection source and observation type (e.g. station observation, marine observation, surface pressure observation from a radiosonde sounding).

Because the ISPDv2 was assimilated as observational input into the 20CR, each record also includes the 20CR quality control and assimilation feedback metadata. These quality control and feedback metadata include the results of the five-step quality control procedure described in Appendix B of Compo *et al.* (2011) and all the relevant statistical quantities returned by the 20CR ensemble data assimilation system. The format of the HDF5 metadata, therefore, is designed to allow traceability of observations from their original source to the ISPDv2, and to permit direct feedback from the 20CR data assimilation back to the original source archives.

Each observation at any single observation time is assigned a unique number that when combined with the time stamp (year, month, date, hour, and minute) forms a unique identifier within the complete dataset. Thus, every observation has an unambiguous reference within the full ISPDv2 dataset, which allows for precise identification and traceability. This data management feature is critical for accurately adding additional metadata from other future reanalyses, and

evolving the ISPDv2 to new versions while maintaining the usage provenance of each record.

The 20CR required hourly files from the ISPDv2 data for the data assimilation system, thus each data file in the ISPDv2 collection contains one hour of global observations. Based on this file organization and data coverage, a maximum of 24 data files per day and 8760 data files per (non-leap) year exist in the ISPDv2 collection. The total dataset volume is 491.32 GB. The HDF5 data are available for download as either individual hourly data files, or as yearly tar files, which are tarballs of one entire year of HDF5 data files.

The ISPDv2 is also available in NetCDF and ASCII column data formats, both of which are derived from the native HDF5 format. The data content within these two formats is not a complete reproduction of the tabular table content and metadata that are contained within the HDF5 data. It does contain, however, essential information, including the 20CR quality control feedback information, for most users to carry out their research projects. The ASCII data are available as monthly tar files for direct download from the National Center for Atmospheric Research (NCAR) Research Data Archive (RDA) website (<http://dx.doi.org/10.5065/D6SQ8XDW>), while the NetCDF data are produced on demand through the data subsetting service described in Section 3.1.

The data availability varies considerably over the period of record. Maps of annual land station distribution (Figure 1) highlight the steady growth in the number of reporting stations, and hence data coverage, since the year 1850 (station coverage maps for every year are available at <http://www.esrl.noaa.gov/psd/data/ISPD/v2.0/>). The sources of land station




Figure 1. Annual distribution of ISPDv2 land station locations for the years 1850, 1900, 1950, and 2000. Station coverage maps for every year in the ISPDv2 are available at <http://www.esrl.noaa.gov/psd/data/ISPD/v2.0/>.

observations in the 19th and early 20th centuries are located primarily in Europe and North America. The data coverage increases dramatically across most of the rest of the globe from 1950 to the present.

Figure 2 shows time series of the total number of land observing stations in each indicated continental region by year from 1850 to 2010. The notable decline in stations during the mid-1960s – primarily in the Asia/Eastern Europe time series – is attributed to the reduction in the Integrated Surface Database records during this period, and is explained in Smith *et al.* (2011) as the result of the transition from keying of observational data to digital transmission and receipt of data.

The Asia time series also exhibits a considerable decline in station coverage starting in the mid-1980s. A listing of the number of stations originating from the Integrated Surface Database for the years 1963–1964 and selected years between 1980 and 2001 is shown in Table 3, which reveals these reductions are primarily associated with the observations originating from the Russian Federation and other former Soviet Union countries. These reductions can be explained by the deterioration of funding of the meteorological observation network in the final years of the Soviet Union existence and thereafter in the 1990s and 2000s. The observation network health during this post-Soviet period was a function of the economic situation in each of the 15 newly independent nations (and, to some extent, a function of foreign sponsorship). The China, North and Central America, and South-West Pacific regions also exhibit notable decreases between 1963 and 1964.

The average number of pressure observations per day contained in the 20CR feedback records from all

ISPDv2 components (station, marine, and tropical cyclone) is illustrated by maps for the years 1900, 1950, and 2000 in Figure 3. The preponderance of observations in the Northern Hemisphere in earlier years is evident, as is the growth of marine observations over time (see Woodruff *et al.*, 2011 for a complete description).

The total number of observations per year in the ISPDv2 increases steadily from approximately 100 000 records in 1870 to 53 million in 2010 (Figure 4). A local increase and subsequent decrease in this trend during the 1878–1894 period is attributed to the addition of 1.8 million records from the US Marine Meteorological Journals and digitized for ICOADS Release 2.5 (Woodruff *et al.*, 2011). A decrease during the World War I years is evident, while an increase during the World War II years is attributed to the inclusion of 1.6 million ICOADS records digitized from the UK Royal Navy Ships' logbooks (Brohan *et al.*, 2009; Woodruff *et al.*, 2011). A decrease during the mid-1960s is associated with the aforementioned decline in station records in the Integrated Surface Database (Smith *et al.*, 2011). A decrease in 2007 is most likely due to a delay in processing, and therefore incomplete records, in the Integrated Surface Database. Other small-scale variations in the curve likely can be correlated with increased scientific interest (local peaks) and research budget reductions (local troughs; Fleming, 2000).

The dataset contains reports from more than 10 000 land stations, which, in addition to the marine observing systems reports, comprise over 1.5 billion observations in total. Between 1860 and 1917, the total number of marine and surface station observations is comparable, but in later years the total count





Figure 2. Time series of the total number of ISPDv2 land observing stations in each indicated continental region by year from 1850 to 2010. The inset panel indicates the boundaries used for the continental regions.


Table 3. Total number of stations originating from the Integrated Surface Database and included in ISPDv2 for selected years between 1963 and 2001.

WMO region	1963	1964	1980	1990	1993	1995	1998	2001
0	47	28	877	1007	925	927	869	1058
1	184	166	717	813	740	748	777	935
2	914	223	800	695	683	577	410	482
3	795	124	1031	878	835	751	607	584
4	466	353	916	932	864	876	867	1049
5	518	347	727	598	592	587	558	638
6	211	222	689	768	758	776	825	887
7	484	280	1524	1594	1671	1697	1560	1770
8	132	132	550	695	661	684	630	671
9	689	626	578	865	874	898	860	841

The columns represent the station count for the years listed and organized by WMO regional block number (0 and 1 = Europe; 2 and 3 = Russian Federation and other former Soviet Union countries; 4 = Asia; 5 = China; 6 = Africa; 7 = North America, Central America, and the Caribbean; 8 = South America and Antarctica; and 9 = South-West Pacific).

**Figure 3.** Maps showing the average number of pressure observations per day contained in the 20CRv2 feedback records from all ISPDv2 components: station, marine, and tropical cyclone for the years 1900 (top), 1950 (middle), and 2000 (bottom). Counts of observations are made in five-degree grid boxes.

is dominated by land surface station reports (not shown). This prevalence of land station observations is primarily due to the input data compiled for the

**Figure 4.** Time series of the number of pressure observations per year in version 2 of the International Surface Pressure Databank (ISPDv2) from 1870 to 2010. Note the logarithmic scale along the y-axis. Inset panel: time series during the same period showing the number of observations in the Northern Hemisphere (blue curve) and Southern Hemisphere (red curve).

National Centers for Environmental Prediction (NCEP)/NCAR Reanalysis project (1948–2003; Kalnay *et al.*, 1996; Kistler *et al.*, 2001).

2. Data usage and application

2.1. Reanalysis

The primary motivation for the creation and development of the ISPD was to facilitate progress on research and understanding of long-term trends and variations in global surface pressure. In particular, a long-term historical dataset, such as the ISPD, becomes essential to providing an observational underpinning to retrospective climate analysis datasets, commonly known as reanalyses. Reanalysis products are used extensively in climate research, applications, and services, including for monitoring and comparing current climate condi-

tions with those of the past, identifying the causes of climate variations and change, and preparing climate predictions. Information derived from reanalyses is also being used increasingly in commercial and business applications in sectors such as energy, agriculture, water resources, and insurance.

Prior to completion of the ISPD, most reanalysis products (including those from the NCEP/NCAR Reanalysis Project (Kalnay *et al.*, 1996; Kistler *et al.*, 2001) and the European Centre for Medium-Range Weather Forecasts [ECMWF; Uppala *et al.*, 2005]) could only extend back to about 1950 since they relied on assimilating a set of observations that included upper-level atmospheric information. The primary input data source for the NCEP/NCAR Reanalysis, for example, are the global rawinsonde observations, which contain a period of record substantial enough to produce that reanalysis back to 1948. The limited time range of these reanalyses restricts their usefulness for many climate research applications. The ISPD adds value by providing a much longer period of record and thus enables the development of reanalyses with longer time spans.

The first reanalysis to make use of the ISPD is the 20CR Project (Compo *et al.*, 2011), which assimilates only the ISPDv2 surface and sea-level pressure observations and prescribes observed monthly sea-surface temperature and sea-ice distributions from HadISST1.1 (Rayner *et al.*, 2003) as boundary conditions. This global reanalysis dataset spans the late 19th century, the entire 20th century, and the early 21st century (1871–2012). It estimates the state of the atmosphere – the ‘analysis’ – by combining the hourly and synoptic ISPDv2 observations in a 6-h time window with a dynamically generated 9-h first guess forecast initialized from the previous analysis. Cycling this procedure with overlapping 6-h analyses and 9-h forecasts has the effect of spreading the observational information from the ISPD three-dimensionally in space and in time.

As the 20CR was produced, the early period ISPDv2 ‘beta’ version was revised to include newly digitized additions and improve the overall coverage of stations. As ICOADS Release 2.5 was then available, it was also included to increase the number of marine observations in the early period. Similarly for IBTrACS, newly digitized records from the most recent versions of IBTrACS were added to ISPDv2 as it evolved and improved over time. The final ISPDv2, therefore, contains multiple source versions of ICOADS and IBTrACS. The newer versions of ICOADS and IBTrACS did not change appreciably except to expand their data inputs and coverage, therefore we do not anticipate that the use of multiple versions introduces inconsistencies in the pressure dataset.

2.2. Twentieth century reanalysis quality control and data assimilation feedback

During the data assimilation processing of the 20CR, the ISPDv2 input data were interrogated by a five-step

quality control procedure that included, among other steps, checking the observations for meteorological plausibility, comparing them with neighbouring data values, and performing a bias correction on the land component (see Compo *et al.*, 2011 for more information on this procedure). The 20CR then applied an Ensemble Kalman Filter data assimilation method which used background first-guess fields supplied by an ensemble of 56 forecasts from an experimental version of the NCEP Global Forecast System (GFS) numerical weather prediction model (Kanamitsu *et al.*, 1991; Moorthi *et al.*, 2001; Saha *et al.*, 2006). These quality control and data assimilation results, which include the first-guess fields, analysis departures, bias estimates, and observation errors, were then written back into the ISPDv2 so that each observational record contains this feedback information. Future users of the ISPDv2, and those who contributed the original observational data, therefore can utilize this information to make an informed decision on the quality and usefulness of each observational record during the 20CR time period.

One project that utilized the 20CR feedback metadata is the ERA-20C reanalysis (Poli *et al.*, 2013), which is the first reanalysis produced under the European Reanalysis of Global Climate Observations database project (ERA-CLIM: www.era-clim.eu). The ERA-20C is a global reanalysis that spans the 20th century (1900–2010) assimilating only surface and sea-level pressure observations from ISPD and ICOADS and marine surface winds from ICOADS. Feedback information from the 20CR was used in the bias correction scheme for ISPD data that were assimilated in the ERA-20C reanalysis; this was based on a break-point analysis using the 20CR first-guess departures (Hersbach *et al.*, 2015). In locations where breakpoints were suspected (e.g. instances of instrument errors or a change in station location), the ERA-20C bias correction scheme assigned less confidence to the 20CR first-guess value, and thus was allowed to be more adaptive compared to cases where no irregularities in long-term departures were identified.

In another study, Wang *et al.* (2014) used the 20CR feedback metadata to show that the strong extratropical cyclone events in the 20CR agree well with the geostrophic wind extremes derived from *in situ* pressure observations. This study illustrates how the 20CR quality control procedure performed well in identifying errors in the observational record. For example, the 20CR quality control system identified and rejected 143 of 146 erroneous observation values from the Aberdeen, Scotland records for the period 1871–1921. This quality control feedback information, therefore, is valuable in identifying observation errors which might otherwise be overlooked and, coupled with continued data rescue efforts such as those facilitated by ACRE (Allan *et al.*, 2011), serves to strengthen and reduce uncertainty in the observational record.

3. Data access and user services

3.1. Research data archive data location and accessibility

The primary repository for ISPDv2 is in the RDA (<http://rda.ucar.edu>) at NCAR in Boulder, Colorado. The RDA is a free and open data collection where data discovery can be achieved through faceted searches based on Global Change Master Directory (GCMD: <http://gcmd.nasa.gov>) metadata keywords, free text queries, and lists highlighting the most used datasets. Data access is free, but requires each user to register through a simple online process that validates the submitted email address.

The ISPDv2 data access is organized by year and month on the RDA website (<http://dx.doi.org/10.5065/D6SQ8XDW>), and users may browse through the pages to locate specific data of interest. Data files are organized in default groups by observation year and month. They can be downloaded directly from the RDA web interface using server-supplied 'wget' scripts. Users may select a collection of files to download from default lists by using the 'Web File Listing' option under the 'Data Access' tab.

A more refined option is provided through the data subset request form, which may be accessed via the 'Get a subset' link under the 'Data Access' tab. This produces a customized data subset of the HDF5 data based on user-provided constraints. Users can specify the temporal limits, spatial domain, observation type (s), and ASCII or NetCDF data output formats through the data subset request form (Figure 5). The output file compression can also be requested on this form (not shown). Observations from individual observing stations may also be requested, which is a useful feature for users who wish to procure time series for specific locations or regions. Once submitted, the data subset requests are produced by a delayed mode data processing procedure. Users are notified and directed to a web download location when their data request output files are accessible.

3.2. Dataset citation

The RDA also provides a data citation service. The dataset homepage provides citation syntax in the standard forms recommended by the Federation of Earth System Information Partners (ESIP), American Meteorological Society (AMS), American Geophysical Union (AGU), DataCite, and the Geoscience Data Journal. This dataset citation is also available in Research Information System (RIS) format so that users may import the citation for this dataset directly into their citation reference management software (e.g. EndNote, Zotero, etc.). One key element in the dataset citation is the data access date ('Accessed dd mmm yyyy'). Leveraging the fact that RDA data users are registered, a customized data citation can be prepared

for the user by using the 'Get a customized data citation' link on the dataset home page. The dates that users access the data are recorded and can be retrieved on demand at later times.

3.3. File content metadata

File content metadata are collected as part of the data processing and preparation of the ISPDv2 for the RDA. The metadata supports additional information services for interested users. The overall ISPDv2 metadata summary is provided from the dataset home page under the 'More Details' link. Here, tabulations of the observing station location, observation type, platform identification, and maps of the global distribution for the full ISPDv2 can be viewed (Figure 6).

In addition, users may view the content metadata for the yearly tar file archives by clicking on a looking glass icon adjacent to the tar file listing, accessible under the 'Data Access' tab. This service enables users to query the ISPDv2 via an interactive map of land station locations, and more efficiently determine the observational data available at any particular time and region of interest (Figure 7).

3.4. Supporting software and documentation

Source code developed at the University of Colorado CIRES and NOAA Earth System Research Laboratory and written in C language to read and decode the HDF5 data into ASCII column output is provided on the RDA website under the 'Software' tab on the dataset home page. Documentation describing the HDF5 tables and ASCII output is also available for download under the 'Documentation' tab on the dataset home page.

3.5. ECMWF data location and accessibility

A subset of the ISPDv2 is also archived at ECMWF (<http://apps.ecmwf.int/datasets>). Here, users may extract temporal subsets from the ISPDv2 archive stored at ECMWF and refine their selection by observation platform. The output from these requests is written in ASCII column format and contains the date, location, observed value, report type, unique identifier which allows traceability to other ISPD versions, and several feedback parameters from the 20CR, including the 20CR bias estimate, the 20CR ensemble mean first guess pressure minus the 'modified' observed pressure, and the ensemble mean analysis pressure minus the 'modified' observed pressure. The 'modified' pressure is the pressure value after the 20CR system adjusted it to be consistent with the orography in the assimilating model. This is similar to a reduction to sea level but instead is an adjustment to the 20CR orography, which could be higher or lower than the station elevation. In addition to the ASCII output, users may produce observation count maps for their data selection on the ECMWF web interface.

Select Temporal Range

Start Date

End Date

1768-01-01

2010-12-31

Reset Range

We recommend that you limit your temporal range to 25 years because of the large number of ISPD records to process. This will avoid the possibility of an interruption while processing your order. An interruption would require a manual restart and delay your order.

Select Spatial Range

Enter latitude & longitude manually

Select location by station ID

Map

Satellite

Interactive Map Instructions:

- Use the 'Pan Map' option to drag and center the map on your area of interest
- Use the 'Draw Box' option to drag a box around your area of interest. You can also manually enter bounding latitudes and longitudes in the text boxes.

North*: 90

West*: -180

East*: 180

South*: -90

Reset

*Latitudes and longitudes must be specified in whole degrees

Zoom: - +

Pan Map

Draw Box

Select Observation Types

Select which observation types to retain in your data request. All will be retained if none are selected.

☐ Radiosonde observations
☐ Dropsonde observations
☐ Marine observations
☐ Surface station observations
☐ Tropical cyclone best track data

or

Select Output Data Format

☒ NetCDF
☐ ASCII

Figure 5. The data subsetting interface found on the ISPDv2 Research Data Archive webpage after user authentication. Users choose temporal range, spatial selection (choices are interactive map functions, manual latitude/longitude entry, or station identifier entry), and also can select one or many observation types, and ASCII or NetCDF output data formats.

4. Future plans

Since the release of ISPDv2, over 22 additional organizations have contributed to the station and marine components of the ISPD (Table 1) and the period of record now extends back to 1755. These new contributions have been incorporated into a newer version 3 of the ISPD (ISPDv3), which will be made available in the near future through the NCAR RDA. The observation feedback archive (OFA) version of this dataset as used in the ERA-20C reanalysis is available from ECMWF (Hersbach *et al.*, 2015).

The gain in observations for ISPDv3 is illustrated by the time series shown in Figure 8. Most of the increase occurs in the Northern Hemisphere, and there are visible increases in the late 19th century and during the World War I years. The 2007 decrease in merged records from the Integrated Surface Database into ISPDv2 (cf. Figure 4) is recovered in ISPDv3. These and increases in other years come from diverse station and marine collections contributed to the ISPD under the auspices of GCOS, WCRP, and the ACRE initiative. This includes the initial efforts of the Old-Weather.org citizen science project that focused on UK





Summary for platform observations in HDF5 format			
Observing platform	Observation type • Data types	Average frequency of data (varies by individual platform ID and data type)	Temporal/geographical coverage (each dot represents a 3° box containing one or more observations)
CMAN station	Surface • Assimilation feedback parameters • Observed sea level pressure (hPa)	<ul style="list-style-type: none"> • 1 to 6 per min • 1 to 51 per h • 1 to 20 per day 	1983-01-01 to 2011-12-31 
Coastal station	Surface • Assimilation feedback parameters • Observed sea level pressure (hPa)	<ul style="list-style-type: none"> • 1 per day • 5 per week 	1986-01-01 to 1989-12-31 
Drifting buoy	Surface • Assimilation feedback parameters • Observed sea level pressure (hPa) • Observed surface pressure (hPa)	<ul style="list-style-type: none"> • 1 to 10 per min • 1 to 50 per h • 1 to 20 per day • 1 to 6 per week • 1 to 4 per month • 1 to 10 per year 	1899-01-01 to 2011-12-31 
Fixed ocean platform	Surface • Assimilation feedback parameters • Observed sea level pressure (hPa)	<ul style="list-style-type: none"> • 1 to 7 per h • 1 to 11 per day • 3 per month 	1971-01-01 to 1997-12-31 

Figure 6. Example of the content metadata interface found on the ISPDv2 Research Data Archive webpage. Here, users can view and investigate tabulations of the location, observation type, platform identification, and maps of the global distribution of observations for the full dataset.

Royal Navy observations during 1914–1923, the citizen-led Todd Project team (<http://www.charlestodd.net>) that focused on recovering observations assembled by Sir Charles Todd, Meteorologist of the Colony of South Australia during 1879–1909, and the efforts of many other university, national, and international organizations (Table 1). Another notable contribution are the North African locations, which arose from the joint effort between the European Union-funded European Reanalysis and Observations for Monitoring (EURO4M: <http://www.euro4m.eu>) project in connection with the WMO/Mediterranean climate

Data REscue (MEDARE) initiative (<http://www.omm.urv.cat/MEDARE>; Brunet *et al.*, 2014). The merging of these contributions results in increases that are particularly dramatic when viewed as a map, as shown for the example year of 1918 (Figure 9). The increases over eastern South America, eastern Africa, China, New Zealand, and selected central Pacific islands are especially evident.

ISPDv3 (specifically, version 3.2.6) was implemented into the European Reanalysis of Global Climate Observations database (ERA-CLIM: www.era-clim.eu), with a goal of using these and additional observations




Figure 7. Example of the file content metadata interface for the yearly tar files found on the ISPDv2 Research Data Archive webpage. This service enables users to probe the ISPDv2 via an interactive map of land station locations, and more efficiently search for the observational data at any particular time and region of interest.

in an ECMWF data assimilation system to generate new global climate reanalyses of the 20th century. The ERA-20C global reanalysis, the first reanalysis under this project, spans the 20th century (1900–2010) assimilating only surface and sea-level pressure observations from ISPDv3 and marine wind observations from ICOADS 2.5 using a 4D-Var assimilation system (Poli *et al.*, 2013).

In addition, ISPD version 3.2.9 is being assimilated into the next generation of the 20CR (version 2c). 20CRv2c uses the same NCEP atmosphere/land coupled model at the same spectral resolution of total wavenumber 62 (about 200 km by 200 km) as

20CRv2 but has different prescribed boundary conditions. Daily sea surface temperatures come from a new Simple Ocean Data Assimilation with sparse input version 2 (SODAsi.2, B.S. Giese, H.F. Seidel, G.P. Compo and P.D. Sardeshmukh, in review). Prescribed monthly averaged sea ice concentrations from the new COBE-SST2 (Hirahara *et al.*, 2014) correct the known misspecification in 20CRv2 (Compo *et al.*, 2011; Brönnimann *et al.*, 2012). This activity is currently underway and will be reported on in due course.

The ISPD also will be used in the planned centennial coupled reanalysis being developed under the Japa-




Figure 8. Time series of the number of pressure observations per year in version 2 (thin blue curve) and version 3.2.8 (red curve) of the International Surface Pressure Databank (ISPD) from 1800 to 2011. Note the logarithmic scale along the y-axis. Inset panel: Time series (shown on a linear scale) during the period 1820 to 1920 showing the number of observations in the Northern Hemisphere (blue curves) and Southern Hemisphere (red curves). The shading highlights the gain in observations between the two versions.




Figure 9. Upper panel: Map showing the daily average number of pressure observations in the new ISPDv3.2.8 for the year 1918. Lower panel: Difference in the daily average number of pressure observations between ISPDv3.2.8 and the ISPDv2 (cf. Figure 3). Counts of observations are made in five-degree grid boxes.

nese research project Program for Risk Information on Climate Change (SOUSEI: http://www.jamstec.go.jp/sousei/eng/research/theme_a.html), by the Meteorological Research Institute of the Japan Meteorological Agency and collaborators (M. Ishii, 2014, pers. comm.).

Comments and questions on the ISPD can be directed to the RDA data specialist or left at the reanalyses.org webpage for this dataset (<http://reanalyses.org/observations/international-surface-pressure-databank>).

Acknowledgements

Construction of the ISPD is supported by the U.S. Department of Energy Office of Science (Biological and Environmental Research program) and the NOAA Climate Program Office. The Twentieth Century Reanalysis Project and the International Surface Pressure Databank used resources of the National Energy Research Scientific Computing Center and of the Oak Ridge Leadership Computing Facility at Oak Ridge National Laboratory, which are supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231 and Contract No. DE-AC05-00OR22725, respectively. Manola Brunet and Philip D. Jones are grateful to the European Union-funded projects EURO4M, FP7-SPACE-2009-1 Project No. 242093) and Uncertainties in Ensembles of Regional Reanalyses (UERRA, FP7-SPACE-2013-1 Project No. 607193). Pavel Groisman was partially supported by Grant 14.B25.31.0026 of the Russian Ministry of Education and Science. Renate Auchmann was supported by the Swiss National Science Foundation (SNF; Project TWIST, 200021_146599/1).

The authors thank the following individuals and organizations for their contributions: (1) Rajmund Przybylak of the Nicolaus Copernicus University, Department of Meteorology and Climatology, Poland. (2) Frank Le Blancq and many others, Jersey Met Service, for Channel Island data. (3) The IT group at CIRES and NOAA/ESRL Physical Sciences Division. (4) Prashant Sardeshmukh of University of Colorado/CIRES for valuable guidance and input. (5) Paul DellaMarta and MeteoSwiss colleagues who originally provided data to EMULATE. (6) Ingeborg Auer of Zentralanstalt für Meteorologie und Geodynamik (ZAMG) who originally provided Austrian data to EMULATE. (7) Clive Wilkinson of NCDC and Climatic Research Unit, University of East Anglia, Norwich, UK, and Eric Freeman of NCDC for World War II UK Royal Navy Data. (8) David Tse of Hong Kong Observatory for Hong Kong data. (9) Volker Wagner for a subset of the Deutscher Wetterdienst HISTOR marine data. (10) Società Meteorologica Italiana (SMI) for the Torino record.

(11) Mac Benoy and the Australian Meteorological Association, Todd Project team. (12) Dick Dee, ECMWF, and the ERA-CLIM project. (13) Manuel Barros, Instituto Geofísico da Universidade do Porto, University of Porto, Vila Nova de Gaia, Portugal. (14) Jonathan Burroughs, Joseph Elms (retired) and Thomas C. Peterson, NOAA National Climatic Data Center, Asheville, NC, USA. (15) Karen Andsager (retired), formerly of Midwestern Regional Climate Center, Cham-

paign, IL USA. (16) Joelle Gergis, University of Melbourne, Australia. (17) Jennifer Hardwick, Met Office Hadley Centre, Exeter, UK. (18) David Jones and Blair Trewin, Bureau of Meteorology, Melbourne, Victoria, Australia. (19) Philip L. Woodworth, Proudman Oceanographic Laboratory, Liverpool, UK. (20) Reinhard Zoellner, Deutscher Wetterdienst, Hamburg, Germany. (21) Dr Birger Tinz, Division Maritime – Climatological Monitoring Centre, Deutscher Wetterdienst, Hamburg, Germany. (22) Ricardo M. Trigo, Instituto Dom Luiz, University of Lisbon, Lisbon, Portugal. (23) Luís Filipe Nunes and Manuel Mendes, Instituto Português do Mar e da Atmosfera, Lisbon, Portugal. (24) Scott D. Woodruff, NOAA Earth System Research Laboratory, Physical Sciences Division, Boulder, CO, USA. (25) Michael Kruk, ERT Inc., Asheville, NC, USA. (26) Leslie Stoecker, Midwestern Regional Climate Center, Champaign, IL, USA. (27) Hui Wan and Val Swail, Environment Canada, Toronto, Canada. (28) Fatou Sima, Division of Meteorology, Department of Water Resources, Banjul, The Gambia. (29) Andrea Lockwood and Philip Mooney, Hollings Scholars, NOAA Earth System Research Laboratory, Boulder, CO, USA. (30) Sam Dean of National Institute of Water and Atmospheric Research (NIWA), Auckland, New Zealand for productive conversations.

The following persons affiliated with the University of Giessen, Germany, digitized subdaily pressure data originating from Europe, Asia, Africa, northern South America, and the Caribbean: Christine Kolbe, Lisa Flander, Marina Ostheimer, Johanna Enghardt, Athanasios Tsikerdekis, Lamprini Dergianli, Despina Xoplaki, Christos Samaras, Eleni Kaimasidou, Nancy Gouta, Elisabet Tsalkitidou, Mary Athanasidou, Christakis Athanasidou, Panagiota Katsaouni, Alexander Theocharis, Stella Dafka, Kevin Pometti, Gail Kelly, Elda Fleitmann, Lisa Theile, Petra Strehlau, Ines Stange, and Jonas Viezens.

References

- Allan R, Brohan P, Compo GP, Stone R, Luterbacher J, Brönnimann S. 2011. The International Atmospheric Circulation Reconstructions over the Earth (ACRE) initiative. *Bulletin of the American Meteorological Society* **92**: 1421–1425, doi:10.1175/2011BAMS3218.1.
- Ansell TJ, Jones PD, Allan RJ, Lister D, Parker DE, Brunet M, Moberg A, Jacobeit J, Brohan P, Rayner NA, Aguilar E, Alexandersson H, Barriendos M, Brandsma T, Cox NJ, Della-Marta PM, Drebs A, Founda D, Gerstengarbe F, Hickey K, Jónsson T, Luterbacher J, Nordli Ø, Oesterle H, Petrakis M, Philipp A, Rodwell MJ, Saladie O, Sigro J, Slonosky V, Srnc L, Swail V, García-Suárez AM, Tuomenvirta H, Wang X, Wanner H, Werner P, Wheeler D, Xoplaki E. 2006. Daily mean sea level pressure reconstructions for the European–North Atlantic region for the period 1850–2003. *Journal of Climate* **19**: 2717–2742, doi:10.1175/JCLI3775.1.
- Brohan P, Allan R, Freeman JE, Waple AM, Wheeler D, Wilkinson C, Woodruff S. 2009. Marine observations of old weather. *Bulletin of the American Meteorological Society* **90**: 219–230, doi:10.1175/2008BAMS2522.1.
- Brönnimann S, Grant AN, Compo GP, Ewen T, Griesser T, Fischer AM, Schraner M, Stickler A. 2012. A multi-data set comparison of the vertical structure of temperature variability and change over the Arctic during the past 100 years. *Climate Dynamics* **39**: 1577–1598, doi:10.1007/s00382-012-1291-6.
- Brunet M, Gilabert A, Jones P, Efthymiadis D. 2014. A historical surface climate dataset from station observations in Mediterranean North Africa and Middle East areas. *Geoscience Data Journal* **1**: 121–128, doi:10.1002/gdj3.12.
- Compo GP, Whitaker JS, Sardeshmukh PD, Matsui N, Allan RJ, Yin X, Gleason BE, Vose RS, Rutledge G, Bessemoulin P, Brönnimann S, Brunet M, Crouthamel RI, Grant AN, Groisman PY, Jones PD, Kruk MC, Kruger AC, Marshall GJ, Maugeri M, Mok HY, Nordli O, Ross TF, Trigo RM, Wang XL, Woodruff SD, Worley SJ. 2011. The Twentieth Century Reanalysis Project. *Quarterly Journal Royal Meteorological Society* **137**: 1–28, doi:10.1002/qj.776.
- Compo GP, Whitaker JS, Sardeshmukh PD, Matsui N, Allan RJ, Yin X, Gleason BE, Vose RS, Rutledge G, Bessemoulin P, Brönnimann S, Brunet M, Crouthamel RI, Grant AN, Groisman PY, Jones PD, Kruk MC, Kruger AC, Marshall GJ, Maugeri M, Mok HY, Nordli O, Ross TF, Trigo RM, Wang XL, Woodruff SD, Worley SJ. 2010. International Surface Pressure Databank (ISPDv2). *Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory, Boulder, Colorado, USA*. doi:10.5065/D6SQ8XDW.
- Dupigny-Giroux L-A, Ross TF, Elms JD, Truesdell R, Doty SR. 2007. RESOURCES - NOAA's Climate Database Modernization Program: rescuing, archiving, and digitizing history. *Bulletin of the American Meteorological Society* **88**: 1015–1017, doi:10.1175/BAMS-88-7-1015.
- Fleming JR. 2000. *Meteorology in America, 1800–1870*. Johns Hopkins University Press: Baltimore, MD; 292.
- Hersbach H, Poli P, Dee D. 2015. The observation feedback archive for the ICOADS and ISPD data sets. ERA Report Series, 18, ECMWF, UK.
- Hirahara S, Masayoshi I, Fukuda Y. 2014. Centennial-scale sea surface temperature analysis and its uncertainty. *Journal of Climate* **27**: 57–75, doi:10.1175/JCLI-D-12-00837.1.
- Kalnay E, Kanamitsu M, Kistler R, Collins W, Deaven D, Gandin L, Iredell M, Saha S, White G, Woollen J, Zhu Y, Leetmaa A, Reynolds R, Chelliah M, Ebisuzaki W, Higgins W, Janowiak J, Mo KC, Ropelewski C, Wang J, Jenne R, Joseph D. 1996. The NCEP/NCAR 40-year reanalysis project. *Bulletin of the American Meteorological Society* **77**: 437–471, doi:10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2.
- Kanamitsu M, Alpert JC, Campana KA, Caplan PM, Deaven DG, Iredell M, Katz B, Pan H-L, Sela J, White GH. 1991. Recent changes implemented into the Global Forecast System at NMC. *Weather Forecasting* **6**: 425–435, doi:10.1175/1520-0434(1991)006%3C0425:RCIITG%3E2.0.CO;2.

- Kistler R, Collins W, Saha S, White G, Woollen J, Kalnay E, Chelliah M, Ebisuzaki W, Kanamitsu M, Kousky V, van den Dool H, Jenne R, Fiorino M. 2001. The NCEP-NCAR 50-year reanalysis: monthly means CD-ROM and documentation. *Bulletin of the American Meteorological Society* **82**: 247–267, doi:10.1175/1520-0477(2001)082<0247:TNNYRM>2.3.CO;2.
- Knapp KR, Kruk MC, Levinson DH, Diamond HJ, Neumann CJ. 2010. The International Best Track Archive for Climate Stewardship (IBTrACS): unifying tropical cyclone best track data. *Bulletin of the American Meteorological Society* **91**: 363–376, doi:10.1175/2009BAMS2755.1.
- Maugeri M, Brunetti M, Monti F, Nanni T. 2004. Sea-level pressure variability in the Po plain (1765–2000) from homogenized daily secular records. *International Journal of Climatology* **24**: 437–455, doi:10.1002/joc.991.
- Maugeri M, Lentini G, Brunetti M, Nanni T. 2008. Availability and quality of Italian secular meteorological records and consistency of still unexploited early data. In *MEDARE—Proceedings of the International Workshop on Rescue and Digitization of Climate Records in the Mediterranean Basin*, Brunet M, Kuglitsch FG (eds). World Meteorological Organization, WCDMP-67, WMO/TD No. 1432, June 2008; 61–70.
- Moorthi S, Pan H-L, Caplan P. 2001. Changes to the 2001 NCEP operational MRF/AVN global analysis/forecast system. Technical Procedures Bulletin 484, NOAA, NWS: Silver Spring, MD. <http://www.nws.noaa.gov/om/tpb/484.htm> (accessed 3 June 2010).
- Parker D, Kent E, Woodruff S, Dehennauw D, Harrison DE, Manabe T, Mietus M, Swail V, Worley S. 2004. The Second JCOMM Workshop on Advances in Marine Climatology (CLIMAR-II). *WMO Bulletin* **53**: 157–159.
- Poli P, Hersbach H, Tan D, Dee D, Thépaut J-N, Simmons A, Peubey C, Laloyaux P, Komori T, Berrisford P, Dragani R, Trémolet Y, Hólm E, Bonavita M, Isaksen L, Fisher M. 2013. The data assimilation system and initial performance evaluation of the ECMWF pilot reanalysis of the 20th-Century assimilating surface observations only (ERA-20C). ERA Report Series, 14, ECMWF, UK.
- Rayner A, Parker DN, Horton EE, Folland CB, Alexander LK, Rowell DV, Kent EP, Kaplan CA. 2003. Global analyses of sea surface temperature, sea ice, night marine air temperature since the late nineteenth century. *Journal of Geophysical Research* **108** (D14): 4407, doi:10.1029/2002JD002670.
- Saha S, Nadiga S, Thiaw C, Wang J, Wang W, Zhang Q, van den Dool HMPan H-L, Moorthi S, Behringer D, Stokes D, Peña M, Lord S, White G, Ebisuzaki W, Peng P, Xie P. 2006. The NCEP climate forecast system. *Journal of Climate* **19**: 3483–3517, doi:10.1175/JCLI3812.1.
- Schmith T, Alexandersson H, Iden K, Tuomenvirta H. 1997. North Atlantic-European pressure observations 1868–1995 (WASA dataset version 1.0). Technical Report 97-3, Danish Meteorological Institute, Copenhagen, Denmark, 13. http://www.dmi.dk/fileadmin/user_upload/Rapporter/TR/1997/tr97-3.pdf (accessed 16 February 2001).
- Schmith T, Kaas E, Li T-S. 1998. Northeast Atlantic winter storminess 1875–1995 re-analysed. *Climate Dynamics* **14**: 529–536, doi:10.1007/s003820050239.
- Smith A, Lott N, Vose RS. 2011. The Integrated Surface Database: recent developments and partnerships. *Bulletin of the American Meteorological Society* **92**: 704–708, doi:10.1175/2011BAMS3015.1.
- Uppala SM, Kållberg PW, Simmons AJ, Andrae U, Bechtold VDC, Fiorino M, Gibson JK, Haseler J, Hernandez A, Kelly GA, Li X, Onogi K, Saarinen S, Sokka N, Allan RP, Andersson E, Arpe K, Balmaseda MA, Beljaars ACM, Berg LVD, Bidlot J, Bormann N, Caires S, Chevallier F, Dethof A, Dragosavac M, Fisher M, Fuentes M, Hagemann S, Hólm E, Hoskins BJ, Isaksen L, Janssen PAEM, Jenne R, McNally AP, Mahfouf J-F, Morcrette J-J, Rayner NA, Saunders RW, Simon P, Sterl A, Trenberth KE, Untch A, Vasiljevic D, Viterbo P, Woollen J. 2005. The ERA-40 re-analysis. *Quarterly Journal of the Royal Meteorological Society* **131**: 2961–3012, doi:10.1256/qj.04.176.
- Valente MA, Trigo R, Barros M, Nunes LF, Alves EI, Pinhal E, Coelho FES, Mendes M, Miranda JM. 2008. Early stages of the recovery of Portuguese historical meteorological data. In *MEDARE—Proceedings of the International Workshop on Rescue and Digitization of Climate Records in the Mediterranean Basin*, Brunet M, Kuglitsch FG (eds). World Meteorological Organization, WCDMP-67, WMO/TD No. 1432, June 2008; 95–102.
- Wang XL, Feng Y, Compo GP, Zwiers FW, Allan RJ, Swail VR, Sardeshmukh PD. 2014. Is the storminess in the Twentieth Century Reanalysis really inconsistent with observations? A reply to the comment by Krueger *et al.* (2013b). *Climate Dynamics* **42**: 1113–1125, doi:10.1007/s00382-013-1828-3.
- Woodruff SD, Diaz HF, Elms JD, Worley JS. 1998. COADS Release 2 data and metadata enhancements for improvements of marine surface flux fields. *Physics and Chemistry of the Earth* **23**: 517–526.
- Woodruff SD, Diaz HF, Worley SJ, Reynolds RW, Lubker SJ. 2005. Early ship observational data and ICOADS. *Climatic Change* **73**: 169–194, doi:10.1007/s10584-005-3456-3.
- Woodruff SD, Worley SJ, Lubker SJ, Ji Z, Eric Freeman J, Berry DI, Brohan P, Kent EC, Reynolds RW, Smith SR, Wilkinson C. 2011. ICOADS Release 2.5: extensions and enhancements to the surface marine meteorological archive. *International Journal of Climatology* **31**: 951–967, doi:10.1002/joc.2103.
- Worley SJ, Woodruff SD, Reynolds RW, Lubker SJ, Lott N. 2005. ICOADS release 2.1 data and products. *International Journal of Climatology* **25**: 823–842, doi:10.1002/joc.1166.
- Yin X, Gleason BE, Compo GP, Matsui N, Vose RS. 2008. The International Surface Pressure Databank (ISPD) land component version 2.2. National Climatic Data Center, Asheville, NC, 1–12. ftp://ftp.ncdc.noaa.gov/pub/data/ispd/doc/ISPD2_2.pdf (accessed 19 August 2010).