

Assessing alcohol consumption through wastewater-based epidemiology: Spain as a case study

Ester López-García¹, Carlos Pérez-López¹, Cristina Postigo^{1*}, Vicente Andreu², Lubertus Bijlsma³, Iria González-Mariño^{4,5}, Félix Hernández³, Rosa Maria Marcé⁶, Rosa Montes⁴, Yolanda Picó², Eva Pocurull⁶, Andreu Rico⁷, Rosario Rodil⁴, María Rosende⁸, Yolanda Valcárcel⁹, Olatz Zuloaga¹⁰, José Benito Quintana⁴, Miren López de Alda^{1*}

¹Water, Environmental, and Food Chemistry Unit (ENFOCHEM), Department of Environmental Chemistry, Institute of Environmental Assessment and Water Research (IDAEA-CSIC), Barcelona

²Food and Environmental Safety Research Group (SAMA-UV), Desertification Research Centre (CIDE), CSIC-Generalitat Valenciana-University of Valencia, Valencia

³Environmental and Public Health Analytical Chemistry, Research Institute for Pesticides and Water, University Jaume I, Castellón

⁴Department of Analytical Chemistry, Nutrition and Food Sciences, Institute of Research on Chemical and Biological Analysis (IAQBUS), Universidade de Santiago de Compostela, Santiago de Compostela

⁵Department of Analytical Chemistry, Nutrition and Bromatology, Faculty of Chemical Sciences, University of Salamanca, Salamanca.

⁶Department of Analytical Chemistry and Organic Chemistry, Universitat Rovira i Virgili, Tarragona

⁷IMDEA Water Institute, Science and Technology Campus of the University of Alcalá, Alcalá de Henares

⁸FI-TRACE group, Department of Chemistry, University of the Balearic Islands, Palma de Mallorca

⁹Research Group in Environmental Toxicology and Risk Assessment (TAyER), Medical specialties and Public Health, Faculty of Health Sciences, Rey Juan Carlos University, Madrid

¹⁰Department of Analytical Chemistry, Faculty of Science and Technology (UPV/EHU) & Plentzia Marine Station of Basque Country University (EHU/UPV), Basque Country

***Corresponding Authors:** Cristina Postigo; Miren López de Alda
Institute of Environmental Assessment and Water Research
(IDAEA-CSIC)
Department of Environmental Chemistry
C/ Jordi Girona 18-26, 08034 Barcelona, Spain.
cprqam@cid.csic.es; miren.lopezdealda@idaea.csic.es
Tel: +34-934-006-100, Fax: +34-932-045-904

Abstract

Background: In this study, an alternative and complementary method to those approaches currently used to estimate alcohol consumption by the population is described. This method, known as wastewater-based epidemiology (WBE), allows back-calculating the alcohol consumption rate in a given population from the concentrations of a selected biomarker measured in wastewater.

Methods: Composite (24-h) wastewater samples were collected at the inlet of 17 wastewater treatment plants located in 13 Spanish cities for seven consecutive days in 2018. The sampled area covered 12.8% of the Spanish population. Wastewater samples were analyzed to determine the concentration of ethyl sulfate, the biomarker used to back-calculate alcohol consumption.

Results: Alcohol consumption ranged from 4.5 to 46 mL/day/inhabitant. Differences in consumption were statistically significant among the investigated cities and between weekdays and weekends. WBE-derived estimates of alcohol consumption were comparable to those reported by its corresponding region in the Spanish National Health Survey in most cases. At the national level, comparable results were obtained between the WBE-derived annual consumption rate (5.7 ± 1.2 L ethanol per capita (aged 15+)) and that reported by the National Health Survey (4.7 L ethanol per capita (aged 15+)).

Conclusions: This is the largest WBE study carried out to date in Spain to estimate alcohol consumption rates. It confirms that this approach is useful for establishing spatial and temporal patterns of alcohol consumption, which could contribute to the development of health care management plans and policies. Contrary to established methods, it allows obtaining information in a fast and relatively economical way.

58

59 *Keywords:* sewage epidemiology, alcohol abuse, liquid chromatography-mass spectrometry,

60 consumption patterns,

61

62

64 1. Introduction

65 In 2016, the consumption of alcohol was responsible for 3 million deaths worldwide and it
66 became one of the main health risk factors for the population, being more harmful than digestive
67 diseases, road injuries, diabetes, or violence (World Health Organization (WHO), 2018). In
68 Spain, alcohol is the psychoactive substance most consumed (Observatorio Español de las
69 Drogas y las Adicciones (OEDA), 2019). In 2017 (last reported year), 91% of the Spanish
70 population aged 15-64 years had consumed alcohol at some point in their lifetime, while 75%
71 had consumed alcohol in the last year, and 63% did it in the last month. Overall, the
72 consumption by men is higher than by women and the average age at which alcohol begins to
73 be consumed is 16.6 years (OEDA, 2019). According to the 2018's Global status report on
74 alcohol and health provided by the WHO, the annual intake of alcohol in Spain in 2016 was 10
75 L of pure alcohol per capita (aged 15+), which is similar to the European average (9.8 L) (WHO,
76 2018). These estimates are traditionally obtained from population surveys, recorded alcohol
77 data (alcohol taxation or sales), and unrecorded alcohol data (homemade or informally
78 produced alcohol, smuggled alcohol, alcohol for industrial or medical uses, alcohol obtained
79 through cross-border shopping, or surrogate alcohol) (WHO, 2018). Through surveys,
80 consumption figures can be disaggregated for specific population groups by age or gender.
81 However, the use of these tools/data to derive alcohol consumption figures is time-consuming
82 and relatively expensive, and consequently, it does not allow obtaining real-time estimates (i.e.,
83 consumption data in Spain are given with a delay of two years). Furthermore, the data obtained
84 by surveys may not be representative of actual population consumption due to misreporting of
85 alcohol consumption by survey participants (Stockwell et al., 2016; van Wel et al., 2016) or to
86 inaccurate estimates of unrecorded alcohol (Probst et al., 2019). Therefore, it is necessary to
87 propose alternative approaches that provide quick and precise information and that, together

with the traditional ones, can help to obtain a more reliable picture of alcohol consumption rates.

Wastewater-based epidemiology (WBE) is a novel approach that has been applied in the last decade to estimate illicit drug use at the city level (González-Mariño et al., 2019). The European Monitoring Centre for Drugs and Drug Addiction (EMCDDA) has adopted it, indeed, as a complementary indicator to established methods for illicit drug use estimation (EMCDDA, 2016). The WBE approach is based on the fact that, after consumption, the substances are excreted via urine and feces, either unaltered or as a metabolite, and conducted through the sewage network to a wastewater treatment plant (WWTP). Thus, a raw wastewater sample contains specific biomarkers of the drugs that can be used to back-calculate the amount of substance that has been consumed. In the case of alcohol, after human consumption, about 95% is metabolized in the liver via oxidation to acetaldehyde and acetic acid, about 5% is excreted unaltered, and a small part (<0.1%) is excreted as ethyl sulfate (EtS) and ethyl glucuronide (EtG) after conjugation with sulfate and glucuronic acid, respectively. EtS and EtG can be detected in urine after 1 hour of alcohol intake (Helander and Beck, 2005), so they have been proposed as good indicators for recent alcohol consumption. However, only EtS is stable in wastewater (Rodríguez-Álvarez et al., 2014) and its occurrence in wastewater is exclusively due to alcohol consumption and not to the metabolism of unaltered alcohol by endogenous bacteria (Reid et al., 2011). Thus, EtS has been pointed out as the best biomarker to estimate alcohol consumption through WBE.

WBE was first applied to estimate alcohol consumption in 2011 in Oslo (Norway) (Reid et al., 2011) and, since then, many studies have been carried out in cities from other European countries (Andrés-Costa et al., 2016; Baz-Lomba et al., 2016; Gatidou et al., 2016; Mastroianni et al., 2014, 2017; Rodríguez-Álvarez et al., 2014, 2015; van Wel et al., 2016) Vietnam (Nguyen et al., 2018), China (Gao et al., 2020), United States (Chen et al., 2019), Canada (Ryu et al.,

2016), and Australia (Zheng et al., 2020). The main objective of these studies was not only to investigate spatial differences of alcohol consumption between populations or to assess changes in alcohol consumption due to special events (Andrés-Costa et al., 2016) but also, to compare WBE-derived alcohol estimates with alcohol consumption figures obtained using traditional methods, such as official data provided by the WHO or by national surveying institutions. In these studies, the alcohol consumption rates were estimated from data gathered from a single WWTP, which only serves a city or part of it, after a sampling period of one week in most of the cases, except for Milan and Santiago (Rodríguez-Álvarez et al., 2015), Oslo (Reid et al., 2011), Lied (Belgium) (van Wel et al., 2016), U.S (Chen et al., 2019) and Australia (Zheng et al., 2020), for which longer sampling periods were used (namely, 2 weeks, 3 weeks, four-two weeks periods, one weekday every month during eleven months, and one week every two months during 6 years, respectively). To date, only three studies have conducted nation-wide investigations by collecting samples from different WWTPs: a study conducted in Australia, in which 18 WWTPs were sampled, covering 45% of the whole population (Lai et al., 2018); one carried out in Belgium, which covered 8 WWTPs and 12.8% of the total population (Boogaerts et al., 2016); and another one in China, which included 48 WWTPs and 3.3% of the whole population (Gao et al., 2020).

The present study is one of the few nation-wide applications of WBE to estimate alcohol consumption rates, and the largest conducted so far in Spain. Wastewater samples were analyzed from 17 WWTPs, covering 12.8% of the Spanish population. The specific objectives of this work were: i) to assess spatial differences in alcohol consumption between the different investigated areas in Spain, ii) to assess weekly consumption patterns, and iii) to extrapolate the estimated alcohol consumption in the studied areas to the whole Spanish population, and to compare it with official data reported by the WHO or national institutions.

2. Material and methods

2.1. Reagents

Analytical standards of ethyl sulfate (EtS) and its isotopically labeled compound, EtS-d₅, were obtained as EtS sodium salt and ethyl-d₅ sulfate salt from Cerilliant (Round Rock, TX, USA) as solutions in methanol (MeOH) at a concentration of 1 mg/mL. Water and MeOH, both HPLC-grade, and acetic acid (98% purity) used as a mobile phase modifier, were purchased from Merck (Darmstadt, Germany). Dibutylamine (>99.5% purity), also used as a mobile phase modifier, was obtained from Sigma Aldrich (Steinheim, Germany).

2.2. Standard solutions

Stock standard solutions were prepared at different concentrations in the range of 10 to 20,000 µg/L by appropriate dilution of the commercial EtS standard in MeOH, with a constant concentration of EtS-d₅ of 2,500 µg/L, and were stored in the dark at -20°C until analysis. Before analysis, working standard solutions were freshly prepared by dilution of these stock standard solutions in HPLC water (1:100, v/v).

2.3. Sample collection and preparation

Influent wastewater samples were collected from 17 WWTPs located in 13 Spanish cities that belong to 7 out of the 17 regions of Spain. Figure 1 shows the location of the sampled WWTPs. The sampling covers populations of various sizes (i.e., between 47,961 and 1,163,154 inhabitants). In total, the population reached with the sampling was 5,981,848 inhabitants, which corresponds to 12.8% of the Spanish population. The cities sampled were Barcelona,

160 Bilbao, Castellón, Guadalajara, Lleida, Madrid, Móstoles, Palma de Mallorca, Reus, Santiago
161 de Compostela, Tarragona, Toledo, and Valencia, including in some cases part of their
162 metropolitan area. Except for Barcelona, Madrid, and Móstoles, where WWTPs only covered
163 35, 30, and 90 % of their total population, respectively, all other main cities were fully covered
164 (100% of their population). Table 1 shows the populations served by each WWTP as well as
165 the sampling protocol carried out in each of them.

166 From each WWTP, 24-h composite influent wastewater samples were collected during seven
167 consecutive days in the spring of 2018 using time or flow proportional techniques (Table 1).
168 The sampling was conducted during a “normal week” so that special events such as holidays or
169 festivals were avoided. After collection, samples were immediately stored at -20°C. They were
170 sent frozen by courier in less than 24 hours to the laboratory in Barcelona, where all samples
171 were analyzed. Once in the laboratory, an aliquot of 10 mL was spiked with EtS-d₅ at a
172 concentration of 25 µg/L and 1 mL of this sample was transferred to a 1.5 mL microcentrifuge
173 tube and centrifuged at 10,000 rpm for 10 minutes at a temperature of 4°C (Eppendorf 5810R,
174 Hamburg, Germany). Then, the supernatant was transferred to a glass vial and stored at -20°C
175 in the darkness until its analysis by liquid chromatography coupled to tandem mass
176 spectrometry (LC-MS/MS).

177 **Table 1.** Description of sampled WWTPs (name, population served and locations/districts covered with main city in bold) and the sampling
178 protocol carried-out (location of autosampler, and sampling mode, start time and period).

Regions	City ^a	WWTP name	Population served by the WWTPs	Method used to estimate the population served ^b	Locations/districts served by the WWTPs	Percentage of the main city covered by the WWTP ^c	Location of autosampler	Sampling mode ^d	Sampling start time	Sampling period
Balearic Islands	Palma de Mallorca	Palma I	406,492	Census 2017	Palma beach, Sant Jordi, El Pil·lari, Son Sant Joan airport, part of Palma de Mallorca	100	After fine screen	T (100 mL/ 15 min)	10:00	10/04/2018 - 16/04/2018
		Palma II	47,961	Census 2017	Palma de Mallorca (main part), Marratxí, Esporles, Bunyola and Son Castelló, Can Valero, Son Rosinyol industrial states		After fine screen	T (100 mL/ 15 min)	10:00	18/04/2018 - 24/04/2018
Basque Country	Bilbao	Galindo	860,237	Census 2016	Abanto-Zierbena, Alontotegi, Arrigorriaga, Barakaldo, Barrika, Basauri, Berango, Bilbao, Derio, Erandio, Etxebarri, Galdakao, Getxo, Leioa, Lezama, Loiu, Ortuella, Portugalete, Santurtzi, Sestao, Sondika, Sopelana, Trapagaran, Ugao-Miravalles, Urduliz, Zamudio, Zaratamo, Zeberio	100	After coarse screens and pumping	T (100mL/ 60 min)	8:00	17/04/2018 - 23/04/2018
Castilla-La Mancha	Toledo	Estiviel	79,793	Average BOD April-May 2018	Toledo	100	After sieving	T (100 mL/ 15 min)	8:00	17/04/2018 - 23/04/2018

	Guadalajara	Guadalajara	94,755	Average BOD Jan-April 2018	Guadalajara	100	Before fine screen	T (200 mL/ 60 min)	10:00	02/05/2018 - 08/05/2018
Catalonia	Barcelona	Baix Llobregat	1,163,154	Census 2017	Barcelona, Cervelló, Cornellà de Llobregat, Esplugues de Llobregat, Hospitalet de Llobregat, El Prat de Llobregat, Sant Boi de Llobregat, San Joan Despí, San Just Desvern	35	Mechanical bar screens	T (50 mL/ 10min)	9:00	14/03/2018 - 20/03/2018
	Lleida	Lleida	143,612	Census 2017	Lleida , Alpicat	100	Before fine screen	T (200 mL/ 60 min)	6:00	07/03/2018 - 13/03/2018
	Reus	Reus	115,000	Census 2017	Reus , Castellvell, Almóster	100	After fine screen	F	20:00	17/04/2018 - 23/04/2018
	Tarragona	Tarragona	142,635	Census 2017	Tarragona , La Canonja, Els Pallaresos	100	Before fine screen	T (450 mL/ 60 min)	8:00-9:00	17/04/2018 - 23/04/2018
Commun ity of Madrid	Madrid	Madrid- Centre	727,176	Average COD for the sampling period	Madrid-Center (Neighborhoods: Chamartín, Tetuán, Moncloa-Aravaca, Chamberí, Centro, Arganzuela, Retiro, Ciudad Lineal, Salamanca, Moratalaz, Puente de Vallecas).	30	After sieving	T (400 mL/ 30 min)	8:00	16/05/2018 - 22/05/2018
	Madrid	Madrid- North	227,869	Average BOD 2016 (with 60 g BOD/d)	Pozuelo y Madrid- North: (Neighborhoods: Chamartín, Tetuán, Moncloa, Aravaca, Fuencarral, El Pardo, Las Rozas, Majadahonda)		After fine screen	T (100 mL/ 60 min)	8:00	20/06/2018 - 26/06/2018

	Móstoles	El Soto	187,281	H x 3.5 (WWTP recomm.)	Móstoles, Alcorcón, Fuenlabrada	90	After fine screen	T (100 mL/ 60 min)	8:00	17/05/2018 - 23/05/2018
Galicia	Santiago de Compostela	Silvouta	136,500	H x 2.5 (WWTP recomm.)	Santiago de Compostela	100	After fine screen	T (150 mL/ 10 min)	9:00	13/03/2018 - 19/03/2018
Communitiy of Valencia	Castellón	Castellón de la Plana	171,669	Census 2015	Castellón	100	Before fine screen	T (100 mL/ 15 min)	8:30	11/04/2018 - 17/04/2018
	Valencia	Pinedo I (Valencia-PI)	527,222	COD	Valencia (main part)	100	After fine screen	T (100 mL/ 60 min)	8:00	10/04/2018 - 16/04/2018
	Valencia	Pinedo II (Valencia-P II)	788,242	COD	Albal, Alcàsser, Alfafar, Benetúser, Beniparrell, Burjassot, Catarroja, Llocnou de la Corona, Massanassa, Mislata, Paiporta, Paterna, Picanya, Picassent, Sedaví, Silla, Torrent, part of Valencia		After fine screen	T (100 mL/ 60 min)	8:00	10/04/2018 - 16/04/2018
	Valencia	Quart-Benager (Valencia-QB)	162,249	COD	Alaquàs, Aldaia, Manises, Mislata, Quart de Poblet, Xirivella		After fine screen	F	8:00	10/04/2018 - 16/04/2018

^aName of the main city served by the WWTPs (some WWTPs receive wastewater from other towns included in the capital metropolitan area). ^bBOD: Biochemical Oxygen Demand; COD: Chemical Oxygen Demand; H: Number of homes connected to the sewage system. WWTP recomm: following WWTP recommendations. ^cWWTPs serving parts of the same main city were considered all together for this calculation. ^dT: time-proportional (volume sampled/frequency of sampling); F: Flow-proportional



187 **Figure 1.** Map of Spain with the location of the sampled WWTPs (regions are indicated in
188 different colors).

2.4. Sample analysis

The analysis of EtS was performed with a previously described and validated methodology based on ion-pair LC-MS/MS (Mastroianni et al., 2014) using a SymbiosisTM Pico System (Spark Holland, Emmen, The Netherlands) equipped with a 100 µL sample loop. The LC system was coupled to a 4000QTRAP hybrid triple quadrupole-linear ion trap (QqLIT) mass spectrometer equipped with a Turbo Ion Spray source (AB-Sciex, Foster City, CA, USA) set in the negative ionization mode (ESI⁻). Chromatographic separation was performed with a Purospher Star RP-18 end-capped column (125 mm × 2 mm, particle size 5 µm) preceded by a guard column of the same packing material and particle size, both from Merck (Darmstadt, Germany) and a mobile phase consisting of MeOH and water both containing 5 mM of dibutylammonium acetate (DBAA) at a constant flow rate of 0.3 mL/min. MS/MS detection was performed in selected reaction monitoring mode (SRM) recording 2 SRM transitions for EtS (125→97, 125→80) and one for EtS-d₅ (130→98). Data acquisition and evaluation was performed with Analyst 1.5 software (AB-Sciex, Foster City, CA, USA). Quantification of the samples was based on the isotope dilution method.

2.5. Quality control and quality assurance

A calibration curve was freshly prepared in water for the analysis of each batch of samples in the range 0.1-200 µg/L. For this, appropriate amounts of stock standard solutions were fortified in water and processed following the sample treatment protocol. The calibration curve was injected at the beginning and the end of each batch of samples, and calibration curves were constructed with the average response, using weighted least square regression models ($1/x^2$ as weight) to reduce the effect of high concentrations in the model. Only those calibration

solutions that did not deviate more than 20% from the theoretical concentration were used to construct the model.

Quality controls, i.e., a standard solution containing EtS and EtS-d5 at concentrations of 5 µg/L and 25 µg/L, respectively, were injected every 6 samples to check the correct operation of the instrument. MS signals for EtS were absent in solvent blanks (HPLC-grade water injected every 3 samples) and method blanks (HPLC-grade water processed following the sample treatment protocol and thus, fortified with EtS-d5 at a concentration of 25 µg/L). Therefore, analyte carryover between injections and cross-contamination during sample preparation could be discarded.

2.6. Alcohol consumption estimates

Back calculation of alcohol consumption was made according to the following equation:

$$\frac{mL\ EtOH}{day * inhabitant} = C_{EtS} \left[\frac{\mu g}{L} \right] * 10^{-6} \left[\frac{g}{\mu g} \right] * Q \left[\frac{m^3}{day} \right] * 10^3 \left[\frac{L}{m^3} \right] * \frac{1}{P} * 3047 * \frac{1}{\rho_{EtOH} \left(\frac{g}{mL} \right)}$$

where C_{EtS} is the concentration of EtS measured in the wastewater sample, Q is the water flow entering the WWTP, P is the total population served by the WWTP (Table 1), 3047 is the correction factor applied which takes into account the molar mass ratio between ethanol (MW: 46.07 g/mol) and EtS (MW: 126.13 g/mol) and the excretion rate of EtS in urine (0.012%) (Rodríguez-Álvarez et al., 2015), and ρ_{EtOH} is ethanol density (0.789 g/mL).

2.7. Statistical data analysis

Data were statistically analyzed to compare alcohol consumption rates between populations, regions, weekdays, and weekends, and between populations grouped according to their size (above or below 300,000 inhabitants). Since data were not normally distributed (after Shapiro

Wilk test, $p\text{-value} < 0.05$) and/or the sample size was too small ($n < 10$) in some cases, non-parametric tests were applied. The Mann-Whitney U test was used to compare two independent samples, whereas the Kruskal-Wallis test was used to compare three or more individual groups. If the latter revealed significant differences among groups, they were subsequently investigated after applying the Mann-Whitney U test to every two populations. False Discovery Rate (FDR) correction for multiple testing was applied to reduce the number of “false positives”. Spearman correlation test was also applied to assess the correlation between WBE-derived data and those reported by established indicators. All the analyses were done using the software R (version R 3.5.3) and considering a 95% confidence level ($\alpha = 0.05$).

3. Results

3.1. Occurrence of EtS in wastewater samples and alcohol consumption estimations

Table 2 shows the concentrations of EtS, the mass loads of EtS that reached each WWTP and the estimated alcohol consumption in each investigated area, expressed as average, median and range; whereas Figure 2 depicts alcohol consumption in the form of boxplots by each investigated population in the various considered regions. EtS was found in all samples above LOQ ($0.07 \mu\text{g/L}$) at concentrations ranging from $1.4 \mu\text{g/L}$ (Santiago de Compostela) to $74 \mu\text{g/L}$ (Tarragona). The average weekly concentrations of EtS ranged from 2.9 to $43 \mu\text{g/L}$, with the lowest values being found in the WWTPs that serve the cities of Santiago de Compostela, Lleida, and Guadalajara (below $10 \mu\text{g/L}$) and the highest values in the WWTPs that serve Móstoles ($31 \mu\text{g/L}$) and Tarragona ($43 \mu\text{g/L}$). The average weekly levels of EtS measured in the remaining WWTPs were between 11 (Toledo) and $21 \mu\text{g/L}$ (Reus).

The alcohol consumption estimated from levels of EtS in the analyzed samples ranged from 4.5 (Santiago de Compostela) to 46 mL/day/inhabitant (Tarragona). The cities with the highest

average alcohol consumption were Tarragona, Bilbao, and Móstoles, with average weekly consumption of 27, 20, and 17 mL/day/inhabitant, respectively. The lowest average alcohol consumptions (<10 mL/day/inhabitant) were estimated in Toledo (7.4), Santiago de Compostela (8.4), Lleida (8.5), Madrid-Centre (8.9), Castellón (9.0), and Valencia-QB (9.4). In the remaining investigated areas (Guadalajara, Barcelona, Reus, Madrid-North, Valencia-PI, Valencia-PII, and Palma de Mallorca), average alcohol consumption was between 11 and 14 mL/day/inhabitant.

Comparing with previous studies conducted in Spain, similar alcohol consumption rates were previously reported in Barcelona (18 mL/day/inhabitant) (Mastroianni et al., 2014) and Castellón (6.6 mL/day /inhabitant) (Baz-Lomba et al., 2016), and higher in Santiago de Compostela (13.6-16.3 mL/day/inhabitant) (Rodríguez-Álvarez et al., 2015, 2014). On the contrary, the alcohol consumption estimated during a normal week in Valencia (Valencia-PI (6.2 mL/day/inhabitant (aged 15+)), Valencia-PII (3.3 mL/day/inhabitant (aged 15+)) and Valencia-QB (5.9 mL/day/inhabitant (aged 15+)) was lower than that estimated in the present study, even though consumption figures in that study were obtained considering only the population aged 15+ (Andrés-Costa et al., 2016).



Comparing with other international studies, the estimated rates in the investigated Spanish populations (average alcohol consumption from 7.4 to 27 mL/day/inhabitant), were similar to those reported by other investigated cities (Table 3) except in Ho Chin Minh (Vietnam) (Nguyen et al., 2018), Lesvos (Greece) (Gatidou et al., 2016), Milan (Italy) (Baz-Lomba et al., 2016; Rodríguez-Álvarez et al., 2015) and Lugano (Switzerland) (Ryu et al., 2016), where alcohol consumption rates (from 3.4 to 6.6 mL/day/inhabitant) were lower than those estimated for Spanish populations. On the contrary, Copenhagen (Denmark) and Granby (Canada) (Ryu et al., 2016), showed higher alcohol consumption rates, 40 and 44 mL/day/inhabitant, respectively.

285 **Table 2.** Frequency of detection of EtS (%), EtS concentration ($\mu\text{g/L}$), EtS load (mg/day/inhabitant) and alcohol consumption (mL/day/inhabitant)
286 in the investigated cities (expressed as average, median and range).

	Freq. (%)	Concentration ($\mu\text{g/L}$)			EtS load (mg/day/inhabitant)			Alcohol (mL/day/inhabitant)				
		Average	Median	Range	Average	Median	Range	Average	Median	Range	Average weekdays	Average weekend
Palma I	100	15	15	11-21	-	-	-	-	-	-	-	-
Palma II	100	18	16	14-26	-	-	-	-	-	-	-	-
Palma de Mallorca ^a	-	-	-	-	3492	3221	2581-4702	14	12	10-18	12	17
Bilbao	100	17	16	18-29	5133	4867	3906-7632	20	19	15-30	19	23
Guadalajara	100	9.3	7.8	6.5-15	2857	2499	2051-4417	11	9.7	7.9-17	9.0	16
Toledo	100	11	9.1	7.8-19	1926	1555	1426-3007	7.4	6.0	5.5-12	5.8	11
Barcelona	100	16	14	5.9-25	3455	3021	2030-5352	13	12	7.8-21	11	20
Lleida	100	7.4	6.9	5.6-10	2208	1807	1663-3333	8.5	7.0	6.4-13	7.2	12
Reus	100	21	13	12-39	3081	2036	1814-5363	12	7.9	7.0-21	8.8	20
Tarragona	100	43	50	11-74	7091	8597	1935-11906	27	33	7.5-46	27	28
Madrid-Centre	100	15	15	9.4-23	2301	2175	1381-3431	8.9	8.4	5.3-13	7.6	12
Madrid-North	100	18	17	9.4-26	3375	3342	1719-5327	13	13	6.6-21	13	14
Móstoles	100	31	28	18-50	4430	4147	2592-7520	17	16	10-29	15	22
Santiago de Compostela	100	2.9	2.7	1.4-4.4	2178	2197	1173-3124	8.4	8.5	4.5-12	7.0	12
Castellón	100	12	11	7.3-23	2325	1964	1635-4101	9.0	7.6	6.3-16	7.4	13
Valencia-PI	100	13	13	7.5-19	2977	2829	1722-4364	12	11	6.6-17	9.6	16
Valencia-PII	100	12	11	6.9-19	2957	3282	2168-3483	11	13	8.4-13	11	13
Valencia-QB	100	14	11	10-22	2438	2339	1693-3770	9.4	9.0	6.5-15	8.0	13

287 ^aDuring sampling period Palma I derived part of its water flow to Palma II, so to calculate EtS load and to estimate alcohol consumption, Palma I
288 and Palma II were jointly treated as Palma de Mallorca.

289



290

291 **Figure 2.** Distribution of alcohol consumption among investigated populations (Figure 2a) and regions (Figure 2b). (In Figure 2a, populations
292 belonging to the same region are shown between vertical lines; * Outlier).

293 **Table 3.** Alcohol consumption rates estimated by means of WBE approach in different cities
 294 worldwide.

City (Country)	Alcohol consumption (mL/day/inhabitant)		Year	Reference
	Average	Range		
Ho Chi Minh (Vietnam)	3.1-3.9		2015	(Nguyen et al., 2018)
Lesvos (Greece)	3.4/5.4	1.7-7.2/2.2-11.2	2015	(Gatidou et al., 2016)
Valencia-PII (Spain)	3.3 ^a	1.1-6.4 ^a	2014	(Andrés-Costa et al., 2016)
Milan (Italy)	5.1	3.2-10.5	2012- 2014	(Rodríguez-Álvarez et al., 2015)
	6.4	5.1-8.1	2014	(Ryu et al., 2016)
	6.6		2015	(Baz-Lomba et al., 2016)
Valencia-QB (Spain)	5.9 ^a	3.3-12.8 ^a	2014	(Andrés-Costa et al., 2016)
Valencia-PII ^b (Spain)	6.1 ^a	4.3-9.1 ^a	2014	(Andrés-Costa et al., 2016)
Valencia-PI (Spain)	6.2 ^a	1.1-18.31 ^a	2014	(Andrés-Costa et al., 2016)
Lugano (Switzerland)	6.5	4.5-8.4	2014	(Ryu et al., 2016)
Toowoomba (Australia)	9.7	6.9-14.5	2014	(Ryu et al., 2016)
Utrecht (The Netherlands)	10.8		2015	(Baz-Lomba et al., 2016)
	12.9	7.7-20.7	2014	(Ryu et al., 2016)
Santiago de Compostela (Spain)	13.6	3.8-22.6	2012- 2014	(Rodríguez-Álvarez et al., 2015)
	16.3	9.3-23.5	2012	(Rodríguez-Álvarez et al., 2014)
Valencia-PII ^b	14.4 ^a	4.9-23.8 ^a	2014	(Andrés-Costa et al., 2016)
Almada (Portugal)	14.6	8.4-24.1	2014	(Ryu et al., 2016)
Canberra (Australia)	14.6	9.3-22.3	2014	(Ryu et al., 2016)
Zurich (Switzerland)	14.7		2015	(Baz-Lomba et al., 2016)
Bristol (The United Kingdom)	16.2		2015	(Baz-Lomba et al., 2016)
Berlin (Germany)	16.9	13.8-22.3	2014	(Ryu et al., 2016)
Oslo (Norway)	16.1		2009	(Reid et al., 2011)
	18.9		2015	(Baz-Lomba et al., 2016)
	19.2	8.8-52.9	2014	(Ryu et al., 2016)
Barcelona (Spain)	18 ^a	7-31 ^a	2011- 2015	(Mastroianni et al., 2017)
Dülmen (Germany)	20.3	5.5-40	2014	(Ryu et al., 2016)
London (United Kingdom)	21.5	10.9-36	2014	(Ryu et al., 2016)
Brussels (Belgium)	21.6		2015	(Baz-Lomba et al., 2016)
Eindhoven (The Netherlands)	21.7	13.7-30.4	2014	(Ryu et al., 2016)
Amsterdam (The Netherlands)	22	14.3-30.5	2014	(Ryu et al., 2016)
Castellón (Spain)	23.4	11.6-61.6	2014	(Ryu et al., 2016)
Dortmund (Germany)	23.6	18.1-34	2014	(Ryu et al., 2016)

Munich (Germany)	29.5	0.5-47.4	2014	(Ryu et al., 2016)
Dresden (Germany)	29.4	15.1-91.7	2014	(Ryu et al., 2016)
Montreal (Canada)	29.2	21.8-38.8	2014	(Ryu et al., 2016)
Copenhagen (Denmark)	29.7		2015	(Baz-Lomba et al., 2016)
	40.2	24.6-74	2014	(Ryu et al., 2016)
Granby (Canada)	44.3	27.3-59.3	2014	(Ryu et al., 2016)
Valencia-QB ^b	40.9 ^a	27.0-56.1 ^a	2014	(Andrés-Costa et al., 2016)

^aAlcohol consumption expressed in mL/day/inhabitant (aged 15+)

^bAlcohol consumption rate during “Fallas festivity”

3.2. Spatial variation in alcohol consumption

The statistical test applied to evaluate spatial variation in alcohol consumption among different population showed that populations belonging to the same region showed no statistically significant differences in alcohol consumption (p-value > 0.05, Mann-Whitney U test) (Table 4) while, statistically significant differences between populations belonging to different regions were found (p-value < 0.05, Mann-Whitney U test) (Table 4). Particularly, alcohol consumption estimated for the population served by Bilbao WWTP was different to that observed in 9 other populations, namely, Castellón, Guadalajara, Lleida, Madrid-Centre, Santiago de Compostela, Toledo, Valencia-PI, Valencia-PHI, and Valencia-QB, with median alcohol consumption in Bilbao between 1.5 (Valencia-PHI) and 3 (Toledo) times higher than in the aforementioned cities. Also, statistically significant differences were observed between Palma de Mallorca and Toledo (consumption in Palma de Mallorca 2 times higher than in Toledo) and between Móstoles and Castellón (consumption in Móstoles 1.7 times higher than in Castellón) (Table 2 and 4).

315 **Table 4.** Comparison of alcohol consumption between pairs of investigated populations (U Mann Whitney test p-values)^a.

	Barcelona	Bilbao	Castellón	Guadalajara	Lleida	Madrid-La China	Madrid-Viveros	Móstoles	Palma de Mallorca	Reus	Santiago de Compostela	Tarragona	Toledo	Valencia-PI	Valencia-PII
Bilbao	0.114														
Castellón	0.095	0.020*													
Guadalajara	0.389	0.045*	0.209												
Lleida	0.114	0.012*	0.789	0.287											
Madrid-La China	0.148	0.012*	0.855	0.389	0.729										
Madrid-Viveros	1.000	0.075	0.237	0.601	0.114	0.171									
Móstoles	0.855	0.389	0.045*	0.095	0.075	0.075	0.534								
Palma de Mallorca	0.925	0.060	0.075	0.237	0.070	0.070	0.855	0.789							
Reus	0.729	0.114	0.662	0.662	0.237	0.729	0.789	0.287	0.662						
Santiago de Compostela	0.171	0.012*	0.925	0.348	1.000	0.855	0.148	0.075	0.060	0.601					
Tarragona	0.389	0.662	0.114	0.209	0.070	0.095	0.287	0.534	0.389	0.171	0.075				
Toledo	0.070	0.012*	0.237	0.171	0.237	0.534	0.070	0.075	0.045*	0.114	0.389	0.060			
Valencia-PI	0.459	0.045*	0.237	0.662	0.237	0.389	0.729	0.389	0.348	0.925	0.287	0.209	0.171		
Valencia-PII	0.925	0.012*	0.171	0.729	0.148	0.171	0.601	0.237	0.601	0.789	0.114	0.237	0.075	1.000	
Valencia-QB	0.171	0.012*	0.789	0.459	0.601	0.855	0.209	0.075	0.095	0.789	0.662	0.095	0.209	0.459	0.348

316
317 ^aFirstly, a non-parametric test (Kruskal Wallis test) was applied in order to compare alcohol consumption among all investigated populations since the number
318 of data per city was $n < 10$. Since $p < 0.05$, (Kruskal Wallis p-value = 0.0003887), the null hypothesis (H_0 : alcohol consumption among all investigated
319 populations is equal) was rejected and a U Mann Whitney test was applied to compare alcohol consumption between pairs of populations. False Discovery Rate
320 (FDR) correction for multiple testing was applied to reduce the number of “false positive”.

321 * $p < 0.05$, null hypothesis in U Mann Whitney test (H_0 : alcohol consumption between pairs of populations is equal) is rejected.

322

At the regional level (Figure 2b, Table 5) differences of alcohol consumption were statistically significant (p -value < 0.05 , Mann-Whitney U test) between Basque Country and all the other investigated regions, except Catalonia, and between the Balearic Islands and the region of Castilla-La Mancha and Galicia (Table 5). The median consumption of alcohol in the Basque Country (19 mL/day/inhabitant) was between 1.5 and 2.2 times higher than the median consumption observed in the Balearic Islands (12), Community of Madrid (11), Valencian Community (9.5), Castilla-La Mancha (8.7) and Galicia (8.5 mL/day/inhabitant). The Balearic Islands presented a median figure of alcohol consumption 1.5 times higher than those obtained in Castilla-La Mancha and Galicia.

As for the city size, small cities, i.e., those with official census populations $< 300,000$ inhabitants (Toledo, Guadalajara, Santiago de Compostela, Reus, Tarragona, Lleida, Castellón and Móstoles), showed significantly lower alcohol consumption rates per capita than large cities, i.e., those with official census population $> 300,000$ (p -value < 0.05 , Mann Whitney U).

Table 5. Comparison of alcohol consumption between pairs of regions (U Mann Whitney test p -values)^a.

	Castilla-La Mancha	Catalonia	Community of Madrid	Valencian Community	Galicia	Balearic Islands
Catalonia	0.088					
Community of Madrid	0.088	1.000				
Valencian Community	0.286	0.335	0.200			
Galicia	1.000	0.169	0.096	0.221		
Balearic Islands	0.029*	0.558	0.406	0.073	0.025*	
Basque Country	0.001*	0.073	0.020*	$<0.001^*$	0.004*	0.025*

^aFirstly, a Kruskal Wallis test was applied in order to compare alcohol consumption among all investigated regions since for 3 regions (Galicia, Balearic Islands and Basque Country), $n < 10$. As p -value < 0.05 (Kruskal Wallis p -value = 0.000588), the null hypothesis (H_0 : alcohol consumption among all regions is equal) was rejected and a U Mann Whitney test was applied to compare alcohol consumption between pairs of regions. False Discovery Rate (FDR) correction for multiple testing was applied to reduce the number of “false positive”.


* $p < 0.05$ and null hypothesis in U Mann Whitney (H_0 : alcohol consumption between pairs of regions is equal) is rejected.

3.3. Weekly patterns

Figure 3 shows the daily alcohol consumption expressed as mL/day/inhabitant or as the contribution of each day to the total weekly consumption observed in each population. The difference in the amount of alcohol consumed during the weekend (Saturday and Sunday) (median=15 mL/day/inhabitant) and during the weekdays (Monday to Friday) (median=9.0 mL/day/inhabitant) was found to be statistically significant ($p\text{-value} < 0.05$, Mann Whitney U).

Figure 4 shows the weekly trends of alcohol consumption in the investigated populations. The strongest differences in alcohol consumption between weekdays and weekends were observed in Reus and Toledo (with average consumption figures 2.2 and 2.0 times higher, respectively, during the weekend than during weekdays), and the weakest in Madrid-North (where Monday is the day of highest consumption) and Tarragona (where, in fact, large variations in alcohol consumption were observed throughout the week (Figure 4)).

Figure 3 also shows a general high contribution of Mondays to total weekly alcohol consumption figures when compared with the other weekdays. According to Høiseth et al., EtS can remain in urine for several hours (between 25 and 48) depending on the dose of ethanol ingested (Høiseth et al., 2008), so, the high value of alcohol consumption estimated on Monday could be attributed to the presence of EtS in wastewater from its consumption during the weekend.



365

366 **Figure 3.** Distribution of alcohol consumption throughout the week expressed as mL/day/inhabitant (Figure 2a) and the contribution of each day
 367 to the total weekly consumption (%) (Figure 2b). (*Outlier)

368

369




Figure 4. Weekly trends of alcohol consumption in the investigated populations.

3.4. Nationwide extrapolation

The total daily alcohol load (kg/day) that arrived at each WWTP was used to back-calculate alcohol consumption at the national level. Data were extrapolated taking into account that the population covered by the study was about 6.0 million inhabitants (12.8% of the Spanish population) and the total population of Spain in 2018 accounted for 46.7 million inhabitants (INE, 2018). The extrapolation resulted in annual consumption of 4.8 ± 1.1 L of pure ethanol per capita in Spain, which increases to 5.7 ± 1.2 L or 5.9 ± 1.3 L of pure ethanol when only population above 15 years (aged 15+) or adult population (aged 18+) is considered, respectively (Table 6). This value is in line with official data reported by the National Health Survey (INE) (Table 7) that reports an average weekly consumption of 13 mL/day/inhabitant (aged 15+) equivalent to an average annual consumption of 4.7 L of pure ethanol per capita (aged 15+), and also with official data published by the Spanish Ministry of Agriculture, Fishing and Food, which indicates consumption of 51.8 L of beer per capita (+18) (MAPA, 2018), equivalent to 4.3 L of pure ethanol per capita (aged 18+) taking into account that alcohol consumption by type of alcoholic beverage is distributed as 54% beer, 18% wine and 28% spirits and the alcohol content in each one is 4.5, 12 and 40%, respectively (WHO, 2018). On the contrary, a higher alcohol consumption rate (10 L of pure ethanol per capita (aged 15+)) was reported for Spain in the WHO report (WHO, 2018).

Table 6. Average alcohol consumption estimated in Spain through WBE.

	Alcohol consumption in the investigated populations	Alcohol consumption in Spain				
	Kg/day	Kg/day	L/day	L/year/ inhabitants	L/year/ inhabitants (aged 15+)	L/year/ inhabitants (aged 18+)
Tuesday	48187	376424	477090	3.7	4.4	4.6
Wednesday	50115	391487	496181	3.9	4.6	4.8
Thursday	55403	432792	548532	4.3	5.1	5.3
Friday	57734	451005	571616	4.5	5.3	5.5
Saturday	84030	656420	831965	6.5	7.7	8.0
Sunday	77172	602852	764071	6.0	7.1	7.3
Monday	62306	486721	616884	4.8	5.7	5.9
<i>Average</i>	<i>62135</i>	<i>485386</i>	<i>615191</i>	<i>4.8</i>	<i>5.7</i>	<i>5.9</i>
<i>SD</i>	<i>13597</i>	<i>106216</i>	<i>134621</i>	<i>1.1</i>	<i>1.2</i>	<i>1.3</i>

Table 7. Average alcohol consumption (mL/day/inhabitant (aged 15+)) in the investigated regions in this study and Spain reported by the National Health Survey (INE).

	Week (Mon-Sun)		Weekdays (Mon-Thurs)		Weekend (Frid-Sun)	
	Average	sd	Average	sd	Average	sd
Balearic Island	18	14	15	14	22	17
Basque Country	19	14	11	15	30	19
Castilla-La Mancha	13	13	7.5	13	20	17
Catalonia	16	13	10	13	23	17
Community of Madrid	14	16	8.0	16	21	18
Galicia	20	12	16	13	25	13
Valencian Community	14	11	8.5	12	22	15
<i>Spain</i>	<i>13</i>	<i>12</i>	<i>8.4</i>	<i>12</i>	<i>19</i>	<i>16</i>

Source: National Health Survey (INE, 2017).

<https://www.ine.es/jaxi/Tabla.htm?path=/t15/p419/a2017/p03/10/&file=03011.px&L=0>

4. Discussion

In this study, alcohol consumption in different populations of Spain was estimated through WBE. The population investigated covers 12.8% of the total Spanish population and is distributed around 13 main cities and 7 different regions. Results showed spatial variations in alcohol consumption among specific populations and regions. Although Tarragona, Bilbao, and Móstoles were the cities with the highest average alcohol consumption figures, Bilbao was the only one where alcohol consumption was significantly different from several other populations (see Table 4 and Figure 2). Also, alcohol consumption in Palma de Mallorca and Móstoles was significantly higher than in Toledo and Castellón, respectively. WBE-derived alcohol consumption figures were compared with the latest data reported by the National Health Survey carried out by the Spanish Ministry of Health, Consumption and Social Welfare in collaboration with the National Institute of Statistics (INE) (INE, 2017) and with prevalence data reported in the Annual Report of the Spanish Observatory on Drugs and Drugs Addiction (OEDA, 2019). Since official data are only provided at the level of regions, the average alcohol consumption obtained in each investigated population was compared with consumption data reported for its corresponding region. Figure 5 compares WBE data and INE National Health Survey data. WBE-derived alcohol consumption figures in five of the investigated populations (Toledo, Lleida, Madrid-Centre, Castellón, and Valencia-QB) showed good correlation with INE official data at the region level, being the differences of consumption figures lower than 13%, whereas a weaker correlation (differences of consumption between 22 and 30%) was observed in 4 populations (Palma de Mallorca, Reus, Valencia-PI, and Valencia-PII). WBE-derived data in the remaining populations (Bilbao, Guadalajara, Barcelona, Tarragona, Madrid-North, Móstoles, and Santiago de Compostela) showed larger differences with official INE data.




Figure 5. Alcohol consumption estimated in the investigated populations through WBE (red square), data reported for the corresponding region in the INE National Health Survey (blue line), and differences of consumption between WBE data and survey data (grated bars) (%). (The bars within the dark green zone delimit consumption differences between both methodologies below 15% and those within the light green zone below 30%)

On the other hand, the comparison of WBE-data with prevalence data of alcohol consumption reported for each region, showed poor correlation when all investigated populations were considered (see Figure 6). However, as shown in Figure 7, when the data from the 7 populations that did not correlate with official INE consumption figures (Bilbao, Guadalajara, Barcelona,

Tarragona, Madrid-North, Móstoles, and Santiago de Compostela) were removed, a significant correlation was observed (r^2 “Lifetime prevalence”: 0.4499, p -value < 0.05 ; r^2 “Last year prevalence”: 0.5407, p -value < 0.05). According to WBE-data the population belonging to the Basque Country presented a significantly higher consumption than populations belonging to the other regions (except Catalonia), and alcohol consumption in the Balearic Islands was significantly higher than in Castilla-La Mancha and Galicia (Figure 2b, Table 5). Compared to prevalence data reported by the Annual Report (Figure 8), WBE results are in agreement with prevalence data only in the case of the Balearic Islands since the Balearic Islands show a higher prevalence of consumption than Castilla-La Mancha and Galicia. On the contrary, in the case of the Basque Country, the prevalence of alcohol consumption, although above the Spanish average, is similar to that reported for the Valencian Community or Galicia (Figure 8).

The differences observed between WBE-derived alcohol consumption figures and established indicators could have different explanations. On the one hand, data reported by established methods may not represent the actual consumption by the population since they are affected by a degree of uncertainty. The two established indicators used to compare the WBE-derived estimates, provided indeed different results, in the sense that the highest prevalence data was reported for the Balearic Islands (see Figure 8) whereas the highest alcohol consumption rate was reported for Galicia in the INE National Health Survey (Table 7). On the other hand, the populations sampled may not be representative of alcohol consumption in the whole region. As previously demonstrated, significant differences in alcohol consumption were observed between small and large populations (section 3.2). In some regions, only one municipality was sampled (i.e., the Balearic Islands and Galicia) which may not adjust to the alcohol consumption patterns of the whole region. This hypothesis is supported by the fact that within the same region, WBE-data derived from some populations correlated well with the INE survey data, whereas others did not (see Castilla-La Mancha, Catalonia, and Community of Madrid in Figure

5). Despite this, at the national level, the annual alcohol consumption rate obtained through WBE was comparable to that reported by the National Health Survey, which may indicate that the sampled population is quite representative of the whole country.




Figure 6. Correlation between average alcohol consumption estimated in each city by WBE (mL/day/inhabitant) and prevalence data (“Lifetime prevalence”, “Last year prevalence” and “Last month prevalence”) reported by its region in the annual Report of the Spanish Observatory on Drugs and Drugs Addiction 2019. (Data from all investigated populations are shown; Spearman correlation p-values < 0.05 were considered statistically significant).





Figure 7. Correlation between average alcohol consumption estimated in each city by WBE (mL/day/inhabitant) and prevalence data (“Lifetime prevalence”, “Last year prevalence” and “Last month prevalence”) reported by its region in the Annual Report of the Spanish Observatory on Drugs and Drugs Addiction 2019. (Data from Guadalajara, Barcelona, Tarragona, Madrid-North, Móstoles, Santiago de Compostela, and Bilbao were excluded; Spearman correlation p-value < 0.05 were considered statistically significant).

480

481 Unlike the Spanish National Health Survey, the national WBE-derived data show a low
 482 correlation to those reported by the WHO. This fact was also observed in the nation-wide study
 483 carried out in Belgium (Boogaerts et al., 2016) in which the national alcohol consumption rate
 484 estimated by the WBE approach was half that reported by the WHO. Such differences could be
 485 attributed to the fact that WHO data may not appropriately represent the actual consumption of
 486 alcohol by the population. WHO data are derived from production, import, export and sale data,
 487 which in countries where there is not a strict control, like Spain, can lead to an overestimation
 488 of consumption, since alcohol can be stored and not consumed shortly after purchase. In
 489 countries like Norway, where sales statistics are among the most accurate in the world, a good
 490 correlation was obtained between WBE and WHO data (Reid et al., 2011).

491



492

493 **Figure 8.** Prevalence data of alcohol consumption in the investigated regions and Spain
 494 reported in the Annual Report of the Spanish Observatory on Drugs and Drugs Addiction 2019.

495

496 As expected, the weekly consumption patterns in most populations showed an increase in
497 alcohol consumption during the weekend. Saturday and Sunday were the days when alcohol
498 consumption contributed the most to the total weekly consumption, with a median contribution
499 of 20%, while the remaining days of the week contributed between 11% (Tuesday) and 14%
500 (Monday) (Figure 2b). Similar results were obtained in Australia, where each weekend day
501 contributed with 20% to the weekly consumption rate, while the rest of the days of the week
502 varied between 11% and 13% (Lai et al., 2018). The increase in alcohol consumption during
503 the weekend was also reported in an international study conducted in 11 different countries
504 worldwide (Baz-Lomba et al., 2016), in Norway (Reid et al., 2011), Belgium (Boogaerts et al.,
505 2016; van Wel et al., 2016), and in Spain, where previous studies, far less ambitious than the
506 present study, were done in Barcelona (Mastroianni et al., 2017, 2014), Santiago de Compostela
507 (Rodríguez-Álvarez et al., 2015, 2014) and Valencia (Andrés-Costa et al., 2016). The increase
508 of alcohol consumption during the weekend was also reported by the INE National Health
509 Survey for all regions investigated in the present study in terms of consumption rate (see Table
510 7) (INE, 2017), so again, a good correlation was obtained between WBE approach and
511 established indicators.

512 Despite the good correlation mostly obtained between WBE-derived data and those obtained
513 with established indicators, the estimates of alcohol consumption through WBE are affected by
514 some degree of uncertainty that should be taken into consideration. On the one hand, it has been
515 shown that EtS is stable in wastewater (one week at room temperature and more than 1 month
516 at -20°C) (Rodríguez-Álvarez et al., 2014); however, EtS could degrade to some extent in
517 sewage systems (Banks et al., 2018; Gao et al., 2018). This could lead to an underestimation of
518 the real alcohol consumption, which could (partially) explain the lower consumption estimates
519 obtained through WBE compared to those reported by the WHO. However, degradation can

be corrected by applying a correction factor, as demonstrated in a recent study conducted in Australia (Zheng et al., 2020). On the other hand, the excretion rate used to back-calculate alcohol consumption was obtained from two studies in which only 10 men (Høiseth et al., 2008) and one man (Wurst et al., 2006) were investigated, respectively. Further studies involving more volunteers of different ages, gender, or race, or studying the excretion rate among the Spanish population could help to obtain a more representative excretion rate which would increase the accuracy of back-calculations. An additional source of uncertainty may come from the sampling (collection of a not representative sample). In this study, WBE data have been obtained from samples collected during only one week, which may not be representative of alcohol consumption throughout the entire year. Increasing the sampling period, several times a year or during consecutive years could be used to obtain temporal trends in alcohol consumption within one year and throughout the years. Furthermore, unlike the estimates at the national level, the differences observed in some regions between WBE-derived data and those reported by established indicators could indicate that population sampled are not representative of the whole region. Increasing the population sampled or sampling populations of different sizes within one region could lead to a more representative picture of the habits of consumption of the whole region. Finally, other sources of uncertainty may come from inaccurate measurement of the water volume entering the plant, and the calculation of the size of the population that contributes to the total EtS load measured in wastewater (Castiglioni et al., 2013). In the present study, the latter was assessed using different methods (census data, population connected to the WWTP, water quality parameters), following in each case the recommendations provided by the experts of the WWTP in order to obtain the value that best reflects the population served by each WWTP.

Regardless of the aforementioned limitations, the WBE approach appears as a promising, convenient tool for alcohol consumption assessment, which surely needs to be refined in the

next few years. WBE is much useful to establish spatial and temporal variations in alcohol consumption in a fast, objective, and inexpensive way, providing data in nearly real-time. WBE can complement in this way the information gained with the established methodologies which are also affected by some uncertainties. In this sense, the use of different indicators and sources of information would improve the alcohol consumption estimates and hence, contribute to better development and evaluation of health care management plans and policies.

5. Conclusions

The present work represents the first nation-wide study conducted in Spain to evaluate alcohol consumption through the application of the WBE approach and is one of the first nation-wide assessments available worldwide. The study has covered 13 main cities (in some cases including surrounding towns) that represent 12.8% of the Spanish total population. The results show that WBE is a useful tool to define spatial and temporal variations in alcohol consumption in a fast, objective, and inexpensive way, providing complementary data to the information gained with the established methodologies. The WBE-derived alcohol consumption data correlated well (within $\pm 15\%$) with official data reported by conventional methods at the regional level in 5 out of the 16 populations investigated (31% of the total population examined), and satisfactorily (within $\pm 30\%$) in 9 of the populations studied (accounting for 56% of the scrutinized population). Also, extrapolation of WBE-derived alcohol consumption estimates to the national territory led to an annual consumption of alcohol in Spain comparable to that reported for Spain by the National Health Survey, although, lower than that reported by the WHO. The comparison of WBE data with those obtained with established consumption indicators should be done with caution because both methodologies are subject to some uncertainties. Increasing the sampling period, the sampled population, and conducting further

studies on alcohol metabolism to establish appropriate correction factors would help to reduce the main uncertainties associated with WBE and, therefore, to improve the accuracy of the consumption estimates.

References

Andrés-Costa, M.J., Escrivá, Ú., Andreu, V., Picó, Y., 2016. Estimation of alcohol consumption during “Fallas” festivity in the wastewater of Valencia city (Spain) using ethyl sulfate as a biomarker. *Sci. Total Environ.* 541, 616–622. <https://doi.org/10.1016/j.scitotenv.2015.09.126>

Banks, A.P.W., Lai, F.Y., Mueller, J.F., Jiang, G., Carter, S., Thai, P.K., 2018. Potential impact of the sewer system on the applicability of alcohol and tobacco biomarkers in wastewater-based epidemiology. *Drug Test. Anal.* 10, 530–538. <https://doi.org/10.1002/dta.2246>

Baz-Lomba, J.A., Salvatore, S., Gracia-Lor, E., Bade, R., Castiglioni, S., Castrignanò, E., Causanilles, A., Hernandez, F., Kasprzyk-Hordern, B., Kinyua, J., McCall, A.-K., van Nuijs, A., Ort, C., Plósz, B.G., Ramin, P., Reid, M., Rousis, N.I., Ryu, Y., de Voogt, P., Bramness, J., Thomas, K., 2016. Comparison of pharmaceutical, illicit drug, alcohol, nicotine and caffeine levels in wastewater with sale, seizure and consumption data for 8 European cities. *BMC Public Health* 16, 1035. <https://doi.org/10.1186/s12889-016-3686-5>

Boogaerts, T., Covaci, A., Kinyua, J., Neels, H., Van Nuijs, A.L.N., 2016. Spatial and temporal trends in alcohol consumption in Belgian cities: A wastewater-based approach. *Drug Alcohol Depend.* 160, 170–176. <https://doi.org/10.1016/j.drugalcdep.2016.01.002>

Castiglioni, S., Bijlsma, L., Covaci, A., Emke, E., Hernández, F., Reid, M., Ort, C., Thomas,

K. V., Van Nuijs, A.L.N., De Voogt, P., Zuccato, E., 2013. Evaluation of uncertainties associated with the determination of community drug use through the measurement of sewage drug biomarkers. *Environ. Sci. Technol.* 47, 1452–1460. <https://doi.org/10.1021/es302722f>

Chen, J., Venkatesan, A.K., Halden, R.U., 2019. Alcohol and nicotine consumption trends in three U.S. communities determined by wastewater-based epidemiology. *Sci. Total Environ.* 656, 174–183. <https://doi.org/10.1016/j.scitotenv.2018.11.350>

European Monitoring Centre for Drugs and Drug Addiction, 2016 (EMCDDA, 2016). http://www.emcdda.europa.eu/system/files/publications/2273/TDXD16022ENC_4.pdf (accessed 3.9.20).

Gao, J., Li, J., Jiang, G., Yuan, Z., Eaglesham, G., Covaci, A., Mueller, J.F., Thai, P.K., 2018. Stability of alcohol and tobacco consumption biomarkers in a real rising main sewer. *Water Res.* 138, 19–26. <https://doi.org/10.1016/j.watres.2018.03.036>

Gao, J., Zheng, Q., Lai, F.Y., Gartner, C., Du, P., Ren, Y., Li, X., Wang, D., Mueller, J.F., Thai, P.K., 2020. Using wastewater-based epidemiology to estimate consumption of alcohol and nicotine in major cities of China in 2014 and 2016. *Environ. Int.* 136, 105492. <https://doi.org/10.1016/j.envint.2020.105492>

Gatidou, G., Kinyua, J., van Nuijs, A.L.N., Gracia-Lor, E., Castiglioni, S., Covaci, A., Stasinakis, A.S., 2016. Drugs of abuse and alcohol consumption among different groups of population on the Greek Island of Lesbos through sewage-based epidemiology. *Sci. Total Environ.* 563–564, 633–640. <https://doi.org/10.1016/j.scitotenv.2016.04.130>

González-Mariño, I., Baz-Lomba, J.A., Alygizakis, N.A., Andrés-Costa, M.J., Bade, R., Barron, L.P., Been, F., Berset, J.-D., Bijlsma, L., Bodík, I., Brenner, A., Brock, A.L.,

615 Burgard, D.A., Castrignanò, E., Christophoridis, C.E., Covaci, A., de Voogt, P., Devault,
 616 D.A., Dias, M.J., Emke, E., Fatta-Kassinos, D., Fedorova, G., Fytianos, K., Gerber, C.,
 617 Grabic, R., Grüner, S., Gunnar, T., Hapeshi, E., Heath, E., Helm, B., Hernández, F.,
 618 Kankaanpää, A., Karolak, S., Kasprzyk-Hordern, B., Krizman-Matasic, I., Lai, F.Y.,
 619 Lechowicz, W., Lopes, A., López de Alda, M., López-García, E., Löve, A.S.C.,
 620 Mastroianni, N., McEneff, G.L., Montes, R., Munro, K., Nefau, T., Oberacher, H.,
 621 O'Brien, J.W., Olafsdottir, K., Picó, Y., Plósz, B.G., Polesel, F., Postigo, C., Quintana,
 622 J.B., Ramin, P., Reid, M.J., Rice, J., Rodil, R., Senta, I., Simões, S.M., Sremacki, M.M.,
 623 Styszko, K., Terzic, S., Thomaidis, N.S., Thomas, K.V., Tschärke, B.J., van Nuijs, A.L.N.,
 624 Yargeau, V., Zuccato, E., Castiglioni, S., Ort, C., 2019. Spatio-temporal assessment of
 625 illicit drug use at large scale: evidence from 7 years of international wastewater
 626 monitoring. *Addiction* 115, 109-120. <https://doi.org/10.1111/add.14767>

627 Helander, A., Beck, O., 2005. Ethyl Sulfate: A Metabolite of Ethanol in Humans and a Potential
 628 Biomarker of Acute Alcohol Intake. *J. Anal. Toxicol.* 29, 270–274.
 629 <https://doi.org/10.1093/jat/29.5.270>

630 Høiseth, G., Bernard, J.P., Stephanson, N., Normann, P.T., Christophersen, A.S., Mørland, J.,
 631 Helander, A., 2008. Comparison between the urinary alcohol markers EtG, EtS, and
 632 GTOL/5-HIAA in a controlled drinking experiment. *Alcohol Alcohol.* 43, 187–191.
 633 <https://doi.org/10.1093/alcalc/agm175>

634 Instituto Nacional de Estadística, 2017 (INE, 2017). Consumo medio diario de alcohol (en
 635 gramos de alcohol puro) según sexo y comunidad autónoma. Media y desviación típica.
 636 Población de 15 y más años que consume bebidas alcohólicas uno o más días a la semana.
 637 <https://www.ine.es/jaxi/Tabla.htm?path=/t15/p419/a2017/p03/10/&file=03011.px&L=0>
 638 (accessed 1.23.20).

639 Instituto Nacional de Estadística, 2018 (INE, 2018). Población residente por fecha, sexo y
 640 edad(9663). <https://www.ine.es/jaxiT3/Datos.htm?t=9663> (accessed 3.3.20).

641 Lai, F.Y., Gartner, C., Hall, W., Carter, S., O'Brien, J., Tschärke, B.J., Been, F., Gerber, C.,
 642 White, J., Thai, P., Bruno, R., Prichard, J., Kirkbride, K.P., Mueller, J.F., 2018. Measuring
 643 spatial and temporal trends of nicotine and alcohol consumption in Australia using
 644 wastewater-based epidemiology. *Addiction* 113, 1127–1136.
 645 <https://doi.org/10.1111/add.14157>

646 Ministerio de Agricultura, Pesca y Alimentación, 2018 (MAPA, 2018)
 647 https://cerveceros.org/uploads/5d1b32241b8be__Informe
 648 *SocioeconomicoCerveza_2018.pdf* (accessed 11.28.19).

649 Mastroianni, N., López de Alda, M., Barcelo, D., 2014. Analysis of ethyl sulfate in raw
 650 wastewater for estimation of alcohol consumption and its correlation with drugs of abuse
 651 in the city of Barcelona. *J. Chromatogr. A* 1360, 93–99.
 652 <https://doi.org/10.1016/j.chroma.2014.07.051>

653 Mastroianni, N., López-García, E., Postigo, C., Barceló, D., López de Alda, M., 2017. Five-
 654 year monitoring of 19 illicit and legal substances of abuse at the inlet of a wastewater
 655 treatment plant in Barcelona (NE Spain) and estimation of drug consumption patterns and
 656 trends. *Sci. Total Environ.* 609, 916–926.
 657 <https://doi.org/10.1016/J.SCITOTENV.2017.07.126>

658 Nguyen, H.T., Thai, P.K., Kaserzon, S.L., O'Brien, J.W., Eaglesham, G., Mueller, J.F., 2018.
 659 Assessment of drugs and personal care products biomarkers in the influent and effluent of
 660 two wastewater treatment plants in Ho Chi Minh City, Vietnam. *Sci. Total Environ.* 631–
 661 632, 469–475. <https://doi.org/10.1016/j.scitotenv.2018.02.309>

662 Observatorio Español de las Drogas y las Adicciones, 2019 (OEDA, 2019). INFORME 2019
 663 Alcohol, tabaco y drogas ilegales en España.

664 Probst, C., Fleischmann, A., Gmel, G., Poznyak, V., Rekve, D., Riley, L., Rylett, M., Shield,
 665 K.D., Rehm, J., 2019. The global proportion and volume of unrecorded alcohol in 2015.
 666 J. Glob. Health 9. <https://doi.org/10.7189/jogh.09.010421>

667 Reid, M.J., Langford, K.H., Mørland, J., Thomas, K. V., 2011. Analysis and Interpretation of
 668 Specific Ethanol Metabolites, Ethyl Sulfate, and Ethyl Glucuronide in Sewage Effluent
 669 for the Quantitative Measurement of Regional Alcohol Consumption. Alcohol. Clin. Exp.
 670 Res. no-no. <https://doi.org/10.1111/j.1530-0277.2011.01505.x>

671 Rodríguez-Álvarez, T., Rodil, R., Cela, R., Quintana, J.B., 2014. Ion-pair reversed-phase liquid
 672 chromatography-quadrupole-time-of-flight and triple-quadrupole-mass spectrometry
 673 determination of ethyl sulfate in wastewater for alcohol consumption tracing. J.
 674 Chromatogr. A 1328, 35–42. <https://doi.org/10.1016/j.chroma.2013.12.076>

675 Rodríguez-Álvarez, T., Racamonde, I., González-Mariño, I., Borsotti, A., Rodil, R., Rodríguez,
 676 I., Zuccato, E., Quintana, J.B., Castiglioni, S., 2015. Alcohol and cocaine co-consumption
 677 in two European cities assessed by wastewater analysis. Sci. Total Environ. 536, 91–98.
 678 <https://doi.org/10.1016/j.scitotenv.2015.07.016>

679 Ryu, Y., Barceló, D., Barron, L.P., Bijlsma, L., Castiglioni, S., de Voogt, P., Emke, E.,
 680 Hernández, F., Lai, F.Y., Lopes, A., López de Alda, M., Mastroianni, N., Munro, K.,
 681 O'Brien, J., Ort, C., Plósz, B.G., Reid, M.J., Yargeau, V., Thomas, K. V., 2016.
 682 Comparative measurement and quantitative risk assessment of alcohol consumption
 683 through wastewater-based epidemiology: An international study in 20 cities. Sci. Total
 684 Environ. 565, 977–983. <https://doi.org/10.1016/j.scitotenv.2016.04.138>

- Stockwell, T., Zhao, J., Greenfield, T., Li, J., Livingston, M., Meng, Y., 2016. Estimating under- and over-reporting of drinking in national surveys of alcohol consumption: identification of consistent biases across four English-speaking countries. *Addiction* 111, 1203–1213. <https://doi.org/10.1111/add.13373>
- van Wel, J.H.P., Gracia-Lor, E., van Nuijs, A.L.N., Kinyua, J., Salvatore, S., Castiglioni, S., Bramness, J.G., Covaci, A., Van Hal, G., 2016. Investigation of agreement between wastewater-based epidemiology and survey data on alcohol and nicotine use in a community. *Drug Alcohol Depend.* 162, 170–175. <https://doi.org/10.1016/j.drugalcdep.2016.03.002>
- World Health Organization, 2018 (WHO, 2018)<https://apps.who.int/iris/bitstream/handle/10665/274603/9789241565639-eng.pdf?ua=1> (accessed 11.15.19).
- Wurst, F.M., Dresen, S., Allen, J.P., Wiesbeck, G., Graf, M., Weinmann, W., 2006. Ethyl sulphate: A direct ethanol metabolite reflecting recent alcohol consumption. *Addiction* 101, 204–211. <https://doi.org/10.1111/j.1360-0443.2005.01245.x>
- Zheng, Q., Tschärke, B.J., Krapp, C., O’Brien, J.W., Mackie, R.S., Connor, J., Mueller, J.F., Thomas, K. V., Thai, P.K., 2020. New approach for the measurement of long-term alcohol consumption trends: Application of wastewater-based epidemiology in an Australian regional city. *Drug Alcohol Depend.* 207. <https://doi.org/10.1016/j.drugalcdep.2019.107795>

708

709

710

Supporting Information

Assessing alcohol consumption through wastewater-based epidemiology: Spain as a case study

Ester López-García¹, Carlos Pérez López¹, Cristina Postigo^{1*}, Vicente Andreu², Lubertus Bijlsma³, Iria González-Mariño^{4,5}, Félix Hernández³, Rosa Maria Marcé⁶, Rosa Montes⁴, Yolanda Picó², Eva Pocurull⁶, Andreu Rico⁷, Rosario Rodil⁴, María Rosende⁸, Yolanda Valcárcel⁹, Olatz Zuloaga¹⁰, José Benito Quintana⁴, Miren López de Alda^{1*}

¹Water, Environmental, and Food Chemistry Unit (ENFOCHEM), Department of Environmental Chemistry, Institute of Environmental Assessment and Water Research (IDAEA-CSIC), Barcelona

²Food and Environmental Safety Research Group (SAMA-UV), Desertification Research Centre (CIDE), CSIC-Generalitat Valenciana-University of Valencia, Valencia

³Environmental and Public Health Analytical Chemistry, Research Institute for Pesticides and Water, University Jaume I, Castellón

⁴Department of Analytical Chemistry, Nutrition and Food Sciences, Institute of Research on Chemical and Biological Analysis (IAQBUS), Universidade de Santiago de Compostela, Santiago de Compostela

⁵Department of Analytical Chemistry, Nutrition and Bromatology, Faculty of Chemical Sciences, University of Salamanca, Salamanca.

⁶Department of Analytical Chemistry and Organic Chemistry, Universitat Rovira i Virgili, Tarragona

⁷IMDEA Water Institute, Science and Technology Campus of the University of Alcalá, Alcalá de Henares

⁸FI-TRACE group, Department of Chemistry, University of the Balearic Islands, Palma de Mallorca

⁹Research Group in Environmental Toxicology and Risk Assessment (TAyER), Medical specialties and Public Health, Faculty of Health Sciences, Rey Juan Carlos University, Madrid

¹⁰Department of Analytical Chemistry, Faculty of Science and Technology (UPV/EHU) & Plentzia Marine Station of Basque Country University (EHU/UPV), Basque Country

***Corresponding Authors:** Cristina Postigo; Miren López de Alda
Institute of Environmental Assessment and Water Research
(IDAEA-CSIC)
Department of Environmental Chemistry
C/ Jordi Girona 18-26, 08034 Barcelona, Spain.
cprqam@cid.csic.es; miren.lopezdealda@idaea.csic.es
Tel: +34-934-006-100, Fax: +34-932-045-904

Number of pages: 16

Number of tables: 6

Number of figures: 4

List of Tables

Table S1. Description of sampled WWTPs (name, population served and locations/districts covered with main city in bold) and the sampling protocol carried out (location of autosampler, and sampling mode, start time and period).....	3
Table S2. Alcohol consumption rates estimated by means of WBE approach in different cities worldwide.....	6
Table S3. Comparison of alcohol consumption between pairs of investigated populations (U Mann Whitney test p-values) ^a	8
Table S4. Comparison of alcohol consumption between pairs of regions (U Mann Whitney test p-values) ^a	9
Table S5. Average alcohol consumption estimated in Spain through WBE.....	10
Table S6. Average alcohol consumption (mL/day/inhabitant (aged 15+)) in the investigated regions in this study and Spain reported by the National Health Survey (INE).	11

List of Figures

Figure S1. Map of Spain with the location of the sampled WWTPs (regions are indicated in different colors).	12
Figure S2. Weekly trends of alcohol consumption in the investigated populations.	13
Figure S3. Correlation between average alcohol consumption estimated in each city by WBE (mL/day/inhabitant) and prevalence data (“Lifetime prevalence”, “Last year prevalence” and “Last month prevalence”) reported by its region in the annual Report of the Spanish Observatory on Drugs and Drugs Addiction 2019. (Data from all investigated populations are shown; Spearman correlation p-values < 0.05 were considered statistically significant).....	14
Figure S4. Prevalence data of alcohol consumption in the investigated regions and Spain reported in the Annual Report of the Spanish Observatory on Drugs and Drugs Addiction 2019.....	15

Table S1. Description of sampled WWTPs (name, population served and locations/districts covered with main city in bold) and the sampling protocol carried out (location of autosampler, and sampling mode, start time and period).

Regions	City ^a	WWTP name	Population served by the WWTPs	Method used to estimate the population served ^b	Locations/districts served by the WWTPs	Percentage of the main city covered by the WWTP ^c	Location of autosampler	Sampling mode ^d	Sampling start time	Sampling period
Balearic Islands	Palma de Mallorca	Palma I	406,492	Census 2017	Palma beach, Sant Jordi, El Pili-lari, Son Sant Joan airport, part of Palma de Mallorca	100	After fine screen	T (100 mL/ 15 min)	10:00	10/04/2018-16/04/2018
		Palma II	47,961	Census 2017	Palma de Mallorca (main part), Marratxí, Esporles, Bunyola and Son Castelló, Can Valero, Son Rosinyol industrial states		After fine screen	T (100 mL/ 15 min)	10:00	18/04/2018-24/04/2018
Basque Country	Bilbao	Galindo	860,237	Census 2016	Abanto-Zierbena, Alontotegi, Arrigorriaga, Barakaldo, Barrika, Basauri, Berango, Bilbao, Derio, Erandio, Etxebarri, Galdakao, Getxo, Leioa, Lezama, Loiu, Ortuella, Portugalete, Santurtzi, Sestao, Sondika, Sopelana, Trapagaran, Ugao-Miravalles, Urduliz, Zamudio, Zaratamo, Zeberio	100	After coarse screens and pumping	T (100mL/ 60 min)	8:00	17/04/2018-23/04/2018

Castilla-La Mancha	Toledo	Estiviel	79,793	Average BOD April-May 2018	Toledo	100	After sieving	T (100 mL/ 15 min)	8:00	17/04/2018-23/04/2018
	Guadalajara	Guadalajara	94,755	Average BOD Jan-April 2018	Guadalajara	100	Before fine screen	T (200 mL/ 60 min)	10:00	02/05/2018-08/05/2018
Catalonia	Barcelona	Baix Llobregat	1,163,154	Census 2017	Barcelona, Cervelló, Cornellà de Llobregat, Esplugues de Llobregat, Hospitalet de Llobregat, El Prat de Llobregat, Sant Boi de Llobregat, San Joan Despí, San Just Desvern	35	Mechanical bar screens	T (50 mL/ 10min)	9:00	14/03/2018-20/03/2018
	Lleida	Lleida	143,612	Census 2017	Lleida , Alpicat	100	Before fine screen	T (200 mL/ 60 min)	6:00	07/03/2018-13/03/2018
	Reus	Reus	115,000	Census 2017	Reus , Castellvell, Almoester	100	After fine screen	F	20:00	17/04/2018-23/04/2018
	Tarragona	Tarragona	142,635	Census 2017	Tarragona , La Canonja, Els Pallaresos	100	Before fine screen	T (450 mL/ 60 min)	8:00-9:00	17/04/2018-23/04/2018
Community of Madrid	Madrid	Madrid-Centre	727,176	Average COD for the sampling period	Madrid-Center (Neighborhoods: Chamartín, Tetuán, Moncloa-Aravaca, Chamberí, Centro, Arganzuela, Retiro, Ciudad Lineal, Salamanca, Moratalaz, Puente de Vallecas).	30	After sieving	T (400 mL/ 30 min)	8:00	16/05/2018-22/05/2018

	Madrid	Madrid-North	227,869	Average BOD 2016 (with 60 g BOD/d)	Pozuelo y Madrid-North: (Neighborhoods: Chamartín, Tetuán, Moncloa, Aravaca, Fuencarral, El Pardo, Las Rozas, Majadahonda)		After fine screen	T (100 mL/ 60 min)	8:00	20/06/2018-26/06/2018
	Móstoles	El Soto	187,281	H x 3.5 (WWTP recomm.)	Móstoles, Alcorcón, Fuenlabrada	90	After fine screen	T (100 mL/ 60 min)	8:00	17/05/2018-23/05/2018
Galicia	Santiago de Compostela	Silvouta	136,500	H x 2.5 (WWTP recomm.)	Santiago de Compostela	100	After fine screen	T (150 mL/ 10 min)	9:00	13/03/2018-19/03/2018
Community of Valencia	Castellón	Castellón de la Plana	171,669	Census 2015	Castellón	100	Before fine screen	T (100 mL/ 15 min)	8:30	11/04/2018-17/04/2018
	Valencia	Pinedo I (Valencia-PI)	527,222	COD	Valencia (main part)	100	After fine screen	T (100 mL/ 60 min)	8:00	10/04/2018-16/04/2018
	Valencia	Pinedo II (Valencia-PII)	788,242	COD	Albal, Alcàsser, Alfafar, Benetúser, Beniparrell, Burjassot, Catarroja, Llocnou de la Corona, Massanassa, Mislata, Paiporta, Paterna, Picanya, Picassent, Sedaví, Silla, Torrent, part of Valencia		After fine screen	T (100 mL/ 60 min)	8:00	10/04/2018-16/04/2018
	Valencia	Quart-Benager (Valencia-QB)	162,249	COD	Alaquàs, Aldaia, Manises, Mislata, Quart de Poblet, Xirivella		After fine screen	F	8:00	10/04/2018-16/04/2018

^aName of the main city served by the WWTPs (some WWTPs receive wastewater from other towns included in the capital metropolitan area). ^bBOD: Biochemical Oxygen Demand; COD: Chemical Oxygen Demand; H: Number of homes connected to the sewage system. WWTP recomm: following WWTP recommendations. ^cWWTPs serving parts of the same main city were considered all together for this calculation. ^dT: time-proportional (volume sampled/frequency of sampling); F: Flow-proportional

Table S2. Alcohol consumption rates estimated by means of WBE approach in different cities worldwide.

City (Country)	Alcohol consumption (mL/day/inhabitant)		Year	Reference
	Average	Range		
Ho Chi Minh (Vietnam)	3.1-3.9		2015	(Nguyen et al., 2018)
Lesvos (Greece)	3.4/5.4	1.7-7.2/2.2-11.2	2015	(Gatidou et al., 2016)
Valencia-PII (Spain)	3.3 ^a	1.1-6.4 ^a	2014	(Andrés-Costa et al., 2016)
Milan (Italy)	5.1	3.2-10.5	2012-2014	(Rodríguez-Álvarez et al., 2015)
	6.4	5.1-8.1	2014	(Ryu et al., 2016)
	6.6		2015	(Baz-Lomba et al., 2016)
Valencia-QB (Spain)	5.9 ^a	3.3-12.8 ^a	2014	(Andrés-Costa et al., 2016)
Valencia-PII ^b (Spain)	6.1 ^a	4.3-9.1 ^a	2014	(Andrés-Costa et al., 2016)
Valencia-PI (Spain)	6.2 ^a	1.1-18.31 ^a	2014	(Andrés-Costa et al., 2016)
Lugano (Switzerland)	6.5	4.5-8.4	2014	(Ryu et al., 2016)
Toowoomba (Australia)	9.7	6.9-14.5	2014	(Ryu et al., 2016)
Utrecht (The Netherlands)	10.8		2015	(Baz-Lomba et al., 2016)
	12.9	7.7-20.7	2014	(Ryu et al., 2016)
Santiago de Compostela (Spain)	13.6	3.8-22.6	2012-2014	(Rodríguez-Álvarez et al., 2015)
	16.3	9.3-23.5	2012	(Rodríguez-Álvarez et al., 2014)
Valencia-PII ^b	14.4 ^a	4.9-23.8 ^a	2014	(Andrés-Costa et al., 2016)
Almada (Portugal)	14.6	8.4-24.1	2014	(Ryu et al., 2016)
Canberra (Australia)	14.6	9.3-22.3	2014	(Ryu et al., 2016)
Zurich (Switzerland)	14.7		2015	(Baz-Lomba et al., 2016)
Bristol (The United Kingdom)	16.2		2015	(Baz-Lomba et al., 2016)
Berlin (Germany)	16.9	13.8-22.3	2014	(Ryu et al., 2016)
Oslo (Norway)	16.1		2009	(Reid et al., 2011)
	18.9		2015	(Baz-Lomba et al., 2016)
	19.2	8.8-52.9	2014	(Ryu et al., 2016)
Barcelona (Spain)	18 ^a	7-31 ^a	2011-2015	(Mastroianni et al., 2017)
Dülmen (Germany)	20.3	5.5-40	2014	(Ryu et al., 2016)
London (United Kingdom)	21.5	10.9-36	2014	(Ryu et al., 2016)
Brussels (Belgium)	21.6		2015	(Baz-Lomba et al., 2016)
Eindhoven (The Netherlands)	21.7	13.7-30.4	2014	(Ryu et al., 2016)
Amsterdam (The Netherlands)	22	14.3-30.5	2014	(Ryu et al., 2016)
Castellón (Spain)	23.4	11.6-61.6	2014	(Ryu et al., 2016)
Dortmund (Germany)	23.6	18.1-34	2014	(Ryu et al., 2016)

Munich (Germany)	29.5	0.5-47.4	2014	(Ryu et al., 2016)
Dresden (Germany)	29.4	15.1-91.7	2014	(Ryu et al., 2016)
Montreal (Canada)	29.2	21.8-38.8	2014	(Ryu et al., 2016)
Copenhagen (Denmark)	29.7		2015	(Baz-Lomba et al., 2016)
	40.2	24.6-74	2014	(Ryu et al., 2016)
Granby (Canada)	44.3	27.3-59.3	2014	(Ryu et al., 2016)
Valencia-QB ^b	40.9 ^a	27.0-56.1 ^a	2014	(Andrés-Costa et al., 2016)

^aAlcohol consumption expressed in mL/day/inhabitant (aged 15+)

^bAlcohol consumption rate during “Fallas festivity”

Table S3. Comparison of alcohol consumption between pairs of investigated populations (U Mann Whitney test p-values)^a.

	Barcelona	Bilbao	Castellón	Guadalajara	Lleida	Madrid-La China	Madrid-Viveros	Móstoles	Palma de Mallorca	Reus	Santiago de Compostela	Tarragona	Toledo	Valencia-PI	Valencia-PII
Bilbao	0.114														
Castellón	0.095	0.020*													
Guadalajara	0.389	0.045*	0.209												
Lleida	0.114	0.012*	0.789	0.287											
Madrid-La China	0.148	0.012*	0.855	0.389	0.729										
Madrid-Viveros	1.000	0.075	0.237	0.601	0.114	0.171									
Móstoles	0.855	0.389	0.045*	0.095	0.075	0.075	0.534								
Palma de Mallorca	0.925	0.060	0.075	0.237	0.070	0.070	0.855	0.789							
Reus	0.729	0.114	0.662	0.662	0.237	0.729	0.789	0.287	0.662						
Santiago de Compostela	0.171	0.012*	0.925	0.348	1.000	0.855	0.148	0.075	0.060	0.601					
Tarragona	0.389	0.662	0.114	0.209	0.070	0.095	0.287	0.534	0.389	0.171	0.075				
Toledo	0.070	0.012*	0.237	0.171	0.237	0.534	0.070	0.075	0.045*	0.114	0.389	0.060			
Valencia-PI	0.459	0.045*	0.237	0.662	0.237	0.389	0.729	0.389	0.348	0.925	0.287	0.209	0.171		
Valencia-PII	0.925	0.012*	0.171	0.729	0.148	0.171	0.601	0.237	0.601	0.789	0.114	0.237	0.075	1.000	
Valencia-QB	0.171	0.012*	0.789	0.459	0.601	0.855	0.209	0.075	0.095	0.789	0.662	0.095	0.209	0.459	0.348

^aFirstly, a non-parametric test (Kruskal Wallis test) was applied in order to compare alcohol consumption among all investigated populations since the number of data per city was $n < 10$. Since $p < 0.05$, (Kruskal Wallis p-value = 0.0003887), the null hypothesis (H_0 : alcohol consumption among all investigated populations is equal) was rejected and a U Mann Whitney test was applied to compare alcohol consumption between pairs of populations. False Discovery Rate (FDR) correction for multiple testing was applied to reduce the number of “false positive”.

* $p < 0.05$, null hypothesis in U Mann Whitney test (H_0 : alcohol consumption between pairs of populations is equal) is rejected.

Table S4. Comparison of alcohol consumption between pairs of regions (U Mann Whitney test p-values)^a.

	Castilla-La Mancha	Catalonia	Community of Madrid	Valencian Community	Galicia	Balearic Islands
Catalonia	0.088					
Community of Madrid	0.088	1.000				
Valencian Community	0.286	0.335	0.200			
Galicia	1.000	0.169	0.096	0.221		
Balearic Islands	0.029 [*]	0.558	0.406	0.073	0.025 [*]	
Basque Country	0.001 [*]	0.073	0.020 [*]	<0.001 [*]	0.004 [*]	0.025 [*]

^aFirstly, a Kruskal Wallis test was applied in order to compare alcohol consumption among all investigated regions since for 3 regions (Galicia, Balearic Islands and Basque Country), $n < 10$. As $p\text{-value} < 0.05$ (Kruskal Wallis $p\text{-value} = 0.000588$), the null hypothesis (H_0 : alcohol consumption among all regions is equal) was rejected and a U Mann Whitney test was applied to compare alcohol consumption between pairs of regions. False Discovery Rate (FDR) correction for multiple testing was applied to reduce the number of “false positive”.

^{*} $p < 0.05$ and null hypothesis in U Mann Whitney (H_0 : alcohol consumption between pairs of regions is equal) is rejected.

Table S5. Average alcohol consumption estimated in Spain through WBE.

	Alcohol consumption in the investigated populations	Alcohol consumption in Spain				
	Kg/day	Kg/day	L/day	L/year/ inhabitants	L/year/ inhabitants (aged 15+)	L/year/ inhabitants (aged 18+)
Tuesday	48187	376424	477090	3.7	4.4	4.6
Wednesday	50115	391487	496181	3.9	4.6	4.8
Thursday	55403	432792	548532	4.3	5.1	5.3
Friday	57734	451005	571616	4.5	5.3	5.5
Saturday	84030	656420	831965	6.5	7.7	8.0
Sunday	77172	602852	764071	6.0	7.1	7.3
Monday	62306	486721	616884	4.8	5.7	5.9
<i>Average</i>	<i>62135</i>	<i>485386</i>	<i>615191</i>	<i>4.8</i>	<i>5.7</i>	<i>5.9</i>
<i>SD</i>	<i>13597</i>	<i>106216</i>	<i>134621</i>	<i>1.1</i>	<i>1.2</i>	<i>1.3</i>

Table S6. Average alcohol consumption (mL/day/inhabitant (aged 15+)) in the investigated regions in this study and Spain reported by the National Health Survey (INE).

	Week (Mon-Sun)		Weekdays (Mon-Thurs)		Weekend (Frid-Sun)	
	Average	sd	Average	sd	Average	sd
Balearic Island	18	14	15	14	22	17
Basque Country	19	14	11	15	30	19
Castilla-La Mancha	13	13	7.5	13	20	17
Catalonia	16	13	10	13	23	17
Community of Madrid	14	16	8.0	16	21	18
Galicia	20	12	16	13	25	13
Valencian Community	14	11	8.5	12	22	15
<i>Spain</i>	<i>13</i>	<i>12</i>	<i>8.4</i>	<i>12</i>	<i>19</i>	<i>16</i>

Source: National Health Survey (INE, 2017).

<https://www.ine.es/jaxi/Tabla.htm?path=/t15/p419/a2017/p03/10/&file=03011.px&L=0>



Figure S1. Map of Spain with the location of the sampled WWTPs (regions are indicated in different colors).




Figure S2. Weekly trends of alcohol consumption in the investigated populations.




Figure S3. Correlation between average alcohol consumption estimated in each city by WBE (mL/day/inhabitant) and prevalence data (“Lifetime prevalence”, “Last year prevalence” and “Last month prevalence”) reported by its region in the annual Report of the Spanish Observatory on Drugs and Drugs Addiction 2019. (Data from all investigated populations are shown; Spearman correlation p-values < 0.05 were considered statistically significant).




Figure S4. Prevalence data of alcohol consumption in the investigated regions and Spain reported in the Annual Report of the Spanish Observatory on Drugs and Drugs Addiction 2019.

References

- Andrés-Costa, M.J., Escrivá, Ú., Andreu, V., Picó, Y., 2016. Estimation of alcohol consumption during “Fallas” festivity in the wastewater of Valencia city (Spain) using ethyl sulfate as a biomarker. *Sci. Total Environ.* 541, 616–622. <https://doi.org/10.1016/j.scitotenv.2015.09.126>
- Baz-Lomba, J.A., Salvatore, S., Gracia-Lor, E., Bade, R., Castiglioni, S., Castrignanò, E., Causanilles, A., Hernandez, F., Kasprzyk-Hordern, B., Kinyua, J., McCall, A.-K., van Nuijs, A., Ort, C., Plósz, B.G., Ramin, P., Reid, M., Rousis, N.I., Ryu, Y., de Voogt, P., Bramness, J., Thomas, K., 2016. Comparison of pharmaceutical, illicit drug, alcohol, nicotine and caffeine levels in wastewater with sale, seizure and consumption data for 8 European cities. *BMC Public Health* 16, 1035. <https://doi.org/10.1186/s12889-016-3686-5>
- Gatidou, G., Kinyua, J., van Nuijs, A.L.N., Gracia-Lor, E., Castiglioni, S., Covaci, A., Stasinakis, A.S., 2016. Drugs of abuse and alcohol consumption among different groups of population on the Greek Island of Lesbos through sewage-based epidemiology. *Sci. Total Environ.* 563–564, 633–640. <https://doi.org/10.1016/j.scitotenv.2016.04.130>
- Mastroianni, N., López-García, E., Postigo, C., Barceló, D., López de Alda, M., 2017. Five-year monitoring of 19 illicit and legal substances of abuse at the inlet of a wastewater treatment plant in Barcelona (NE Spain) and estimation of drug consumption patterns and trends. *Sci. Total Environ.* 609, 916–926. <https://doi.org/10.1016/j.scitotenv.2017.07.126>
- Nguyen, H.T., Thai, P.K., Kaserzon, S.L., O'Brien, J.W., Eaglesham, G., Mueller, J.F., 2018. Assessment of drugs and personal care products biomarkers in the influent and effluent of two wastewater treatment plants in Ho Chi Minh City, Vietnam. *Sci. Total Environ.* 631–632, 469–475. <https://doi.org/10.1016/j.scitotenv.2018.02.309>
- Reid, M.J., Langford, K.H., Mørland, J., Thomas, K. V., 2011. Analysis and Interpretation of Specific Ethanol Metabolites, Ethyl Sulfate, and Ethyl Glucuronide in Sewage Effluent for the Quantitative Measurement of Regional Alcohol Consumption. *Alcohol. Clin. Exp. Res.* no-no. <https://doi.org/10.1111/j.1530-0277.2011.01505.x>
- Rodríguez-Álvarez, T., Racamonde, I., González-Mariño, I., Borsotti, A., Rodil, R., Rodríguez, I., Zuccato, E., Quintana, J.B., Castiglioni, S., 2015. Alcohol and cocaine co-consumption in two European cities assessed by wastewater analysis. *Sci. Total Environ.* 536, 91–98. <https://doi.org/10.1016/j.scitotenv.2015.07.016>
- Rodríguez-Álvarez, T., Rodil, R., Cela, R., Quintana, J.B., 2014. Ion-pair reversed-phase liquid chromatography-quadrupole-time-of-flight and triple-quadrupole-mass spectrometry determination of ethyl sulfate in wastewater for alcohol consumption tracing. *J. Chromatogr. A* 1328, 35–42. <https://doi.org/10.1016/j.chroma.2013.12.076>
- Ryu, Y., Barceló, D., Barron, L.P., Bijlsma, L., Castiglioni, S., de Voogt, P., Emke, E., Hernández, F., Lai, F.Y., Lopes, A., López de Alda, M., Mastroianni, N., Munro, K., O'Brien, J., Ort, C., Plósz, B.G., Reid, M.J., Yargeau, V., Thomas, K. V., 2016. Comparative measurement and quantitative risk assessment of alcohol consumption through wastewater-based epidemiology: An international study in 20 cities. *Sci. Total Environ.* 565, 977–983. <https://doi.org/10.1016/j.scitotenv.2016.04.138>