This document is the Accepted Manuscript version of a Published Work that appeared in final form in *Revista Iberoamericana de Micologia*, July 2020.

Online version:

https://www.sciencedirect.com/science/article/abs/pii/S1130140620300322

DOI: https://doi.org/10.1016/j.riam.2020.07.002

# Identification of *Mucor circinelloides* antigens recognized by sera from immunocompromised infected mice

Maialen Areitio<sup>1</sup>, Aitana Arbizu<sup>1</sup>, Adela Martin-Vicente<sup>2</sup>, Aitziber Antoran<sup>1</sup>, Leire Aparicio-Fernandez<sup>1</sup>, Idoia Buldain<sup>1</sup>, Leire Martin-Souto<sup>1</sup>, Aitor Rementeria<sup>1</sup>, Javier Capilla<sup>2</sup>, Fernando L. Hernando<sup>1</sup> and Andoni Ramirez-Garcia<sup>1</sup>

<sup>1</sup>Fungal and Bacterial Biomics Research Group, Department of Immunology, Microbiology and Parasitology. Faculty of Science and Technology. University of the Basque Country (UPV/EHU), Leioa 48940, Spain.

<sup>2</sup>Mycology Unit, Medical School and IISPV, Universitat Rovira i Virgili, Reus, Spain.

**Corresponding author:** Fernando L. Hernando (tel.: +34946015407, E-mail address: fl.hernando@ehu.es)

Keywords: proteomics, antigen, fungal disease, secreted, characterization

### ABSTRACT

*Mucor circinelloides* is an opportunistic fungus capable of causing mucormycosis, a highly aggressive infection of quick spreading. Besides, it also has a high mortality rate due to late diagnosis and difficult treatment.

Aims: In this study we have identified the most immunoreactive proteins of the secretome and the total protein extract of *M. circinelloides* using sera from immunocompromised infected mice.

Methods: The proteins of the secretome and the total extract were analyzed by two-dimensional electrophoresis and the most immunoreactive antigens were detected by Western Blot, facing the sera of immunocompromised infected mice to the proteins obtained in both extracts of *M. circinelloides*.

Results: Seven antigens were detected in the secretome extract, and two in the total extract, all of them corresponding only to three proteins. The enzyme enolase was detected in both extracts, while triosephosphate isomerase was detected in the secretome, and heat shock protein HSS1 in the total extract.

Conclusions: In this work the most immunoreactive antigens of the secretome and the total extract of *M. circinelloides* were identified. The identified proteins are well known fungal antigens and, therefore, these findings can be useful for future research into alternatives for the diagnosis and treatment of mucormycosis.

#### INTRODUCTION

Mucor circinelloides is a saprophyte and pathogenic fungus which affects, above all, immunocompromised individuals, both human and animal. According to a recent review of cases of mucormycosis between January 2000 and January 2017, Mucor species are the second most common causal agent of mucormycosis, only outnumbered by fungi within the genus Rhizopus.<sup>9</sup> The incidence of mucormycosis has been constantly increasing, the principal cause being the marked rise in debilitating diseases of the immune system, which includes pathologies such as bone marrow or solid organ transplantations, immunodeficiencies related to HIV, diabetes and primary or iatrogenic neutropenia.<sup>10</sup> Recently, it has also caused an outbreak in burned patients.<sup>6</sup> Mucormycosis can occur by inhalation of airborne conidia or inoculation on an open wound,<sup>18</sup> it is highly aggressive and spreads quickly, invading the blood vessels, causing hemorrhages, thrombosis, heart attacks and tissue necrosis. In addition, it must be taken into account the fact that diagnosis is usually carried out at a late stage of infection, and thattreatment strategies arenot clear.<sup>21,22</sup> Because of these reasons outlined above and the fact that it is an emerging pathology, more research into this disease is of paramount importance.<sup>7</sup> Therefore, the aim of this work was to achieve the identification of the most immunogenic antigens of *M. circinelloides*. These findings are useful for future researches on alternatives for diagnosis and treatment of mucormycosis

#### MATERIALS AND METHODS

#### Strains and culture conditions

The strain of *M. circinelloides* used in this study was CBS 125721, isolated from human maxillary tissue. It was cryopreserved at  $-80 \circ$ C and cultured onto Potato Dextrose Agar (PDAat 28  $\circ$ C for 7 days before use. The conidiospores were obtained by washing the plates with sterile saline (0.9% NaCl), filtered through gauze and centrifuged. The concentration of conidiospores was adjusted as needed using a hemocytometer.

### Animal models

The sera used for the detection of the most immunoreactive antigens were obtained from two groups of 5 mice each immunosupressed with cyclophosphamide: one was infected with fungal conidia, and the other one was injected with saline solution, according to a method previously described.13 Briefly, the mice were immunosuppressed 2 days prior the infection by intraperitoneal injection of 200 mg/kg body weight of cyclophosphamide and once every 5 days thereafter, and inoculated intravenously with a density of 105 conidia or saline solution and sacrificed 20 days later. All experiments involving animals were in accordance with the ethical standards of Universitat Rovira i Virgili Animal Welfare and Ethics Comittee.

### Obtention of total protein extract and secretome of M. circinelloides

For the total extract and the secretome,  $5 \times 10^5$  conidia/ml and  $10^6$  conidia/ml were grown, respectively, for 24 h at 37 °C and 120 rpm in 150 ml Potato Dextrose Broth (PDB). After culture, both conidia and hyphae morphologies were obtained. Then, the fungus was centrifuged (12,000 × g, 5 min, 4 °C), and washed with PBS. For the total extract the pellet was resuspended in PBS with 1% -mercaptoethanol and 1% pharmalytes, then it was lysed using crystal beads (0.5 mm diameter) at 30 Hz for 20 min using the Millmix 20 Bead-Beater (Thetnica, Eslovenia, Europe). The secretome was obtained as described by da Silva et al.<sup>5</sup> for *Scedosporium boydii*, with a few modifications. After growing in PDB, the fungus was collected and cultured in PBS-

2% glucose for 20 h at 37 °C and 120 rpm. Finally, the obtained culture was centrifuged at 11.000 × g for 20 min and the supernatant was sterilized through 0.22 m filters (Merck Millipore, MAS, USA). To confirm the absence of cytoplasmic proteins in the secretome the activity of the intracellular enzyme lactate dehydrogenase (LDH) was measured, as previously described,<sup>23</sup> and the cellular integrity of the fungus was observed under optical microscopy.

# Protein detection by two-dimensional electrophoresis

The proteins were precipitated and two-dimensional electrophoresis was carried out following the protocol described in a previous work,<sup>15</sup> using 18 cm long Immobiline DryStrip gels (pH 3– 10, GE Healthcare). The conditions of the isoelectric focusing were modified for the secretome: 12 h rehydration, 150V for 300Vhr (Step and Hold, S&H), 500V for 2000Vhr (S&H), 1000V for 9000Vhr (gradient), 8000V for 20,000Vhr (gradient) and 8000V for 100,000Vhr (S&H). The second dimension was carried outin the PROTEAN II xi Cell system (Bio-Rad, CA, USA) at 45 mA using 10% polyacrylamide gels. The 2-DE gels were stained with Coomassie Brilliant Blue to visualize proteins.

# Antigen detection

Proteins were electrotransferred to Hybond-P PVDF membranes, which were stained with Ponceau red to check the correct transfer of the proteins. Then, Western Blot was performed in accordance with the protocol described by Pellon et al.<sup>15</sup> using the sera of infected mice, pooled and diluted 1/100 as sample. The experiments were carried out in triplicate and the most immunoreactive antigens were selected using ImageMaster 2D Platinum software (GE Healthcare). Identification of immunoreactive proteins The identified spots were extracted from the gels for identification by mass spectrometry (LC–MS/MS)in the service of proteomics SGlker of the University of the Basque Country (UPV/EHU), as described by Buldain et al.<sup>2</sup> Briefly, the extracted gel pieces were swollen in digestion buffer (50 mM NH4HCO3, 12.5 ng/l trypsin and incubated at 37 °C overnight. After that, the peptides were extracted, first with 25 mM NH<sub>4</sub>HCO<sub>3</sub> and acetonitrile, and secondly with 0.1% (v/v) trifluoroacetic acid and ACN. LC–MS/MS was carried out on a SYNAPT HDMS mass spectrometer (Waters, Milford, MA, USA)interfaced with a nanoAcquity UPLC System (Waters). The search for protein identification was made in the non-redundant database of the NCBI, restricted to fungi, using the online server MASCOT (Matrix Science Ltd., London, UK (http://www.matrixscience.com).

# Bioinformatic analysis of M. circinelloides antígens

In order to study the relation between the identified antigens and their homologues in other organisms, the BLAST database (https://blast.ncbi.nlm.nih.gov/Blast.cgi) was used. Then, the similarity values found between the proteins were compared.

# RESULTS

# Inoculation of immunosuppressed mice with *M. circinelloides*

In this study, mice immunosuppressed with cyclophosphamide were used to reduce the immune response and, in consequence, to avoid excessive immunoreactivity in WBs. Besides, we selected a strain of *M. circinelloides* previously classified as avirulent, which allowed us to collect the sera from all mice as the mortality rate in both infected and non-infected groups was zero. All the data of the infection process using this avirulent strain were described in a previous study.<sup>13</sup>

# Identification of the immunoreactive antigens of the secretome recognized by sera of immunocompromised infected mica

The proteomics study of the secretome of *M. circinelloides* by 2- DE showed that secreted proteins were localized throughout the whole isoelectric point (pl) range used and with molecular weights (Mr) smaller than 130 kDa. Specifically, the majority of the proteins contained in the secretome presented a pl between 4.5 and 7, and a Mr between 25 and 70 kDa (Fig. 1a). Sera of immunocompromised infected mice were used against the obtained extract; the immunoreactivity was analyzed and the sevenmostimmunoreactive antigens were identified bymass spectrometry (Fig. 1). Three of these spots corresponded to enolase and four to triosephosphate isomerase, as shown in Table 1. They are almost certainly isoforms of the enolase and triosephosphate isomerase, respectively, as they have the same molecular weights and very similar isoelectric points, as it can be observed in Fig. 1a. The sera of mice injected with saline solution were also used at the same dilution and, as expected, no reactivity was observed (data not shown)

# Identification of the immunoreactive antigens of the total extract recognized by sera of immunocompromised infected mica

Regarding the study of the fungal cell proteome of the total extract by 2-DE, the proteins detected were also localized on the whole range of pl and Mr used. However, the protein distribution pattern observed was very different from the secretome and almost all proteins were present in the ranges of pl of 5–9 and Mr of 40–70 kDa (Fig. 2a). The immunome detected by two-dimensional WB showed a very few number of immunoreactive proteins recognized by sera from immunosuppressed mice and, therefore, only the two most immunoreactive spots were selected for their further identification(Fig. 2). These proteins were identified bymass spectrometry as enolase andheat shock protein HSS1 (Table 2).Inthe same way than for the secretome, the sera of mice injected with saline solution did not show any immunoreactivity (data not shown).

# Bioinformatic analysis of the identified proteins

When comparing the sequence of the identified triosephosphate isomerase with the genome of *M. circinelloides*, an identity value of 99.48% was found. The Hsp70 identified in our work has similarity values of 78.58%, 76.43% and 66.39% with the proteins SSA1 and SSA2 found in Candida albicans, and the Hsp70 found in Rhizopus arrhizus, respectively. Finally, the sequence of the enolase of *M. circinelloides* was compared with the enolase of *C. albicans*, *R. arrhizus* and *Homo sapiens*, and similarity values of 69%, 81.88% and 71% were found, respectively.

### DISCUSSION

Due to the increase of immune system debilitating diseases in the last years, the prevalence of many infections has arisen. Mucormycosis is one ofthese infectious diseases, which can be caused by the fungus *M. circinelloides*. Mucormycosis is known for its high aggressiveness and quick spreading. Besides, due to the late diagnosis in the majority of the cases and the presence of high percentage of resistant isolates, this disease has high mortality rates. Therefore, the aim of this work was to identify the most immunoreactive antigens of *M. circinelloides*, analyzing the immune response produced by sera obtained from immunocompromised infected mice. We used a cyclophosphamide-based immunosuppressive treatment to reduce the humoral immune response and, therefore, avoid the excess of signal which might mask and difficult the interpretation of the results. In this way, we proceeded to the identification of only the most

immunogenic antigens by LC–MS/MS. As a result, we identified in the secretome extract, from seven immunoreactive spots, two proteins, being them enolase and triosephosphate isomerase; in the total extract, from two spots, two proteins were identified, enolase and heat-shock protein HSS1. Among the proteins identified in the secretome extract, three isoforms of enolase and four of triosephosphate isomerase were detected. The spectrometric identifications showed that the proteins are probably the product of the same gene and, therefore, the little differences found among the pl values of these proteins might be due to slight modifications during the secretion process. As enolase and triosephosphate isomerase, usually cytoplasmic proteins, were detected in the secretome, the mechanism of their secretion and function are not completely understood. The secretion process involved in M. circinelloides could be through extracellular vesicles, as seen with triosephosphate isomerase of the dimorphic fungus Paracoccidioides brasiliensis. <sup>12</sup> Besides, taking into account the plasminogen binding activities of these two proteins, their presence in the external medium could be related to pathogen dissemination.<sup>4</sup> Although triosephosphate isomerase was detected in the secretome extract of *M. circinelloides*, the identification by LC–MS/MS of the selected spot resulted in the fungus of the same genus Mucor ambiguus, because the used database determined this protein as the one found in this microorganism. When comparing this protein sequence with the genome of *M. circinelloides*, an identity value of 99.48% was found, indicating the high degree of similarity between the sequences of triosephosphate isomerase in both species. This protein was also identified in the fungi C. albican and P. brasiliensis, and this is considered an important antigen, capable of binding the laminin and fibronectin of the extracellular matrix<sup>17</sup> Moreover, it can be found in the cell wall, but also in the secreted extracellular vesicles.<sup>12</sup> Regarding HSS1, it belongs to the same heat shock protein family as the Hsp70,<sup>1</sup> which in fungal species is overexpressed during infection to prevent he denaturalization of proteins and has been described as immunoreactive against serum IgGs15 and mucosal IgAs.<sup>16</sup> In fact, HSS1 could be considered an orthologue of the Hsp70 proteins, SSA1 and SSA2, found in the ascomycetous fungi C. albicans, and the Hsp70 of *R. arrhizus*, as it shares with those proteins similarity values higher than 65%. Although there is a high percentage of homology, there are some specific regions when compared with the sequences of SSA1 of C. albicans and Hsp70 of R. arrhizus. This fact could be of great interest for diagnosis. Heat-shock proteinshave beenreported as antigens and virulence factors in he cell surface15 and the secretome of important pathogenic fungi, such as Lomentospora prolificans.<sup>3</sup> In this sense, the location of the protein could ease the access of the immune cells, contributing to the high immunogenicity presented. Besides, it has been proved that the Hsp70 of Mycobacterium tuberculosis is capable of inducing the immune response, stimulating human monocytes to produce chemokines and cytokines.<sup>24</sup> Given the major functions that Hsp70 has on infective processes and the homology between this protein and the one found in our study, it could be hypothesized that the protein HSS1 could have a similar role in the infections caused by *M. circinelloides*. Enolase deserves a special mention as it has been identified in both analyzed protein extracts. Enolase has been also identified in the secretome of the mycelia and yeast cells of the dimorphic fungus Paracoccidioides lutzii.<sup>25</sup> This metabolic enzyme was also previously identified by our research group as an antigen recognized by salivary IgA in L. prolificans, <sup>2</sup> and by serum IgGs in C. albicans. <sup>8</sup> Enolase was also associated with the cell wall in *L. prolificans*<sup>15</sup> and C. albicans, <sup>14</sup> where it performs transglutaminase activity, indicating the possible role of this enzyme in infection processes.<sup>20</sup> Enolase, therefore, seems to be an important antigen for many of the most common pathogenic fungi, which makes it a reasonable target in panfungal diagnosis or a key to design new treatments or vaccines. In fact, the use of this enzyme for therapy was tested as a vaccine against C. albicans, lowering both the fungal burden and the amount of tissue damage.<sup>11</sup> In order to determine the possible use of this protein in diagnosis or treatment strategies, its sequence was compared with the sequence in *C. albicans, R. arrhizus* and *H. sapiens*, and the similarity values found make probable a cross-reactivity with other fungi, as it happens between enolase of *L. prolificans* and *Scedosporium apiospermum, S. boydii* and *Aspergillus fumigatus*. <sup>14</sup> Hence, although it could not be used for a specific diagnosis or therapeutic target, it could be used as a panfungal antigen. In conclusion, in this work the most immunoreactive antigens of the secretome and the total extract of *M. circinelloides* were identified. The proteins identified were the HSS1 protein, along with enolase and triosephosphate isomerase, which are well-known fungal antigens. These proteins might be useful in the future for the development of a vaccine, antifungaltreatments and/or for diagnosis, allowing the rapid detection and treatment of the disease and, therefore, lowering the unacceptable mortality rates.

## **Authors contributions**

All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by MA, LMS and ARG. The first draft of the manuscript was written by MA and all authors commented on previous versions of the manuscript. All authors read and approved the final version.

•	5		_		-			
Spot					())) <b>(</b>		Theorethical	Experimental
number		Name of the protein	MICLOOLGANISH	Matches	COVER (%)	score	pl/Mr(kDa)	pl/Mr(kDa)
Secretome								
-	EPB85979.1	Enolase	Mucor circinelloides	10	34	855	5,56/47,17	5,44/63,14
7	EPB85979.1	Enolase	Mucor circinelloides	14	36	1093	5,56/47,17	5,65/63,66
ę	EPB85979.1	Enolase	Mucor circinelloides	4	12	317	5,56/47,17	5,76/63,66
4	GAN00908.1	Triose phosphate isomerase	Mucor ambiguus	9	27	319	5,26/26,95	5,23/26,28
5	GAN00908.1	Triose phosphate isomerase	Mucor ambiguus	10	32	537	5,26/26,95	5,00/26,57
9	GAN00908.1	Triose phosphate isomerase	Mucor ambiguus	14	48	807	5,26/26,95	5,24/26,79
7	GAN00908.1	Triose phosphate isomerase	Mucor ambiguus	11	40	663	5,26/26,95	5,50/26,79
Total extract	t							
-	EPB85979.1	Enolase	Mucor circinelloides	45	46	1346	5,56/47,17	5,79/60,88
2	EPB91082.1	Heat-shock protein HSS1	Mucor circinelloides	23	33	1530	5,07/70,92	5,79/89,23





## References

1. Acioli-Santos B, Sebastiana M, Pessoa F, Sousa L, Figueiredo A, Fortes AM, et al. Fungal transcript pattern during the preinfection stage (12 h) of ectomycorrhiza formed between Pisolithus tinctorius and Castanea sativa roots, identified using cDNA microarrays. Curr Microbiol. 2008;57:620–5.

2. Buldain I, Ramirez-Garcia A, Pellon A, Antoran A, Sevilla MJ, Rementeria A, et al. Cyclophilin and enolase are the most prevalent conidial antigens of Lomentospora prolificans recognized by healthy human salivary IgA and cross-react with Aspergillus fumigatus. PROTEOMICS – Clin Appl. 2016;10:1058–67.

3. Buldain I, Pellon A, Zaldibar B, Antoran A, Martin-Souto L, Aparicio-Fernandez L, et al. Study of humoral responses against Lomentospora/Scedosporium spp. and Aspergillus fumigatus to identify L. prolificans antigens of interest for diagnosis and treatment. Vaccines (Basel). 2019;7:212–34.

4. Chaves EG, Weber SS, Báo SN, Pereira LA, Bailão AM, Borges CL, et al. Analysis of Paracoccidioides secreted proteins reveals fructose 1,6-bisphosphate aldolase as a plasminogen-binding protein. BMC Microbiol. 2015;15:53–67.

5. Da Silva BA, Sodré CL, Souza-Gonc, alves AL, Aor AC, Kneipp LF, Fonseca BB, et al. Proteomic analysis of the secretions of Pseudallescheria boydii, a human fungal pathogen with unknown genome. J Proteome Res. 2012;11:172–88.

6. Garcia-Hermoso D, Criscuolo A, Lee SC, Legrand M, Chaouat M, Denis B, et al. Outbreak of invasive wound mucormycosis in a burn unit due to multiple trains of Mucor circinelloides f. circinelloides resolved by whole-genome sequencing. MBio. 2018;9:e00673–718.

7. Gomes MZR, Lewis RE, Kontoyiannis DP. Mucormycosis caused by unusual mucormycetes, non-Rhizopus, -Mucor, and -Lichtheimia species. Clin Microbiol Rev. 2011;24:411–45.

8. Hernando FL, Calvo E, Abad A, Ramirez A, Rementeria A, Sevilla MJ, et al. Identification of protein and mannoprotein antigens of Candida albicans of relevance for the serodiagnosis of invasive candidiasis. Int Microbiol. 2007;10:103–8.

9. Jeong W, Keighley C, Wolfe R, Lee WL, Slavin MA, Kong DCM. Chen SC-A. The epidemiology and clinical manifestations of mucormycosis: a systematic review and meta-analysis of case reports. Clin Microbiol Infect. 2019;25:26–34.

10. Lee SC, Billmyre RB, Li A, Carson S, Sykes SM, Huh EY, et al. Analysis of a foodborne fungal pathogen outbreak: virulence and genome of a Mucor circinelloides isolate from yogurt. MBio. 2014;5:1–12.

11. LiWQ, Hu XC, Zhang X, Ge Y, Zhao S, Hu Y, et al. Immunisation with the glycolytic enzyme enolase confers effective protection against Candida albicans infection in mice. Vaccine. 2011;29:5526–33.

12. Longo LVG, da Cunha JPC, Sobreira TJP, Puccia R. Proteome of cell wall-extracts frompathogenic Paracoccidioides brasiliensis:Comparisonamongmorphological phases, isolates, and reported fungal extracellular vesicle proteins. EuPA Open Proteomics. 2014;3:216–28.

13. Lopez-Fernandez L, Sanchis M, Navarro-Rodríguez P, Nicolás FE, Silva-Franco F, Guarro J, et al. Understanding Mucor circinelloides pathogenesis by comparative genomics and phenotypical studies. Virulence. 2018;9:707–20.

14. Maresca B, Kobayashi GS. Hsp70 in parasites: as an inducible protective protein and as an antigen. Experientia. 1994;50:1067–74.

15. Pellon A, Ramirez-Garcia A, Buldain I, Antoran A, Rementeria A, Hernando FL. Immunoproteomics-based analysis of the immunocompetent serological response to Lomentospora prolificans. J Proteome Res. 2016;15:595–607.

16. Pellon A, Ramirez-Garcia A, Antoran A, Fernandez-Molina JV, Abad-Diaz-DeCerio A, Montanez ~ D, et al. Scedosporium prolificans immunomes against human salivary immunoglobulin A. Fungal Biol. 2014;118:94–105.

17. Pereira LA, Báo SN, Barbosa MS, da Silva JLM, Felipe MSS, de Santana JM, et al. Analysis of the Paracoccidioides brasiliensis triosephosphate isomerase suggests the potential for adhesin function. FEMS Yeast Res. 2007;7:1381–8.

18. Petrikkos G, Skiada A, Lortholary O, Roilides E, Walsh TJ, Kontoyiannis DP. Epidemiology and clinical manifestations of mucormycosis. Clin Infect Dis. 2012;54 Suppl. 1:23–34.

19. Pitarch A, Díez-Orejas R, Molero G, Pardo M, Sánchez M, Gil C, et al. Analysis of the serologic response to systemic Candida albicans infection in a murine model. Proteomics. 2001;1:550–9.

20. Reyna-Beltran E, Iranzo M, Calderón-González KG, Mondragón-Flores R, LabraBarrios ML, Mormeneo S, et al. The Candida albicans ENO1 gene encodes a transglutaminase involved in growth, cell division, morphogenesis and osmotic protection. J Biol Chem. 2018;293:4304–23.

21. Roden MM, Zaoutis TE, Buchanan WL, Knudsen TA, Sarkisova TA, Schaufele RL, et al. Epidemiology and outcome of zygomycosis: a review of 929 reported cases. Clin Infect Dis. 2005;41:634–53.

22. Schwarz P, Bretagne S, Gantier JC, García-Hermoso D, Lortholary O, Dromer F, et al. Molecular identification of zygomycetes from culture and experimentally infected tissues. J Clin Microbiol. 2006;44:340–9.

23. Silva BA, Pinto MR, Soares RMA, Barreto-Bergter E, Santos ALS. Pseudallescheria boydii releases metallopeptidases capable of cleaving several proteinaceous compounds. Res Microbiol. 2006;157:425–32.

24. Wang Y, Kelly CG, Singh M, McGowan EG, Carrara AS, Bergmeier LA, et al. Stimulation of Th1-polarizing cytokines. C–C chemokines, maturation of dendritic cells, and adjuvant function by the peptide binding fragment of heat shock protein 70. J Immunol. 2002;169:2422–9.

25. Weber SS, Parente AF, Borges CL, Parente JA, Bailão AM, de Almeida Soares CM. Analysis of the secretomes of Paracoccidioides mycelia and yeast cells. PLoS One. 2012;7:e52470