

Dietary procyanidins lower triglyceride levels signaling through the nuclear receptor small heterodimer partner

Authors:

Josep Maria Del Bas(1), Marie Louise Ricketts(2), Isabel Baiges(1), Helena Quesada(1), Anna Ardevol(1), Maria Josepa Salvadó(1), Gerard Pujadas(1), Mayte Blay(1), Lluís Arola(1), Cinta Bladé(1), David D. Moore(2) and Juan Fernandez-Larrea(1)

(1) Departament de Bioquímica i Biotecnologia, Universitat Rovira i Virgili, Campus Sescelades, Tarragona, Spain

(2) Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, TX, USA

Correspondence:

Dr. Cinta Bladé, Departament de Bioquímica i Biotecnologia, Universitat Rovira i Virgili, Campus Sescelades s/n, 43007 Tarragona, Spain

E-mail: mariacinta.blade@urv.cat

Fax: +34977558232

Abbreviations:

ApoA5, apolipoprotein A5; ApoB, apolipoprotein B; CPT-1A, carnitine palmitoyltransferase-1A; FA, fatty acid; GSPE, grape seed procyanidin extract; NR0B2/SHP, nuclear receptor small heterodimer partner; SREBP-1c, steroid response element binding protein 1c; TG, triglyceride

Abstract

Hypertriglyceridemia is an independent risk factor in the development of cardiovascular diseases, and we

have previously reported that oral administration of a grape seed procyanidin extract (GSPE) drastically decreases plasma levels of triglycerides (TG) and apolipoprotein B (ApoB) in normolipidemic rats, with a concomitant induction in the hepatic expression of the nuclear receptor small heterodimer partner (NR0B2/SHP). Our objective in this study was to elucidate whether SHP is the mediator of the reduction of TG-rich ApoB-containing lipoproteins triggered by GSPE. We show that GSPE inhibited TG and ApoB secretion in human hepatocarcinoma HepG2 cells and had a hypotriglyceridemic effect in wild-type mouse. The TG-lowering action of GSPE was abolished in HepG2 cells transfected with a SHP-specific siRNA and in a SHP-null mouse. Moreover, in mouse liver, GSPE downregulated several lipogenic genes, including steroid response element binding protein 1c (SREBP-1c), and upregulated carnitine palmitoyltransferase-1A (CPT-1A) and apolipoprotein A5 (ApoA5), in a SHP-dependent manner. In HepG2 cells GSPE also inhibited ApoB secretion, but in a SHP-independent manner. In conclusion, SHP is a key mediator of the hypotriglyceridemic response triggered by GSPE. This novel signaling pathway of procyanidins through SHP may be relevant to explain the health effects ascribed to the regular consumption of dietary flavonoids.

Keywords: Liver / NR0B2/SHP / Procyanidins / SREBP-1c / Triglycerides

1. Introduction

The regular long-term consumption of dietary flavonoids has been associated with reduced mortality and risk of cardiovascular diseases (CVD) [1-3]. Hypertriglyceridemia is an important contributing factor for atherosclerosis and represents an independent risk factor for CVD [4-8] and several studies have shown beneficial effects of flavonoids regarding atherosclerosis prevention, reducing apolipoprotein B (ApoB), and triglyceride (TG) levels in vivo [9, 10] and in vitro [11, 12]. Using in vitro models, it has been demonstrated that specific flavonoids decrease hepatocyte ApoB secretion acting through phosphoinositide-3 kinase (PI3K) and mitogen-activated protein kinase/extracellular signal-regulated kinase (MAPK/ERK) signaling pathways [13, 14]. In contrast, there are not studies about the mechanisms

used by flavonoids to reduce TG production and secretion by hepatic cells.

We have previously shown that oral administration of procyanidins, a class of flavonoids, drastically reduce the postprandial levels of TG-rich, ApoB-containing lipoproteins, and improve different atherosclerotic risk indexes in normolipidemic rats [15]. These changes in plasma lipid profile were paralleled by a significant increase in the expression of the orphan nuclear receptor small heterodimer partner (NR0B2/SHP) in the liver. SHP is involved in bile acid, cholesterol, TG, and glucose homeostasis [16-18] and mediates the hypotriglyceridemic effect of bile acids [19] and guggulsterone [20]. Thus, we hypothesized that SHP could be a key mediator in the hypotriglyceridemic effect triggered by procyanidins.

In this work we demonstrate that GSPE reduces TG secretion in the human hepatocarcinoma cell line HepG2 and has a hypotriglyceridemic effect in wild-type mouse. This hypotriglyceridemic effect of GSPE is abolished in two SHP deficient models: HepG2 cells transfected with a SHP-specific siRNA and a SHP knockout mouse, thus showing this nuclear receptor as a key mediator of the hypotriglyceridemic response triggered by procyanidins in the liver. Additionally, we have identified SHP target genes, including the steroid receptor element-binding protein 1c (SREBP1c), as putative downstream effectors of the TG lowering response triggered by procyanidins in hepatic cells in vivo.

2. Materials and methods

2.1 Chemicals

Grape seed procyanidin extract (GSPE) was kindly provided by Les Dérives Résiniques et Terpéniques (Dax, France). According to the manufacturer, this procyanidin extract contains monomeric (16.55%), dimeric (18.77%), trimeric (16%), tetrameric (9.3%), oligomeric (5-13 units) (35.7%) procyanidins, and phenolic acids (4.22%).

2.2 Cell cultures and cytotoxicity assays

HepG2 cells (ATCC code HB-8065) were grown until 70% confluence in DMEM (Cambrex) medium supplemented with 10% FBS (Cambrex). Twelve hours before GSPE treatment, media was replaced with serum-depleted media (DMEM supplemented with 0.25% oleic acid: albumin) (Sigma). Cytotoxicity of GSPE was assessed by measuring LDH leakage as described [21] and by the Alamar Blue assay (Biosource, USA), in cells treated 24 h with different concentrations of GSPE diluted in aqueous solution.

2.3 In vivo feeding studies

Mice were housed under standard conditions. Experimental procedures were approved by the local Committee for Care and Use of Laboratory Animals at Baylor College of Medicine. SHP-deficient mice were generated by gene targeting as previously described [22], and were backcrossed with C57BL6 mice to the tenth generation. The correct genotype was verified using previously reported primer sequences and reaction conditions [23]. Age-matched groups of 8-10 wk-old male mice were used in all experiments (n=5 per experimental group). Mice were fed a standard rodent chow and water ad libitum. On experimental day, mice were fed either vehicle (water), or procyanidins (250 mg/kg) via oral gavage. A first dose was administered at 9:00 pm and a second dose at 9:00 am; food was then retired and mice were sacrificed 2 h later. Blood was collected from the orbital plexus after isoflurane anesthesia. Livers were snap-frozen and stored at -80°C until use.

2.4 Lipid analysis

To study the synthesis of TG, HepG2 cells were seeded in 12-well plates and grown to 70% confluence. Growth medium was replaced by serum free medium 12 h before treatment with GSPE (50 mg/L) or vehicle. Twelve hours after the simultaneous addition of GSPE (50 mg/L) and ¹⁴C-acetate (1 µCi/well), media and cells were collected and lipids extracted using chloroform/methanol. TLC was performed as previously described [24] with an additional separation using a Hexane/MTBE/NH₃ solvent to obtain the TG fraction. ¹⁴C-labeled TG were scraped and determined by scintillation counting. Values were normalized per mg of total cellular protein, determined by a colorimetric assay (BioRad). For

nonradioactive measurements, cell media and plasma TG or cholesterol were assayed using enzymatic kits (QCA).

2.5 Quantification of ApoB and SHP proteins

ApoB protein secreted by HepG2 cells into the culture media was determined by immunodetection using Western blot analysis. At the end of the treatments, media were collected and concentrated using Ultrafree-4 centrifugal filter units (Millipore). Equal volumes of media were subjected to SDS-PAGE and transferred to a nitrocellulose membrane (Millipore, USA). ApoB was immunodetected using an anti-apoB antibody (S-14, Santa Cruz Biotechnology) as the primary antibody, HRP-conjugated anti-IgG (sc-2020, Santa Cruz Biotechnology) as the secondary antibody, and immobilon chemiluminescent HRP substrate kit (Millipore, USA) as the chemiluminescent reagent. Membranes were exposed to Hyperfilm ECL (Amersham), and ApoB-specific signals quantified using Quantity One software (BioRad); intensities values were always normalized per cell number. For quantification of SHP protein, HepG2 cells were lysed using RIPA buffer (50 mM Tris-HCl, pH 7.4, 1% NP-40, 0.25% Na-deoxycholate, 150 mM NaCl, 1 mM EDTA) containing a protease-inhibitor cocktail (Sigma). Protein was quantified using a protein assay reagent (BioRad). Equal amounts of cellular protein were separated by SDS-PAGE and electrotransferred to a nitrocellulose membrane. SHP protein was detected using anti-SHP antibody (SHP Q14, Santa Cruz Biotechnology) and HRP-conjugated anti-IgG antibody (sc-2020, Santa Cruz Biotechnology). Signal intensities were quantified using Quantity One software (BioRad). SHP protein intensities in each sample were normalized with respect to GAPDH protein, detected with anti-GAPDH antibody (V-18, Santa Cruz Biotechnology) and anti-IgG antibody.

2.6 siRNA experiments

HepG2 cells were transfected with either siRNA oligonucleotides targeting SHP (Ambion no. 103854) or with negative control siRNA (Ambion no. AM4636) using Ambion silencer siRNA transfection kit. Briefly, siRNAs were incubated with siPort NeoFx reagent (Ambion) to obtain transfection complexes. A

total of 10^5 cells were incubated with 0.03 nmol of siRNA and seeded in 12-well plates. Eight hours after transfection, transfection media were replaced with serum free media and cells were grown for additional 24 h before extraction of lipids, protein, and RNA. To assess interference efficiency, SHP mRNA was quantified by quantitative PCR.

2.7 Gene expression analysis

HepG2 total RNA was isolated using NucleoSpin RNA2 kit (Macherie-Naegel). Liver total RNA was obtained using Trizol reagent (Invitrogen). Reverse transcription reactions were performed on an ABI Prism 7300 SDS Real Time PCR system (Applied Biosystems) using the Taqman Reverse Transcription reagent kit (Applied Biosystems) and specific primers and Taqman probes (Applied Biosystems). Primer sequences are available upon request. Results were normalized to cyclophilin or RPLPO. For microarray analysis, Cy3- or Cy5-labelled cRNA was obtained from each RNA pool sample, hybridized against Agilent Mouse Oligo Microarrays, and data from the microarray images were obtained and analyzed with the Agilent Feature Extraction software. Two differential hybridizations (dye swap) were performed in each microarray experiment.

2.8 Statistical analysis

For statistical analysis in the lipid, gene expression, immunodetection studies, t-test, and ANOVA analyses were performed using SPSS software.

3. Results

3.1 GSPE decreases ApoB and TG synthesis and secretion in HepG2 cells

GSPE presented no cytotoxic effects in concentrations up to 150 mg/L on HepG2 cells, as measured by LDH leakage and Alamar Blue quantification. In this work GSPE was used in the range of 20-100 mg/L.

In order to assess whether HepG2 cells are a valid system to study the bioactivity of GSPE on TG and

ApoB secretion, we performed dose- and time-response experiments. As assessed by immunoblotting analysis, ApoB levels in the culture media, normalized by cell number, decreased dose-dependently 24 h after addition of GSPE, being 20 and 40% lower in cells treated with 50 and 100 mg/L GSPE, respectively, than in vehicle-treated cells. Treatment of cells with 50 mg/L GSPE, completely blocked the accumulation of ApoB in the media during the first 12 h of GSPE treatment, while during the following 12 h, this secretion was partially restored. In this same set of experiments, accumulation of TG in the media was drastically reduced upon treatment with 50 mg/L GSPE for 24 h, reaching 30% the value of vehicle-treated cells. In order to assess the action of GSPE on the de novo synthesis and secretion of TG, HepG2 cells were treated for 12 h with ¹⁴C-acetate and either vehicle or GSPE. Compared to controls, cells treated with GSPE showed a reduction in total ¹⁴C incorporation into TG, a significant decrease in secreted ¹⁴C-TG, without significant intracellular ¹⁴C-TG accumulation. Altogether, these results strongly suggest that GSPE lowers TG-rich lipoprotein secretion. Therefore, GSPE reduces TG and an ApoB secretion in HepG2 cells, which may be regarded as a valid system to study the mechanisms employed by procyanidins to exert these effects in vitro.

3.2 GSPE modulates SHP expression in HepG2 cells

In healthy normolipidemic rats, the hypotriglyceridemic effect of GSPE in vivo occurs concomitantly with an upregulation of SHP expression in the liver, 5 h after oral administration of GSPE [15]. To assess whether SHP expression is also modulated by GSPE in HepG2 cells we first treated HepG2 cells with different doses of GSPE and analyzed changes in SHP mRNA levels by quantitative PCR after 2 h of treatment. These GSPE treatments resulted in dose-dependent upregulation of SHP mRNA levels, reaching a two-fold increase versus control values with a dose of 100 mg/L. The increase in SHP mRNA level was followed by an increase in the level of SHP protein, as assessed by immunoblotting, in HepG2 cells treated during 3 h with a dose of 100 mg/L. Longer periods of GSPE treatment, assayed with a dose of 50 mg/L resulted in a reduction of SHP mRNA, which reached 50% of control values after 6 h of treatment. Therefore, GSPE modulates the expression of SHP in HepG2 cells, triggering a rapid and

transient induction of SHP expression.

3.3 Silencing SHP abolishes the effect of GSPE on TG secretion in HepG2 cells

To study the relevance of SHP as a mediator of the effects of GSPE on ApoB and TG secretion in HepG2, we inhibited SHP expression by transfecting the cells with SHP-specific siRNA. Transfection of HepG2 cells with SHP-siRNA resulted in a 60% reduction in SHP mRNA levels 32 h after transfection. Concomitantly with this reduction in SHP expression, ApoB, and TG release into the media increased 1.5- and 4-fold, respectively, strongly suggesting that SHP activity constrains ApoB and TG secretion in HepG2 cells in basal conditions. In SHP-silenced HepG2 cells, the TG-lowering effect of GSPE was completely abolished. In contrast, the effect of GSPE lowering ApoB secretion remained unaffected in SHP-knocked down HepG2 cells. These results show that the inhibition of ApoB and TG secretion by GSPE in HepG2 cells follow at least two different mechanisms: the action of GSPE on TG secretion involves a SHP-dependent mechanism, whereas repression of ApoB release follows a SHP-independent pathway.

3.4 SHP is a key mediator of the hypotriglyceridemic effect of GSPE in mice

In order to gain insight into the relevance of SHP as a mediator of GSPE hypotriglyceridemic actions in an in vivo system, we compared the effects of GSPE administration in wild-type versus SHP^{-/-} mice. Mice were feed two oral doses of GSPE (250 mg/kg), with a 12 h interval and plasma TG and total cholesterol were analyzed 2 h after the second oral gavage. In wild-type mice, GSPE gavage triggered a 40% reduction in plasma TG levels, a similar response to that previously found in rats [15]. This hypotriglyceridemic effect of GSPE was abolished by the SHP^{-/-} genotype. Therefore, SHP is a key mediator of the hypotriglyceridemic effects exerted by procyanidins in vivo. In contrast, GSPE gavage produced no effects on plasma total cholesterol levels, neither in wild-type nor in SHP-null mice.

3.5 GSPE modulates the hepatic expression of numerous genes related to lipid, fatty acid

(FA), and steroid metabolism in wild-type but not in SHP-/- mice

To gain insight into the SHP-dependent actions of GSPE that accompany the hypotriglyceridemic effect of GSPE in vivo, we compared the changes in gene expression induced by GSPE in the livers of wild-type and SHP-/- mice, using oligonucleotide microarray hybridization. In order to identify those genes whose expression in liver is modulated by GSPE in a SHP-dependent manner and are putatively involved in the hypotriglyceridemic effect of GSPE, we selected those genes clustered into the "lipid, FA, and steroid metabolism" group by Panther software [25], whose expression was changed by GSPE treatment in wild-type mice but remained unaltered in SHP-null mice, setting a fold-change threshold of 1.4 for upregulated and 0.7 for downregulated genes (Table 1). Several genes related to FA, TG, and cholesterol synthesis were identified which show SHP-dependent repression by GSPE, including key regulators of lipid synthesis pathways such as SREBP1C, 3-hydroxy-3-methylglutaryl-coenzyme A synthase, acyl-CoA synthetase, stearoyl-coenzyme A desaturase, and phosphatidic acid phosphatase 2a. Also the expression of several genes involved in FA oxidation and in lipoprotein metabolism was modulated by GSPE in a SHP-dependent manner, including carnitine palmitoyltransferase-1a (CPT-1A), apolipoprotein A5 (ApoA5), CCAAT/enhancer binding protein beta (C/EBP β), and phospholipid transfer protein (Pltp). Remarkably, numerous genes (marked with an asterisk in Table 1) which show a SHP-dependent response to GSPE, have been described as targets of SREBP1, a master regulator of lipid and lipoprotein metabolism [26, 27]. Therefore, SREBP1 emerges as a procyanidin target gene in the liver and as a putative SHP-dependent downstream effector of the hypotriglyceridemic response triggered by procyanidins in hepatic cells in vivo.

4. Discussion

Elevated TG levels are prevalent among the western population, often occurring in persons who are overweight or obese, or who have type 2 diabetes or the metabolic syndrome [5, 6, 8]. Compelling evidence from multiple clinical studies has established hypertriglyceridemia as an independent risk factor

for atherosclerosis [4-7]. Thus, using strategies to manage TG levels, along with low-density lipoprotein cholesterol levels, is warranted to help reduce the risk of CVD. Previously [15] we have shown that oral administration of GSPE triggered a 50% reduction in plasma ApoB and TG levels with a concomitant three-fold increase in liver SHP mRNA expression, in the postprandial state in healthy rats. In the liver, the orphan NR0B2/SHP regulates, among other processes, cholesterol and bile acid metabolism, steroidogenesis, and lipogenesis [16-18]. Therefore, SHP emerged as a putative mediator of the observed hypotriglyceridemic response triggered by GSPE.

In this work, we have shown that GSPE exerts hypotriglyceridemic activity when administered orally to mouse and reduces TG secretion in HepG2 cells, confirming the TG-lowering effect of GSPE previously observed in rats [15]. The similarity in the effects of GSPE in vitro and in vivo suggests that procyanidins species that exert the TG-lowering effect have a low polymerization degree, because only monomeric to pentameric procyanidins have been detected in plasma after oral consumption of procyanidin mixtures in murine models [28, 29]. The TG-lowering effect of GSPE is abolished in the in vitro and in vivo models when SHP expression is diminished or abolished: in HepG2 cells transfected with a SHP-specific siRNA and in SHP knockout mouse. Therefore, GSPE needs the activity of SHP to reduce TG secretion in vitro and triglyceridemia in vivo. In contrast, silencing of SHP expression in HepG2 cells did not abolish the ApoB-lowering effect of GSPE in this cells. Thus, repression of TG-rich ApoB-containing lipoproteins secretion by GSPE in hepatocytes involves both a SHP-dependent pathway, that affects TG secretion, and SHP-independent mechanisms, that affects ApoB release. Some flavonoids have been shown to decrease ApoB release, a marker of lipoprotein secretion, in in vitro hepatocyte models. Although the intracellular mediators involved in this effect are not fully elucidated, the inhibition of P13-K and MAPK/ERK signaling pathways and repression of microsomal TG transfer protein (MTP) activity have been demonstrated [12-14, 30-32].

Comparison of the gene expression changes elicited by GSPE in the liver of wild-type mouse to those elicited in the SHP-null mouse has allowed the identification of SHP-dependent procyanidin-target genes

putatively implicated in the hypotriglyceridemic effect of GSPE. Among these genes, it stands out SREBP-1c, that is downregulated by GSPE in the liver of wild-type mouse, and ApoA5 and CPT-1A, which are upregulated by GSPE.

SREBP-1c is a master regulator of FA synthesis and a mediator for insulin/glucose signaling to lipogenesis, and could be involved in the development of insulin resistance and fatty livers, being its inhibition a proposed target for lowering lipogenesis and TG levels [33-35]. Transgenic mouse overexpressing SREBP1c displays increased expression of FA synthase (FAS), and higher levels of hepatic FAs, and TG [36]; in contrast, SREBP-1c knockout mouse displays diminished FAS expression, lower hepatic FA amounts, and lower plasma TG levels [37]. It is well established that bile acids and synthetic FXR ligands downregulate the hepatic secretion of TG by a pathway that involves induction of SHP and subsequent inhibition of SREBP-1c expression by SHP [19]. In a similar way, SREBP-1c could be a mediator of the hypotriglyceridemic effect of GSPE acting downstream of SHP. Among the genes that we have found to be downregulated by GSPE in mouse liver in a SHP-dependent manner there are several which are known or putative targets of SREBP1c. These include genes involved in the synthesis of monounsaturated and PUFAs: a long chain FA elongase (Elov13) [38, 39], two acyl-CoA synthetases (Acsm2 and Acss2) [26, 33, 35], and a FA desaturase, the stearoyl-coenzyme A desaturase 2 (Scd2) [40, 41]. Scd2 is a microsomal enzyme required for the biosynthesis of oleate and palmitoleate, which are the major MUFAs of membrane phospholipids, TG, and cholesterol esters. Deficiency in FA desaturase activity greatly reduces hepatic TG synthesis [42] and protects mouse against hypertriglyceridemia induced by liver X receptor (LXR) activation [43]. Other genes which we have found to be downregulated by GSPE in a SHP dependent manner, although not known targets of SREBP-1c, include genes encoding enzymes directly involved in TG synthesis such as the phosphatidic acid phosphatase 2. Altogether, this pattern of gene expression induced by GSPE in mouse liver, suggest a diminished rate of TG and FA synthesis.

Along with the downregulation of FA and TG synthetic genes, GSPE upregulated the hepatic expression

of CPT-1A, the rate limiting enzyme in the beta-oxidation of FA. Enhancement of CPT-1 expression and activation of the FA oxidation pathway are potential targets for the treatment of hypertriglyceridemia, type-2 diabetes, and obesity [44, 45]. Our results suggest that SHP activity is necessary for the GSPE-induced upregulation of CPT-1A expression in mouse liver, establishing CPT-1A as a putative target of SHP activity.

Besides the effects of GSPE in reducing TG secretion by the liver, increasing plasma TG catabolism could also contribute to the hypotriglyceridemic effect of GSPE. In this regard, we have found that GSPE upregulates ApoA5 expression in mouse liver, in a SHP-dependent manner. Overexpression of ApoA5 is known to reduce plasma TG and cholesterol in hypertriglyceridemic mouse [46, 47] whereas ApoA5 deficiency results in marked hypertriglyceridemia [48]. The hypotriglyceridemic action of ApoA5 is attributed to the activation of lipase-mediated VLDL-TG hydrolysis and consequent acceleration of VLDL catabolism [48-50]. Again, this result suggests that ApoA5 expression may be under the control of SHP activity. It is known that ApoA5 gene expression is regulated by LXR ligands in a negative manner through upregulation of SREBP-1c [51].

In summary, we have shown here that grape procyanidins exert hypotriglyceridemic effects when orally administered to mouse, and when applied to HepG2 cells. In both models, procyanidins require the presence of SHP to reduce TG-levels. We have demonstrated the SHP-dependent hepatic downregulation of expression of SREBP-1c and upregulation of CPT-1 and ApoA5 in the liver of GSPE-treated mouse, strongly suggesting that procyanidins repress hepatic FA synthesis, enhance hepatic FA oxidation, and activate the catabolism of TG-rich ApoB-containing lipoproteins, by a mechanism that depends on the activity of SHP. In addition, we have shown in HepG2 cells that procyanidins downregulate the secretion of ApoB in a SHP-independent way. Altogether, these results show that procyanidins, using SHP-dependent and SHP-independent mechanisms, may lower the secretion of TG-rich ApoB-containing lipoproteins by hepatic cells. Both mechanisms would act in concert to reduce the number of circulating VLDLs and also their TG content providing a plausible explanation for the hypolipidemic properties of

grape polyphenols present in foods, which lower TG and ApoB levels in human subjects [52, 53] when administered chronically at lower doses than those used in this study. Due to the relevance of postprandial triglyceridemia in the development of atherosclerosis and type 2 diabetes [7, 8], the elucidation of how nutrients can modulate plasma lipid levels has emerged as an important target of nutrition research [54]. In this regard, this work provides new hints for understanding the mechanisms associated with the beneficial effects ascribed to the regular consumption of dietary procyanidins.

Acknowledgments

This study was supported by grant number AGL2005-04889 from the Spanish Dirección General de Investigación del Ministerio de Educación y Ciencia. J. M. del Bas is the recipient of a fellowship from the Spanish Government. We gratefully acknowledge the expert technical assistance of the Centro Nacional de Investigaciones Oncológicas (Madrid, Spain) in performing microarray hybridizations and data acquisition. The authors have declared no conflict of interest.

References

- [1] Crozier, A., Burns, J., Aziz, A. A., Stewart, A. J., et al., Antioxidant flavonols from fruits, vegetables and beverages: Measurements and bioavailability. *Biol. Res.* 2000, 33, 79-88.
- [2] Ariga, T., The antioxidative function, preventive action on disease and utilization of proanthocyanidins. *Biofactors* 2004, 21, 197-201.
- [3] Rasmussen, S. E., Frederiksen, H., Krogholm, K. S., Poulsen, L., Dietary proanthocyanidins: Occurrence, dietary intake, bioavailability, and protection against cardiovascular disease. *Mol. Nutr. Food Res.* 2005, 49, 159-174.
- [4] Bersot, T., Haffner, S., Harris, W. S., Kellick, K. A., Morris, C. M., Hypertriglyceridemia: Management of atherogenic dyslipidemia. *J. Fam. Pract.* 2006, 55, S1-S8.

308 [5] Le, N. A., Walter, M. F., The role of hypertriglyceridemia in atherosclerosis. *Curr. Atheroscler. Rep.*
309 2007, 9, 110-115.

310 [6] Jacobson, T. A., Miller, M., Schaefer, E. J., Hypertriglyceridemia and cardiovascular risk reduction.
311 *Clin. Ther.* 2007, 29,763-777.

312 [7] Hyson, D., Rutledge, J. C., Berglund, L., Postprandial lipemia and cardiovascular disease. *Curr.*
313 *Atheroscler. Rep.* 2003, 5, 437-444.

314 [8] Grundy, S. M., Brewer, H. B., Jr., Cleeman, J. I., Smith, S. C., Jr., Lenfant, C., Definition of metabolic
315 syndrome: Report of the National Heart, Lung, and Blood Institute/American Heart Association
316 conference on scientific issues related to definition. *Circulation* 2004, 109,433-438.

317 [9] Vinson, J. A., Teufel, K., Wu, N., Red wine, dealcoholized red wine, and especially grape juice,
318 inhibit atherosclerosis in a hamster model. *Atherosclerosis* 2001, 156, 67-72.

319 [10] Auger, C., Caporiccio, B., Landrault, N., Teissedre, P. L., et al., Red wine phenolic compounds
320 reduce plasma lipids and apolipoprotein B and prevent early aortic atherosclerosis in
321 hypercholesterolemic golden Syrian hamsters (*Mesocricetus auratus*). *J. Nutr.* 2002, 132, 1207-1213.

322 [11] Pal, S., Ho, N., Santos, C., Dubois, P., et al., Red wine polyphenolics increase LDL receptor
323 expression and activity and suppress the secretion of ApoB100 from human HepG2 cells. *J. Nutr.* 2003,
324 133, 700-706.

325 [12] Davalos, A., Fernandez-Hernando, C., Cerrato, F., Martinez-Botas, J., et al., Red grape juice
326 polyphenols alter cholesterol homeostasis and increase LDL-receptor activity in human cells in vitro. *J.*
327 *Nutr.* 2006, 136, 1766-1773.

328 [13] Allister, E. M., Borradaile, N. M., Edwards, J. Y., Huff, M. W., Inhibition of microsomal triglyceride
329 transfer protein expression and apolipoprotein B100 secretion by the citrus flavonoid naringenin and by

330 insulin involves activation of the mitogen-activated protein kinase pathway in hepatocytes. *Diabetes*
331 2005, 54, 1676-1683.

332 [14] Borradaile, N. M., de Dreu, L. E., Huff, M. W., Inhibition of net HepG2 cell apolipoprotein B
333 secretion by the citrus flavonoid naringenin involves activation of phosphatidylinositol 3-kinase,
334 independent of insulin receptor substrate-1 phosphorylation. *Diabetes* 2003, 52, 2554-2561.

335 [15] Del Bas, J. M., Fernandez-Larrea, J., Blay, M., Ardevol, A., et al., Grape seed procyanidins improve
336 atherosclerotic risk index and induce liver CYP7A1 and SHP expression in healthy rats. *FASEB J.* 2005,
337 19, 479-481.

338 [16] Bavner, A., Sanyal, S., Gustafsson, J. A., Treuter, E., Transcriptional corepression by SHP:
339 Molecular mechanisms and physiological consequences. *Trends Endocrinol. Metab.* 2005, 16, 478-488.

340 [17] Boulias, K., Katrakili, N., Bamberg, K., Underhill, P., et al., Regulation of hepatic metabolic
341 pathways by the orphan nuclear receptor SHP. *EMBO J.* 2005, 24, 2624-2633.

342 [18] Lee, Y. S., Chanda, D., Sim, J., Park, Y. Y., Choi, H. S., Structure and function of the atypical
343 orphan nuclear receptor small heterodimer partner. *Int. Rev. Cytol.* 2007, 261, 117-158.

344 [19] Watanabe, M., Houten, S. M., Wang, L., Moschetta, A., et al., Bile acids lower triglyceride levels via
345 a pathway involving FXR, SHP, and SREBP-1c. *J. Clin. Invest.* 2004, 113, 1408-1418.

346 [20] Urizar, N. L., Moore, D. D., GUGULIPID: A natural cholesterol-lowering agent. *Ann. Rev. Nutr.*
347 2003, 23, 303-313.

348 [21] Puiggros, F., Llopiz, N., Ardevol, A., Blade, C., et al., Grape seed procyanidins prevent oxidative
349 injury by modulating the expression of antioxidant enzyme systems. *J. Agric. Food Chem.* 2005, 53,
350 6080-6086.

351 [22] Wang, L., Lee, Y. K., Bundman, D., Han, Y., et al., Redundant pathways for negative feedback
352 regulation of bile acid production. *Dev. Cell* 2002, 2, 721-731.

353 [23] Urizar, N. L., Liverman, A. B., Dodds, D. T., Silva, F. V., et al., A natural product that lowers
354 cholesterol as an antagonist ligand for FXR. *Science* 2002, 296, 1703-1706.

355 [24] Pill, J., Aufenanger, J., Stegmeier, K., Schmidt, F. H., Müller, D., Thin-layer chromatography of
356 radioactively labelled cholesterol and precursors from biological material. *Fresenius J. Anal. Chem.*
357 1987,327,558-560.

358 [25] Thomas, P. D., Campbell, M. J., Kejariwal, A., Mi, H., et al., PANTHER: A library of protein
359 families and subfamilies indexed by function. *Genome Res.* 2003, 13, 2129-2141.

360 [26] Horton, J. D., Shah, N. A., Warrington, J. A., Anderson, N. N., et al., Combined analysis of
361 oligonucleotide microarray data from transgenic and knockout mice identifies direct SREBP target genes.
362 *Proc. Natl. Acad. Sci. USA* 2003, 100, 12027-12032.

363 [27] Maxwell, K. N., Soccio, R. E., Duncan, E. M., Sehayek, E., Breslow, J. L., Novel putative SREBP
364 and LXR target genes identified by microarray analysis in liver of cholesterol-fed mice. *J. Lipid Res.*
365 2003, 44, 2109-2119.

366 [28] Garcia-Ramirez, B., Fernandez-Larrea, J., Salvado, M. J., Ardevol, A., et al., Tetramethylated
367 dimeric procyanidins are detected in rat plasma and liver early after oral administration of synthetic
368 oligomeric procyanidins. *J. Agric. Food Chem.* 2006, 54, 2543-2551.

369 [29] Shoji, T., Masumoto, S., Moriichi, N., Akiyama, H., et al., Apple procyanidin oligomers absorption
370 in rats after oral administration: Analysis of procyanidins in plasma using the porter method and high-
371 performance liquid chromatography/tandem mass spectrometry. *J. Agric. Food Chem.* 2006, 54, 884-892.

372 [30] Wilcox, L. J., Borradaile, N. M., de Dreu, L. E., Huff, M. W., Secretion of hepatocyte apoB is

373 inhibited by the flavonoids naringenin and hesperetin, via reduced activity and expression of ACAT2 and
374 MTP. *J. Lipid Res.* 2001, 42, 725-734.

375 [31] Kurowska, E. M., Manthey, J. A., Regulation of lipoprotein metabolism in HepG2 cells by citrus
376 flavonoids. *Adv. Exp. Med. Biol.* 2002, 505, 173-179.

377 [32] Yee, W. L., Wang, Q., Agdinaoy, T., Dang, K., et al., Green tea catechins decrease apolipoprotein
378 B-100 secretion from HepG2 cells. *Mol. Cell. Biochem.* 2002, 229, 85-92.

379 [33] Shimano, H., Sterol regulatory element-binding proteins (SREBPs): Transcriptional regulators of
380 lipid synthetic genes. *Prog. Lipid Res.* 2001, 40, 439-452.

381 [34] Shimano, H., Sterol regulatory element-binding protein family as global regulators of lipid synthetic
382 genes in energy metabolism. *Vitam. Horm.* 2002, 65, 167-194.

383 [35] Horton, J. D., Goldstein, J. L., Brown, M. S., SREBPs: Activators of the complete program of
384 cholesterol and fatty acid synthesis in the liver. *J. Clin. Invest.* 2002, 109, 1125-1131.

385 [36] Shimano, H., Horton, J. D., Shimomura, I., Hammer, R. E., et al., Isoform 1c of sterol regulatory
386 element binding protein is less active than isoform 1a in livers of transgenic mice and in cultured cells. *J.*
387 *Clin. Invest.* 1997, 99, 846-854.

388 [37] Liang, G., Yang, J., Horton, J. D., Hammer, R. E., et al., Diminished hepatic response to
389 fasting/refeeding and liver X receptor agonists in mice with selective deficiency of sterol regulatory
390 element-binding protein-1c. *J. Biol. Chem.* 2002, 277, 9520-9528.

391 [38] Matsuzaka, T., Shimano, H., Yahagi, N., Yoshikawa, T., et al., Cloning and characterization of a
392 mammalian fatty acyl-CoA elongase as a lipogenic enzyme regulated by SREBPs. *J. Lipid Res.* 2002,
393 43, 911-920.

394 [39] Wang, Y., Botolin, D., Xu, J., Christian, B., et al., Regulation of hepatic fatty acid elongase and
395 desaturase expression in diabetes and obesity. *J. Lipid Res.* 2006, 47, 2028-2041.

396 [40] Tabor, D. E., Kim, J. B., Spiegelman, B. M., Edwards, P. A., Transcriptional activation of the
397 stearoyl-CoA desaturase 2 gene by sterol regulatory element-binding protein/adipocyte determination and
398 differentiation factor 1. *J. Biol. Chem.* 1998, 273, 22052-22058.

399 [41] Bene, H., Lasky, D., Ntambi, J. M., Cloning and characterization of the human stearoyl-CoA
400 desaturase gene promoter: Transcriptional activation by sterol regulatory element binding protein and
401 repression by polyunsaturated fatty acids and cholesterol. *Biochem. Biophys. Res. Commun.* 2001, 284,
402 1194-1198.

403 [42] Attie, A. D., Krauss, R. M., Gray-Keller, M. P., Brownlie, A. et al., Relationship between stearoyl-
404 CoA desaturase activity and plasma triglycerides in human and mouse hypertriglyceridemia. *J. Lipid Res.*
405 2002, 43, 1899-1907.

406 [43] Chu, K., Miyazaki, M., Man, W. C., Ntambi, J. M., Stearoyl-coenzyme A desaturase 1 deficiency
407 protects against hypertriglyceridemia and increases plasma high-density lipoprotein cholesterol induced
408 by liver X receptor activation. *Mol. Cell. Biol.* 2006, 26, 6786-6798.

409 [44] Chakrabarti, R., Vikramadithyan, R. K., Misra, P., Hiriyan, J., et al., Ragaglitazar: A novel PPAR
410 alpha PPAR gamma agonist with potent lipid-lowering and insulin-sensitizing efficacy in animal models.
411 *Br. J. Pharmacol.* 2003, 140, 527-537.

412 [45] Kuhajda, F. P., Ronnett, G. V., Modulation of carnitine palmitoyltransferase-1 for the treatment of
413 obesity. *Curr. Opin. Investig. Drugs* 2007, 8, 312-317.

414 [46] van der Vliet, H. N., Schaap, F. G., Levels, J. H., Ottenhoff, R., et al., Adenoviral overexpression of
415 apolipoprotein A-V reduces serum levels of triglycerides and cholesterol in mice. *Biochem. Biophys. Res.*

416 Commun. 2002, 295, 1156-1159.

417 [47] Huang, W., Bi, N., Zhang, X., Wang, Y., et al., Overexpression of apolipoprotein AV in the liver
418 reduces plasma triglyceride and cholesterol but not HDL in ApoE deficient mice. Biochem. Biophys. Res.
419 Commun. 2006, 346, 14-18.

420 [48] Grosskopf, I., Baroukh, N., Lee, S. J., Kamari, Y., et al., Apolipoprotein A-V deficiency results in
421 marked hypertriglyceridemia attributable to decreased lipolysis of triglyceride-rich lipoproteins and
422 removal of their remnants. Arterioscler. Thromb. Vasc. Biol. 2005, 25, 2573-2579.

423 [49] Schaap, F. G., Rensen, P. C., Voshol, P. J., Vrans, C., et al., ApoAV reduces plasma triglycerides by
424 inhibiting very low density lipoprotein-triglyceride (VLDL-TG) production and stimulating lipoprotein
425 lipase-mediated VLDL-TG hydrolysis. J. Biol. Chem. 2004, 279, 27941-27947.

426 [50] Merkel, M., Loeffler, B., Kluger, M., Fabig, N., et al., Apolipoprotein AV accelerates plasma
427 hydrolysis of triglyceride-rich lipoproteins by interaction with proteoglycan-bound lipoprotein lipase. J.
428 Biol. Chem. 2005, 280, 21553-21560.

429 [51] Jakel, H., Nowak, M., Moitrot, E., Dehondt, H., et al., The liver X receptor ligand T0901317 down-
430 regulates APOA5 gene expression through activation of SREBP-Ic. J. Biol. Chem. 2004, 279, 45462-
431 45469.

432 [52] Castilla, P., Echarri, R., Davalos, A., Cerrato, F., et al., Concentrated red grape juice exerts
433 antioxidant, hypolipidemic, and antiinflammatory effects in both hemodialysis patients and healthy
434 subjects. Am. J. Clin. Nutr. 2006, 84, 252-262.

435 [53] Zern, T. L., Wood, R. J., Greene, C., West, K. L., et al., Grape polyphenols exert a cardioprotective
436 effect in pre and postmenopausal women by lowering plasma lipids and reducing oxidative stress. J. Nutr.
437 2005, 135, 1911-1917.

[54] Muller, M., Kersten, S., Nutrigenomics: Goals and strategies. Nat. Rev. Genet. 2003, 4,315-322.

Figure Legends

Figure 1. Effect of GSPE on ApoB release in HepG2 cells.

(A) Changes in media ApoB levels in response to 20, 50, and 100 mg/L GSPE doses. HepG2 cells were treated with either vehicle (line 0) or GSPE at the indicated concentrations. After 24 h, ApoB in the media was quantified by immunoblotting as described in Section 2. Relative intensities of the ApoB bands were normalized by cell number. The immunoblot of one from three independent experiments is shown.

(B) Time-response dependent changes in media ApoB accumulation induced by GSPE. HepG2 cells were treated with 50 mg/L GSPE or vehicle (control) and media was collected after 6, 12, and 24 h. ApoB levels were analyzed as in (A). All values are the mean $\pm$ SEM of three independent experiments; * on a GSPE-treated sample denotes a significant difference ($p < 0.05$) between the value of that sample and its control counterpart.

Figure 2. Effect of GSPE on TG synthesis and secretion in HepG2 cells.

(A) Changes in TG accumulation in media of cells treated with GSPE. HepG2 cells were treated with 50 mg/L of GSPE or vehicle (control) for 24 h. TG were assayed as indicated in Section 2 and values were normalized to mg of cell protein.

(B) Changes induced by GSPE in the de novo synthesis and secretion of TG. HepG2 cells were simultaneously incubated with ^{14}C -labelled acetate and 50 mg/L of GSPE or vehicle (control). After 12 h of treatment, radioactivity incorporated into media and cellular TG was measured as described in Section 2. All values are the mean $\pm$ SEM of three independent experiments; * on a GSPE-treated sample denotes a significant difference ($p < 0.05$) between the value of that sample and its control counterpart.

Figure 3. Effects of GSPE on SHP expression in HepG2 cells.

(A) Fold-change of SHP gene expression in response to 20, 50, and 100 mg/L GSPE relative to vehicle treated cells. HepG2 cells were treated with the indicated concentrations of GSPE for 2 h. SHP mRNA levels were determined by RT-qPCR and normalized to RPLP0 mRNA levels.

(B) Changes in SHP protein levels in response to GSPE treatment monitored by Western blot. HepG2 cells were treated with either vehicle (control) or 100 mg/L GSPE. After 3 h, cell lysates were obtained and SHP levels were analyzed by immunoblotting, using GAPDH as the loading control.

(C) Time course of SHP gene expression in response to treatment with 50 mg/L of GSPE. HepG2 total RNA was obtained at the indicated times and analyzed as in (A). All values are the mean $\pm$ SEM of three independent experiments; * on a GSPE-treated sample denotes a significant difference ($p < 0.05$) between the value of that sample and its control counterpart.

Figure 4. Effect of SHP silencing on GSPE modulation of ApoB and TG secretion in HepG2 cells.

HepG2 cells were transfected with a SHP-specific siRNA (+ bars) or negative control, scramble siRNA (- bars). Thirty-two hours after transfection, media was replaced with serum depleted medium, and 12 h afterwards, cells were treated with 50 mg/L GSPE (indicated by +) or vehicle (indicated by -). After GSPE treatment, cells and media were harvested for SHP, ApoB, and TG analysis.

(A) SHP gene expression, determined as in Fig. 3A, after 2 h of GSPE treatment.

(B) ApoB in the cell culture media after 24 h of GSPE treatment, analyzed as in Fig. 1A. A representative ApoB immunoblot is shown.

(C) TG levels in culture media after 24 h of GSPE treatment, analyzed as in Fig. 2A. All values are the mean $\pm$ SEM of three independent experiments. a, b, or c denotes significant differences between groups, as determined by ANOVA; * on a GSPE-treated sample denotes a significant difference ($p < 0.05$) between the value of that sample and its control counterpart.

Figure 5. Effect of GSPE on plasma TG and total cholesterol levels in wild-type and SHP-null mouse.

Wild-type (SHP^{+/+}) and SHP-null (SHP^{-/-}) mice were fed with either vehicle (-bars) or GSPE (250 mg/Kg) (+ bars) via oral gavage ($n = 5$ in each group, age 8-10 wks) as described in Section 2. Plasma was obtained 14 h after the first oral gavage (2 h after the second oral gavage) and TG and total cholesterol were determined as described in Section 2. All values are the mean $\pm$ SEM of three independent experiments; * on a GSPE-treated sample denotes a significant difference ($p < 0.05$) between

489 the value of that sample and its control counterpart.

490 **Tables**

491 **Table 1. SHP-dependent changes induced by GSPE in the expression of lipid-related genes in mouse**
492 **liver**

Name / Gene symbol	Genbank ID	WT	SHP-/-
FA synthesis			
Acyl-CoA synthetase medium-chain family member 2; Acsm2	NM_146197	0.6	1
Acyl-CoA synthetase short-chain family member 2; Acss2*	NM_019811	0.6	0.8
Stearoyl-coenzyme A desaturase 2; Scd2*	NM_009128	0.7	0.9
Sterol regulatory element binding factor 1; Srebp1c*	NM_011480	0.7	1.0
Cytochrome P450, family 2, subfamily c, polypeptide 55;	NM_028089	0.5 ± 0.1	1.0 ± 0.2

Cyp2c55			
Elongation of very long chain FAs (FEN1/Elo2, SUR4/Elo3, yeast)-like 3; Elovi3*	NM_007703	0.7	1
Glycerophospholipid, diacylglycerol, and triacylglycerol synthesis			
Phosphatidic acid phosphatase 2a; Ppap2a	NM_008903	0.7	0.9
DNA segment, Chr 3, Brigham and amp; D3Bwg0562e	NM_177664	0.5	1
Phosphatidylinositol-4-phosphate 5-kinase, type II, a; Pip5k2a	NM_008845	0.7	1
Ethanolamine kinase 2; Etnk2	NM_175443	0.7	0.9

Choline kinase a: Chka	NM_013490	0.7	0.9
Steroid and cholesterol synthesis			
Mevalonate kinase; Mvk	NM_023556	0.7	1.2
Sterol-C5-desaturase; Sc5d*	NM_172769	0.7	1
3-Hydroxy-3- methylglutaryl- Coenzyme A synthase 1; Hmgcs1*	NM_145942	0.6	1.2
NAD(P)-dependent steroid dehydrogenase-like; Nsdhl	NM_010941	0.7	1
7-Dehydrocholesterol reductase; Dhcr7*	NM_007856	0.7	1
Farnesyl diphosphate synthetase; Fdps	NM_134469	0.8	1.6
Phosphomevalonate	NM_026784	0.7	0.9



kinase; Pmvk			
Cytochrome b5 reductase 1; Cyb5r1	NM_028057	0.7	1.3
FA oxidation			
Carnitine palmitoyltransferase 1a, liver; Cpt1a	NM_013495	1.4	1.0
Other lipid-related genes			
Phospholipid transfer protein; Pltp	NM_011125	1.6 ± 0.2	0.9 ± 0.2
Apolipoprotein A5; ApoA5*	NM_080434	2.1 ± 0.2	1.0 ± 0.2
Phosphatidylinositol 3-kinase, C2 domain containing, g polypeptide; Pik3c2g	NM_207683	0.7	1
ST3 b-galactoside alpha-2,3- sialyltransferase 6; St3gal6	NM_018784	0.7	0.8

CCAAT/enhancer binding protein (C/EBP), b; Cebpb*	NM_009883	0.7	0.8
---	-----------	-----	-----

Wild-type and SHP-/- mice were fed either vehicle or GSPE (250 mg/Kg) via oral gavage (five animals in each treatment group) as described in Section 2. Total RNA was obtained from livers and pooled for each treatment group. Two different microarray hybridizations (dye swap) were performed to assess changes induced by GSPE in WT or SHP-/- mice. The values for each gene are the mean fold change obtained from two microarray hybridization experiments (dye swap labeling) comparing signal intensities obtained from GSPE-treated versus control groups. The table lists those genes which are involved in "lipid, FA, and steroid metabolism" following the classification of Panther software [25], and whose expression is changed by GSPE treatment in wild-type mice (mean fold changes in column "WT") but not in SHP-null mice (mean fold changes in column "SHP-/-"). Bold characters denote the mean fold change values obtained by RT-qPCR using individual RNAs instead of pooled RNAs for quantification. Known SREBP-1c target genes are denoted by asterisk.


Figures

Figure 1



506

507 Figure 2



508

