

**Circulating metabolites associated with objectively-measured sleep duration and sleep variability in overweight/obese subjects: a metabolomics approach within the SATIN study**

Christopher Papandreou<sup>1,2</sup>, Lucia Camacho-Barcia<sup>1,2</sup>, Jesús García-Gavilán<sup>1,2</sup>, Thea Toft Hansen<sup>3</sup>, Mads F Hjorth<sup>3</sup>, Jason C G Halford<sup>4</sup>, Jordi Salas-Salvadó<sup>1,2</sup>, Anders Sjödin<sup>3</sup>, Mónica Bulló<sup>1,2</sup>.

<sup>1</sup>  
Human Nutrition Unit, Faculty of Medicine and Health Sciences, Institut d'Investigació Sanitària Pere Virgili, Rovira i Virgili University, Reus, Spain

<sup>2</sup>  
CIBER Fisiopatología de la Obesidad y Nutrición (CIBEROBN), Instituto de Salud Carlos III, Madrid, Spain

<sup>3</sup>  
Department of Nutrition, Exercise and Sports, University of Copenhagen, Rolighedsvej 26, 1958 Frederiksberg C, Denmark

<sup>4</sup>  
Department of Psychological Sciences, Institute of Psychology Health and Society, University of Liverpool, Liverpool, UK

**Corresponding author:** Mònica Bulló, BSc, PhD, Human Nutrition Unit, Faculty of Medicine and Health Sciences, Universitat Rovira i Virgili, St/SantLlorenç 21, 43201 R(E-mail: [monica.bullo@urv.cat](mailto:monica.bullo@urv.cat)).

## Abstract

**Objectives:** To investigate the associations of circulating metabolites with sleep duration and sleep variability. We also assessed the ability of metabolites to discriminate between sleep duration and sleep variability categories.

**Methods:** Cross-sectional analysis of 205 participants with overweight/obesity from the “Satiety Innovation” (SATIN) study. A targeted metabolite profiling (n = 159 metabolites) approach was applied using three different platforms (NMR, LC-MS, GC-MS). Associations between circulating metabolite concentrations and accelerometer-derived sleep duration and variability in sleep duration were assessed using elastic-net regression analysis. Ten-fold cross-validation was used to estimate the discriminative accuracy of metabolites for sleep duration, and sleep variability categories.

**Results:** A metabolite profile, including acyl-carnitines (C11:0/C5:1-DC/iso-C11:0, 2-M-C4:1/3-M-C4:1, C4:0); sphingomyelins (42:1, 33:1); glycerol; stearic acid; 2- and 3- hydroxyl-butyric acid; docosahexaenoic acid; serotonin; and phosphatidylcholine (34:2), was significantly associated with high sleep duration (4<sup>th</sup> plus 5<sup>th</sup> quintile). Ten metabolites, including acyl-carnitines (C18:1, C7:0, C6-OH); phosphatidylcholine (40:6, 37:4, 42:5); lyso-phosphatidylcholine, (20:1); sucrose; glutamic acid, and triacylglycerol (52:4), were significantly associated with high sleep variability (4<sup>th</sup> plus 5<sup>th</sup> quintile). The area under the curve was 0.69 (95% CI 0.62–0.75), and 0.63 (95% CI 0.53–0.72), in the multi-metabolite score for high sleep duration, and sleep variability, respectively. The variance in sleep duration explained by metabolites was 7%. No metabolites were selected for prediction of sleep variability (continuous).

**Conclusion:** A small set of metabolites within distinct biochemical pathways were associated with high sleep duration and sleep variability. These metabolites appeared to moderately discriminate sleep duration and sleep variability categories.

**Keywords:** metabolomics, sleep duration, sleep variability, SATIN

### **Statement of significance**

Sleep restriction and irregular sleep-patterns have been associated with metabolic impairments. However, the metabolic dysregulations associated with these sleep disturbances remain to be determined under “real life”, free-living conditions. The present study provides data showing associations between circulating metabolites and objectively-measured sleep duration and sleep variability in overweight/obese adults. These findings may provide a deeper understanding of metabolic responses to sleep. Future research should assess the involvement of the identified metabolites in sleep-related metabolic health.

### **Introduction**

In humans, adequate sleep and a steady sleep regimen confer a wide range of health benefits, while both sleep deprivation and irregular sleep-patterns have been associated with metabolic disorders, including obesity and type 2 diabetes<sup>1-6</sup>. The biological mechanisms linking sleep deprivation/disruption and long-term health consequences have not yet been fully elucidated<sup>7</sup>.

Metabolic impairments associated with sleep loss have been observed in several experimental studies. Total (24h) sleep deprivation results in increased plasma levels

of glycerophospholipids, acylcarnitines, sphingolipids and amino acids (tryptophan, serotonin, taurine)<sup>8</sup>. Partial<sup>9,10</sup> sleep deprivation has also been shown to induce plasma and serum metabolite changes in various pathways. However, the effects of sleep deprivation under controlled laboratory conditions on human metabolism may not reflect the natural variation in sleep patterns observed in “real-life” conditions. In addition, previous studies suggest that irregular sleep-wake rhythms may be associated with metabolic dysfunction<sup>11,12</sup>. Recently, a cross-sectional study found 64 plasma metabolites across multiple biochemical pathways significantly associated with sleep timing (midpoint between bedtime and wake up time)<sup>13</sup>.

Studies combining objectively measured, “real-life” levels of free-living sleep-deprivation and high sleep-variability with systemic metabolite profiling, would provide a deeper understanding of metabolic responses to sleep and also help to identify relevant metabolites able to predict sleep-patterns.

Using targeted metabolite profiling, cross-sectional associations between plasma/serum concentrations of metabolites with sleep duration and sleep variability were examined in participants with overweight/obesity included in the EU project Satiety Innovation (SATIN) work package 5, prior to the implementation of a dietary intervention. The ability of metabolite markers to discriminate between sleep duration and sleep variability categories was also investigated.

## **Subjects and Methods**

### **Study Design**

A cross-sectional analysis of baseline data from the SATIN study, a medium-term clinical trial focusing on satiation / satiety enhancement, appetite control and weight management in overweight and obese participants was performed. All procedures

were conducted in accordance with the ethical principles set forth in the current version of the Declaration of Helsinki (Fortaleza, Brazil, October 2013) and the International Conference on Harmonization E6 Good Clinical Practice (ICH-GCP). The protocol was approved by the local institutional review boards and Ethics Committees of all the recruiting centres and all participants provided written informed consent. This trial was registered in: [clinicaltrials.gov](https://clinicaltrials.gov), with the identifier: NCT02485743.

### **Study population**

Eligible participants were community-dwelling adults (aged between 20 and 65 years), with an initial body mass index of 27.0 to 35.0 kg/m<sup>2</sup>, fat mass  $\geq 23\%$  and without comorbidities. Exclusion criteria included: weight changes  $\pm 3$ kg in the last three months, cardiometabolic diseases (type 1 or 2 diabetes, cardiovascular disease, hypertension). Cardiovascular disease included previous diagnosis of angina pectoris; myocardial infarction; coronary revascularization procedures; stroke (either ischemic or hemorrhagic, including transient ischemic attacks); symptomatic peripheral artery disease that required surgery or was diagnosed with vascular imaging techniques; ventricular arrhythmia; uncontrolled atrial fibrillation; congestive heart failure (New York Heart Association Class II or IV); hypertrophic cardiomyopathy; and history of aortic aneurism  $\geq 5.5$  cm in diameter or aortic aneurism surgery within the past six months, as diagnosed by a medical doctor. Other exclusion criteria were: chronic kidney and liver pathologies, diseases which may affect energy expenditure, current or former ( $< 3$ months) smoking and regular consumption of alcohol above the recommendations. A total of 236 participants from Reus (n=122) and Copenhagen (n=114) recruiting centers with blood samples available at baseline were considered. Thirty one participants were excluded due to the lack of actigraphy data.

### **Targeted metabolomics**

The current study used three different platforms to analyse metabolites. Proton nuclear magnetic resonance ( $^1\text{H}$  NMR), liquid chromatography coupled to mass spectrometry (LC-MS) and gas chromatography coupled to mass spectrometry (GC-MS) were used to perform optimal profiling of a wide range of metabolites. Fasting blood samples for GC-MS analysis were dried and stored at  $-80^\circ\text{C}$  until analysis, whereas supernatants of samples for LC-MS and NMR were directly analysed, maintained at  $4^\circ\text{C}$  during their sequence run.”

### **Automation of multiple plasma sample extraction**

Aqueous extractions of  $250\mu\text{L}$  of plasma were performed with a methanol/water solution in a Bravo automated liquid handling platform (Agilent Technologies, CA). Lipid extractions of  $100\mu\text{L}$  of plasma were performed by a biphasic extraction with methanol/methyl-tert-butyl ether (MTBE). Solvents were added automatically to the samples and after the appropriate shaking and centrifugation steps the supernatants were dispensed in 96-well plates and stored until analysis with GC-MS, LC-MS and NMR. Internal standards for GC and LC were previously dispensed by the robot to the same plates where supernatants were collected. Quality controls (i.e. pool of samples) were used in both GC and LC to discard drift in the instrumental response.

### **Lipid $^1\text{H}$ -NMR profiling**

Samples were prepared following the procedure previously published<sup>14</sup>.  $^1\text{H}$ -NMR spectra were recorded at 300K on an Avance III 600 spectrometer (Bruker, Germany) operating at a proton frequency of 600.20 MHz using a 5 mm PBBO gradient probe. Lipid samples were measured and recorded in PROCNO 11 using a simple pre-saturation sequence (recycle delay (RD)– $90^\circ$ –ACQ pre-saturation pulse (zgpr) program) to eliminate the residual water moisture of deuterated methanol. After pre-

processing and visual checking of the NMR dataset, specific  $^1\text{H}$  regions of diacylglycerols, triglycerides and total lipids based on terminal methyl and methylene signals were identified in the spectra using a comparison in the AMIX 3.9 software (Bruker, Germany). Curated identified regions across the spectra were integrated using the same AMIX 3.9 software package and exported to excel spreadsheets to obtain relative concentrations.

### **Lipid LC-MS profiling**

The lipid species in plasma samples were determined by ultra-high performance liquid chromatography (UHPLC) coupled to quadrupole-time of flight (qTOF) high resolution mass spectrometry (MS) (6550 iFunnel series, Agilent Technologies, Spain). The ionization was performed in positive electrospray, and the mass calibration reference was used in all the analyses to keep the mass accuracy below 5 ppm. Lipids were separated in a C18 reversed phase column (Kinetex C18-EVO from Phenomenex) and a ternary mobile phase (water/methanol/2-propanol) was used. The lipids measurements were generated from specific RT, isotope peaks relation and the most intense adduct form observed. Each lipid was quantified with an internal standard calibration method using one analytical standard and one deuterated internal standard for each lipid family. Specific vendor software was used (Quantitative Mass Hunter from Agilent).

### **Aqueous GC-MS profiling**

Samples were analysed in a 7890A Series GC coupled to a triple quadrupole (QqQ) (7000 series; Agilent Technologies, Barcelona, Spain). The chromatographic column was a J&W Scientific HP5-MS (30 m x 0.25 mm i.d., 0.25  $\mu\text{m}$  film; Agilent Technologies, Barcelona, Spain), and helium (99.999% purity) was used as a carrier

gas. Ionization was carried out with electronic impact recording data in “Full Scan” mode.

Metabolites measurements were based on specific RT plus ion fragmentation pattern. Quantification was performed by internal standard calibration, using the corresponding analytical standard for each determined metabolite and a deuterated internal standard depending on the family of metabolite. The internal standards used were succinic d<sub>4</sub> acid, glycerol <sup>13</sup>C<sub>3</sub>, norvaline, L-methionine-(carboxy-<sup>13</sup>C, methyl-d<sub>3</sub>), D-glucose <sup>13</sup>C<sub>6</sub>, myristic-d<sub>27</sub> acid and alpha-tocopherol d<sub>6</sub>.

### **Sleep, physical activity and sedentary time**

Subjects were asked to wear an ActiGraph™ tri-axis accelerometer monitor (GT3X+, Pensacola, FL, USA) tightly on the right hip using an elastic belt for seven consecutive days and eight nights (entire 24-hour period); they were only allowed to remove it during water activities (i.e. showering or swimming). At the end of the observation period, data were re-integrated to 60-second epochs and analyzed using ActiLife6 (the ActiGraph 2012, ActiLife version 6). Subjects were instructed to keep logs for bedtime (“lights off” and trying to sleep) and waking time (“lights on”) during the week in which the monitor was worn. To estimate accelerometer determined sleep duration, the self-reported bedtimes and waking times were used as the possible window of sleep and accelerometer data within this window were scored in ActiLife6 using the algorithm by Sadeh et al. 1994<sup>15</sup>. The weekly average of sleep duration was calculated in the proportion of five to two between weekdays (Sunday to Thursday) and weekend days (Friday and Saturday). Sleep duration was only considered valid if it was measured for a minimum of one weekday and one weekend night. The intra-subject standard deviation of the sleep duration was used to represent sleep variability.

Before analysis of physical activity and sedentary time, self-reported sleep duration was removed as well as non-wear time defined as 60 min of consecutive zeros using vector magnitude, allowing for 2 min of non-zero interruptions with a maximum of 100 counts/minute (cpm). Total physical activity was expressed as vector magnitude of the total tri-axial counts from monitor wear time divided by monitor wear time. Time spent in a sedentary state, doing light physical activity and moderate-to-vigorous physical activity (MVPA) were defined as all minutes showing <200, 200 to 2689 and  $\geq 2690$  vector magnitude cpm, respectively<sup>16</sup>. The percentage of time spent in a sedentary state was calculated by dividing sedentary time with monitor wear time and multiplying by 100. The percentage of time spent in light physical activity and MVPA was calculated in a similar way. The weekly averages of total physical activity, light physical activity, MVPA and sedentary time were calculated in the proportion of five to two between weekdays (Monday to Friday) and weekend days (Saturday and Sunday) and were only considered valid if monitor wear time was at least 10 hours/day (excluding sleep duration) for a minimum of one weekday and one weekend day.

## **2.5. Anthropometrical and biochemical measurements**

Anthropometric measures (body mass index, waist circumference and sagittal diameter) were determined by trained personnel with calibrated equipment. Body composition was assessed by dual-energy X-ray absorptiometry (DXA) (Lunar Prodigy Primo, GEHealthcare, Little Chalfond, UK (in Reus participants) and GE Lunar iDXA, Encore software version 16.2 (in Copenhagen participants)). Blood pressure was measured using a validated semiautomatic oscillometer (Omron HEM-705CP; Omron Healthcare). Glucose and insulin concentrations were measured using standard enzymatic automated methods and insulin resistance was estimated<sup>17</sup>. Plasma

CRP (C-reactive protein), IL-6, leptin and adiponectin were determined using a MILLIPLEX MAP Plex Kit (Merck Millipore, Billerica, MA).

### **Dietary assessment**

The dietary intake was based on three-day dietary record (3DDR) including two workdays and a weekend day. The records were inspected for clarification immediately after receiving it by trained dietitians. All foods noted in these diaries were coded to foods listed in country specific food databases<sup>18-21</sup>.

### **Statistical analyses**

Baseline characteristics of study participants were described as means and standard deviations (SD) (normally distributed continuous variables) or median and interquartile range (not normally distributed continuous variables), and percentages or numbers for categorical variables. Individual metabolites with more than 20% missing values were excluded and for the rest of metabolites missing values were imputed using the random forest imputation approach (“missForest” R package version 3.3.1). The levels of 159 metabolites were normalized and scaled to multiples of 1 SD with the rank-based inverse normal transformation. Participants were grouped into two groups based on sleep duration and variability quintiles: "low sleep duration or variability" (1<sup>st</sup>-2<sup>nd</sup> quintile) and "high sleep duration or variability" (4<sup>th</sup>-5<sup>th</sup> quintile). Due to the number of metabolites exceeding the number of observations and the collinear nature of the data, logistic regression with elastic net penalty implemented in the “glmnet” R package version 3.3.3 (alpha = 0.5) was used to build a discrimination model for high sleep duration and high sleep variability. We performed 10-fold cross-validation to find the optimal value of the tuning parameter [ $\lambda$  (lambda)] that yielded the minimum mean-squared error (MSE)<sup>22</sup>. The values minMSE and minMSE + 1SE were estimated using the argument s = “lambda.min” in the cv.glmnet function. The

discrimination model scores were computed as the weighted sum of all metabolites with weights equal to the regression coefficients from the discrimination models.

To estimate the prediction accuracy we split the data into 90% set and 10% set. Within the 90% set, we used the same elastic net procedure we used to build the model. Another 10-fold cross-validation was used to tune the model parameters. Then, we used the outer 10% set to evaluate the model built at the previous step. This procedure guarantees that the outer 10% set is completely separated from the model building procedure, so the discriminative accuracy estimated in this step is unbiased. We then repeated all these steps for 10 times and averaged their discriminative accuracy in the 10% set. Since each of them is unbiased estimate of discriminative accuracy, the average is also unbiased. In addition, we calculated the receiver operator characteristic (ROC) curves to discriminate between sleep duration and sleep variability categories in the 10% set. The area under the curve (AUC) was calculated to estimate the discriminative accuracy.

We also treated sleep duration and sleep variability as continuous variables (sleep duration was normally distributed and sleep variability was log transformed prior to analyses), and linear regression with elastic net penalty was performed in order to examine associations with metabolites. The variance of aforementioned sleep variables explained by the metabolites was estimated from the adjusted R squared by including all selected metabolites in the model. All analyses were performed using R statistical package (version 3.1.1; R Development Core Team, 2011; <http://cran.r-project.org>).

## Results

### Participant characteristics

The general characteristics of study subjects are shown in **Table 1**. The sleep duration and sleep variability groups were similar with respect to most characteristics with the exception of age and MVPA, which were higher in the group of low sleep duration as compared with the high sleep duration group, while in the latter group participants were more likely to be women ( $p < 0.05$ ) (**Table 1**). In the high sleep duration group, sleep duration was also higher ( $8.14 \pm 0.68\text{h}$ ) than the other group ( $6.49 \pm 0.48\text{h}$ ) ( $p < 0.001$ ), but sleep variability did not significantly differ. On the other hand, sleep variability was higher in the high sleep variability group ( $1.52 \pm 0.42\text{h}$ ) compared with the other group ( $0.61 \pm 0.18\text{h}$ ) ( $p < 0.001$ ). However, sleep duration was not significantly different between the two sleep variability groups.

### Associations between metabolites and high sleep duration

**Supplemental figure S1a** shows both the number of metabolites in the models and MSE as functions of  $\lambda$ . The figures also show the location of minMSE and minMSE + 1SE. **Table 2** shows selected metabolites ( $n=12$ ) ranked from the highest to the lowest elastic net positive and negative regression coefficients for high sleep duration. Undecanoylcarnitine/glutaconylcarnitine, isoundecanoylcarnitine, stearic acid, glycerol and docosahexaenoic acid were positively associated; while tiglylcarnitine/3-methylcrotonylcarnitine, 2-hydroxybutyric acid, sphingomyelin 42:1, butyrylcarnitine, sphingomyelin 33:1, phosphatidylcholine 34:2, and 3-hydroxybutyric acid, were negatively associated with high sleep duration.

### Associations between metabolites and high sleep variability

**Supplemental figure S1b** shows both the number of metabolites in the models, the MSE and the location of minMSE and minMSE + 1SE. **Table 3** shows selected

metabolites (n=10) ranked from the highest to the lowest elastic net positive and negative regression coefficients for high sleep variability. Positive regression coefficients were found for six metabolites, one lysophosphatidylcholine (20:1), one phosphatidylcholine (42:5), sucrose, two acylcarnitines (octadecenyl-, hydroxyhexadecanoyl-), and glutamic acid; while negative for two phosphatidylcholine species (40:6, 37:4), heptanoylcarnitine, and triacylglycerol 52:4.

#### **Discrimination of sleep duration and sleep variability categories [high (4<sup>th</sup> plus 5<sup>th</sup> quintile) vs. low (1<sup>st</sup> plus 2<sup>nd</sup> quintile)]**

To explore the discriminative ability of the multi-metabolite models, AUC analyses were carried out. The AUC was 0.69 (95% CI 0.62–0.75) and 0.63 (95% CI 0.53–0.72) in the multi-metabolite score, for high sleep duration and sleep variability, respectively. **Supplemental figure S2a** shows ROC curves for each of the 10 iterations together with the 10-fold cross-validated ROC curve for sleep duration categories and **supplemental figure S2b** shows ROC curves for each of the 7 iterations for sleep variability categories.

#### **Associations between metabolites and sleep duration / sleep variability (continuous)**

When we treated sleep duration as a continuous variable, three metabolites were selected from the elastic net regression with positive and negative regression coefficients (**Table 4**). These results were consistent with those of the primary analysis in relation to selected metabolites. In addition, the metabolites explained 7% of variance in sleep duration. On the other hand, the elastic regression model did not select any metabolites for sleep variability.

## Discussion

In the present study specific metabolites associated with sleep duration and sleep variability were identified among adults of the SATIN study. It was also demonstrated that these metabolite traits, together, contribute information both distinct and complementary to each other, with potential implications for discriminating these sleep characteristics. Despite of these associations, the discriminative ability of the multi-metabolite model for high sleep duration and sleep variability was moderate.

Concerning metabolites positively associated with high sleep duration, acylcarnitines are known transmembrane transporters of fatty acids across the mitochondrial membrane<sup>23</sup>. The highest association with sleep duration, concerning acylcarnitines was found for a mono-unsaturated short-chain species 5:1-DC: undecanoylcarnitine (mesaconylcarnitine) or glutaconylcarnitine, as well as an odd-numbered medium-chain species, is oundecanoylcarnitine (iso-C11:0). Changes in acylcarnitines during sleep or sleep deprivation indicate a role for the carnitine system and fatty-acids  $\beta$ -oxidation in sleep/wake regulation<sup>8</sup>. However, we cannot exclude that a potential increase in energy demands induced by low sleep duration leads a higher oxidation of fatty acids, may increase intermediary acylcarnitines production. It has been previously shown that 5 d of insufficient sleep (5-h sleep loss condition) increases energy needs. Specifically, an increase of  $\sim 5\%$  ( $\sim 111$  kcal/d) in 24-h energy expenditure was observed during sleep loss compared with 9-h control<sup>24</sup>. On the other hand, this physiologically meaningful difference was not detected in prior research<sup>25,26</sup>. In our study, we have two sleep categories (8.14h vs. 6.49h) and it is unknown whether a habitual sleep of 6.49h would induce significant increases in energy demands as compared to 8.14h.

Stearic acid (C18:0), another metabolite positively associated to sleep duration in the present study, was among fatty acids (14:0, 16:0, 16:1, 18:1, 18:2, 18:3, 22:5, C22:6) found significantly reduced in chronically sleep-restricted rats; possibly associated with lower insulin secretion from pancreas islets<sup>27</sup>. Concerning glycerol, the backbone of triacylglycerol, on which fatty acids esterify, it was positively associated with sleep duration. Overall, glycerol and free fatty acids were reported to decrease progressively during sleep; however, they may decrease during early sleep, and subsequently rise in a growth hormone-dependent manner<sup>28</sup>.

Finally, docosahexaenoic acid (DHA), an omega-3 fatty acid and major component of neural tissues, is necessary for the transformation of serotonin into melatonin and it has been suggested to influence sleep physiology<sup>29</sup>. Higher maternal plasma DHA levels during pregnancy have been associated with more mature sleep patterns in newborn babies<sup>30</sup>. Lower blood concentrations of DHA can predict more serious sleep problems in otherwise healthy children; while, dietary supplementation with DHA for 16 weeks can improve parent-rated sleep in children with such problems, and increase sleep duration by almost one hour compared to placebo<sup>31</sup>. The evidence from these investigations indicates that DHA supplementation may play a positive role in improving sleep parameters, at least in children<sup>31</sup>.

Concerning metabolites negatively associated with high sleep duration, among acylcarnitine species (tiglyl- and butyryl-) the main was tiglylcarnitine or 3-methylcrotonylcarnitine (C5:1), a molecule associated with beta-ketothiolase deficiency and ketoacidosis<sup>32</sup>. Another metabolite, 2-hydroxybutyric acid, was reported to be raised significantly in the blood following sleep deprivation; while it was also identified as an early metabolic marker of type 2 diabetes, inhibiting the release of insulin from  $\beta$ -cells<sup>33</sup>. On the other hand, elevated 2-hydroxybutyric acid

may be a by-product of the breakdown of the sulphur amino-acid metabolite, cystathionine, formed in excess during oxidative stress<sup>34</sup>. These associations are consistent with the viewpoint that sleep deprivation contributes to metabolic stress and diabetes development.

Two sphingomyelins (SM 42:1, 33:1) and one phosphatidylcholine (PC 34:2) were negatively associated with sleep duration. SMs are hydrolysed by sphingomyelinases activated by inflammatory cytokines and oxidative stress, to produce ceramides that further activate enzymes involved in stress signalling and apoptosis<sup>35</sup>. PCs are direct substrates for the formation of SMs by SM-synthase, which transfers the choline-phosphate group to ceramides<sup>36</sup>. Finally, 3-hydroxybutyric acid (beta hydroxybutyrate) is one of the ketone bodies and metabolism of these substances affects sleep homeostasis in mice<sup>37</sup> or is affected by ketosis due to low sleep duration. Among metabolites positively associated with high sleep-variability, acylcarnitines octadecenylcarnitine (acylcarnitine C18:1) and hydroxyhexadecanoyl-carnitine were positively associated, while heptanoylcarnitine negatively associated. Phosphatidylcholines PC 42:5 and LPC 20:1 were positively associated, while PC 40:6 and PC 37:4 were negatively associated. The mechanisms linking the aforementioned carnitine, PC and LPC molecules to sleep require further investigation.

Sucrose and glutamic acid were recorded to be positively associated with high sleep-variability. The levels of sucrose in the blood may reflect gastric permeability to sucrose<sup>38</sup>. Finally, glutamate is associated with sleep/wake regulation and cortical projections of the glutamatergic system are important for cortical activation and wakefulness<sup>39</sup>. In the brain itself, in numerous regions, levels of glutamate are highest

during wakefulness or rapid eye movement sleep<sup>40</sup>. Abnormal glutamate homeostasis may be an important contributor to a higher risk of developing diabetes<sup>41</sup>.

The results of the present study should be interpreted in the context of its limitations and strengths. First, the cross-sectional design does not allow making any causal inference of the observed associations, and therefore both directions are plausible. Second, despite participants were not diagnosed for obstructive sleep apnea (OSA) we cannot completely discard it. An exponential pattern was previously observed with the hazard ratio for OSA development increasing with body mass index (BMI)<sup>42</sup>. Since BMI did not significantly differ between sleep duration or sleep variability categories we assume that OSA prevalence is not substantially different between these categories and therefore may not influence the reported findings. Third, the lack of external validation in our study may have increased the risk of overfitting of the model. To decrease the risk of overfitting and get an unbiased estimate of prediction accuracy, we have internally cross validated the models using a 10-fold cross-validation approach. Although we found a moderate discriminating performance, this multi-metabolite model should be tested in a larger sample size with potentially better discriminating accuracy. Fourth, participants were overweight/obese individuals without comorbidities and this may limit the generalizability of the findings to other populations. Regarding strengths, we have used a multi-metabolomics approach in order to cover a wide range of metabolites.

In conclusion, the present study revealed associations of specific metabolites with sleep duration and sleep variability in overweight/obese adults. The exact mechanisms of association for the selected metabolites and sleep-patterns remain to be determined.

Further studies are needed to confirm our findings and assess the potential involvement of the identified metabolites in sleep-related metabolic health.

**Abbreviations:**  $^1\text{H}$  NMR: proton nuclear magnetic resonance; LC-MS: liquid chromatography coupled to mass spectrometry; GC-MS: gas chromatography coupled to mass spectrometry; LPC: lysophosphatidylcholine; PC: phosphatidylcholine; DHA: docosahexaenoic acid; SM: sphingomyelins

#### **Funding sources**

The SATIN project is funded by a grant of the Seventh Framework Programme for "Cooperation": Food, Agriculture and Biotechnology of the European Union (2012-2016).

#### **Acknowledgments**

Metabolomic analyses were performed with the technological infrastructure of the Centre for Omic Sciences (COS). The authors are indebted to Jo Harrold (Institute of Psychology Health and Society, University of Liverpool, UK) for her intellectual contributions to this manuscript.

#### **Disclosure statement**

Financial Disclosure: none.

Non-financial Disclosure: none.

## References

1. Ogilvie RP, Redline S, Bertoni AG, et al. Actigraphy Measured Sleep Indices and Adiposity: The Multi-Ethnic Study of Atherosclerosis (MESA). *Sleep*. 2016;39(9):1701-1708.
2. Garfield V, Llewellyn CH, Steptoe A, Kumari M. Investigating the Bidirectional Associations of Adiposity with Sleep Duration in Older Adults: The English Longitudinal Study of Ageing (ELSA). *Sci Rep*. 2017;7:40250.
3. Shan Z, Ma H, Xie M, et al. Sleep duration and risk of type 2 diabetes: a meta-analysis of prospective studies. *Diabetes Care*. 2015;38(3):529-537.
4. Scheer FA, Hilton MF, Mantzoros CS, Shea SA. Adverse metabolic and cardiovascular consequences of circadian misalignment. *Proc Natl Acad Sci U S A*. 2009;106(11):4453-4458.
5. Gale JE, Cox HI, Qian J, Block GD, Colwell CS, Matveyenko AV. Disruption of circadian rhythms accelerates development of diabetes through pancreatic beta-cell loss and dysfunction. *J Biol Rhythms*. 2011;26(5):423-433.
6. Kim M, Sasai H, Kojima N, Kim H. Objectively measured night-to-night sleep variations are associated with body composition in very elderly women. *J Sleep Res*. 2015;24(6):639-647.
7. Medic G, Wille M, Hemels ME. Short- and long-term health consequences of sleep disruption. *Nat Sci Sleep*. 2017;9:151-161.
8. Davies SK, Ang JE, Revell VL, et al. Effect of sleep deprivation on the human metabolome. *Proc Natl Acad Sci U S A*. 2014;111(29):10761-10766.
9. Weljie AM, Meerlo P, Goel N, et al. Oxalic acid and diacylglycerol 36:3 are cross-species markers of sleep debt. *Proc Natl Acad Sci U S A*. 2015;112(8):2569-2574.
10. Bell LN, Kilkus JM, Booth JN 3rd, Bromley LE, Imperial JG, Penev PD. Effects of sleep restriction on the human plasma metabolome. *Physiol Behav*. 2013;122:25-31.
11. Bailey SM, Udoh US, Young ME. Circadian regulation of metabolism. *J Endocrinol*. 2014;222(2):75-96.
12. Ray S, Reddy AB. Cross-talk between circadian clocks, sleep-wake cycles, and metabolic networks: Dispelling the darkness. *Bioessays*. 2016;38(4):394-405.
13. Xiao Q, Derkach A, Moore SC, Zheng W, Shu X, Gu F, et al. Habitual Sleep and human plasma metabolomics. *Metabolomics*. 2017;13:63. DOI <https://doi.org/10.1007/s11306-017-1205-z>

14. Mallol R, Rodríguez MA, Heras M, et al. Surface fitting of 2D diffusion-edited <sup>1</sup>H NMR spectroscopy data for the characterisation of human plasma lipoproteins. *Metabolomics*. 2011;7:572-582.
15. Sadeh A, Sharkey M, Carskadon MA. Activity-based sleep-wake identification: an empirical test of methodological issues. *Sleep*. 1994;17(3):201-207.
16. Sasaki JE, John D, Freedson PS. Validation and comparison of ActiGraph activity monitors. *J Sci Med Sport*. 2011;14(5):411-406.
17. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 1985;28(7):412-419.
18. Mataix J. *Tablas de Composición de Alimentos*. 4. Granada, Spain: Universidad de Granada; 2003.
19. Moreiras O, Carvajal A, Cabrera L (Eds) *Tablas de Composición de Alimentos [FoodCompositionTables]* 9. Madrid, Spain: Ediciones Pirámide; 2005.
20. Centre d'Ensenyament Superior de Nutrició i Dietética. *Universitat de Barcelona-McGraw-Hill*; Barcelona, Spain: 2004. *Tablas de composición de alimentos del CESNID*.
21. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4628270/>
22. Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via Coordinate Descent. *J Stat Softw*. 2010;33(1):1-22.
23. Longo N, Frigeni M, Pasquali M. Carnitine transport and fatty acid oxidation. *Biochim Biophys Acta*. 2016;1863(10):2422-2435.
24. Markwald RR, Melanson EL, Smith MR, Higgins J, Perreault L, Eckel RH, Wright KP Jr. Impact of insufficient sleep on total daily energy expenditure, food intake, and weight gain. *Proc Natl Acad Sci U S A*. 2013;110(14):5695-5700.
25. Nedeltcheva AV, Kilkus JM, Imperial J, Kasza K, Schoeller DA, Penev PD. Sleep curtailment is accompanied by increased intake of calories from snacks. *Am J Clin Nutr*. 2009;89(1):126-133.
26. St-Onge MP, Roberts AL, Chen J, Kelleman M, O'Keeffe M, RoyChoudhury A, Jones PJ. Short sleep duration increases energy intakes but does not change energy expenditure in normal-weight individuals. *Am J Clin Nutr*. 2011;94(2):410-416.
27. Zhan S, Wu Y, Sun P, Lin H, Zhu Y, Han X. Decrease in Circulating Fatty Acids Is Associated with Islet Dysfunction in Chronically Sleep-Restricted Rats. *Int J Mol Sci*. 2016;17(12). pii: E2102.
28. Mesarwi O, Polak J, Jun J, Polotsky VY. Sleep disorders and the development of insulin resistance and obesity. *Endocrinol Metab Clin North Am*. 2013;42(3):617-634.
29. Peuhkuri K, Sihvola N, Korpela R. Diet promotes sleep duration and quality. *Nutr Res*. 2012;32(5):309-319.
30. Cheruku SR, Montgomery-Downs HE, Farkas SL, Thoman EB, Lammi-Keefe CJ. Higher maternal plasma docosahexaenoic acid during pregnancy is associated with more mature neonatal sleep-state patterning. *Am J Clin Nutr*. 2002;76(3):608-613.

31. Montgomery P, Burton JR, Sewell RP, Spreckelsen TF, Richardson AJ. Fatty acids and sleep in UK children: subjective and pilot objective sleep results from the DOLAB study-a randomized controlled trial. *J Sleep Res.* 2014;23(4):364-388.
32. Abdelkreem E, Otsuka H, Sasai H, Aoyama Y, Hori T, Abd El Aal M, Mahmoud S, Fukao T. Beta-Ketothiolase Deficiency: Resolving Challenges in Diagnosis. *Journal of Inborn Errors of Metabolism & Screening.* 2016;4:1-9.
33. Cobb J, Eckhart A, Motsinger-Reif A, Carr B, Groop L, Ferrannini E.  $\alpha$ -Hydroxybutyric Acid Is a Selective Metabolite Biomarker of Impaired Glucose Tolerance. *Diabetes Care.* 2016;39(6):988-995.
34. Gall WE, Beebe K, Lawton KA, et al. alpha-hydroxybutyrate is an early biomarker of insulin resistance and glucose intolerance in a nondiabetic population. *PLoS One.* 2010;5(5):e10883.
35. Cutler RG, Kelly J, Storie K, et al. Involvement of oxidative stress-induced abnormalities in ceramide and cholesterol metabolism in brain aging and Alzheimer's disease. *Proc Natl Acad Sci U S A.* 2004;101(7):2070-2075.
36. Li Z, Vance DE. Phosphatidylcholine and choline homeostasis. *J Lipid Res.* 2008;49(6):1187-1194.
37. Chikahisa S, Shimizu N, Shiuchi T, Séi H. Ketone body metabolism and sleep homeostasis in mice. *Neuropharmacology.* 2014;79:399-404.
38. Giofre MR, Meduri G, Pallio S, et al. Gastric permeability to sucrose is increased in portal hypertensive gastropathy. *Eur J Gastroenterol Hepatol.* 2000;12:529-533.
39. Brown RE, Basheer R, McKenna JT, Strecker RE, McCarley RW. Control of sleep and wakefulness. *Physiol Rev.* 2012;92(3):1087-1187.
40. Watson CJ, Lydic R, Baghdoyan HA. Sleep duration varies as a function of glutamate and GABA in rat pontine reticular formation. *J Neurochem.* 2011;118(4):571-580.
41. Xiao Q, Moore SC, Keadle SK, et al. Objectively measured physical activity and plasma metabolomics in the Shanghai Physical Activity Study. *Int J Epidemiol.* 2016; 45(5): 1433-1444.
42. Langton RS, Neyra J, Downs JW, Niebuhr DW. The Relationship Between Enlistment Body Mass Index and the Development of Obstructive Sleep Apnea in the U.S. Military. *Mil Med.* 2016;181(8):913-919.

**Table 1.** Characteristics of the study population

	Low sleep duration (n=82)	High sleep duration (n=82)	Low sleep variability (n=82)	High sleep variability (n=82)
<b>Sleep parameters</b>				
Sleep duration (hour)	6.49 (0.48)	8.14 (0.68) <sup>b</sup>	7.27 (0.79)	7.28 (1.04)
Sleep variability (hour)	1.06 (0.46)	1.08 (0.59)	0.61 (0.18)	1.52 (0.42) <sup>b</sup>
Age (years)	48 (10)	43 (11) <sup>a</sup>	47 (10)	46 (11)
Sex (% Women)	71.9	86.6 <sup>a</sup>	81.7	75.6
Body mass index (kg/m <sup>2</sup> )	31.07 (2.23)	30.87 (1.91)	31.14 (2.26)	30.92 (1.91)
Waist circumference (cm)	101.94 (9.44)	99.85 (8.71)	100.74 (9.11)	101.16 (9.15)
Sagittal diameter	23.15 (2.46)	22.96 (2.29)	22.96 (2.29)	23.11 (2.33)
Total fat (kg)	36.32 (5.83)	37.04 (5.28)	36.81 (6.14)	36.04 (4.89)
Lean mass (kg)	47.32 (10.92)	46.30 (7.91)	47.03 (8.22)	47.21 (10.34)
Moderate-to-vigorous PA (min/d)	59.94 (29.42) <sup>a</sup>	49.70 (25.71)	54.72 (28.08)	51.34 (29.71)
Sedentary time (min/day)	509.35 (116.82)	503.55 (101.36)	509.99 (98.20)	517.21 (111.64)
Total body fat (%E)	35.50 (5.45)	35.24 (6.78)	35.43 (5.51)	35.38 (6.46)
Saturated fat (%E)	11.67 (0.34)	11.85 (0.37)	11.63 (3.39)	11.55 (3.00)
Monounsaturated fat (%E)	13.41 (0.37)	12.87 (0.44)	13.35 (3.01)	13.00 (4.13)
Polyunsaturated fat (%E)	5.05 (0.19)	4.96 (0.17)	4.96 (1.54)	5.01 (1.72)
Protein (%E)	19.43 (5.21)	18.04 (3.35)	18.43 (3.83)	19.06 (4.81)
Carbohydrate (%E)	42.90 (6.64)	44.21 (6.60)	44.49 (6.65)	43.37 (6.62)
Sugar intake (g)	83.61 (50.43)	80.60 (38.76)	84.76 (36.99)	79.41 (53.07)
HOMA-IR	1.93 (1.04; 3.72)	1.68 (1.05; 3.13)	2.07 (1.29; 3.62)	1.68 (1.04; 3.06)
IL-6 (pg/mL)	2.04 (1.02; 2.93)	1.93 (1.36; 2.84)	1.91 (0.95; 2.54)	1.88 (1.09; 2.84)
Adiponectin (µg/mL)	11.62 (8.56)	10.01 (5.18)	10.14 (6.01)	11.13 (6.70)
Leptin (ng/mL)	3.57 (1.33; 7.91)	2.97 (1.22; 8.00)	2.81 (1.33; 7.21)	3.47 (1.24; 8.76)

Data are presented as mean (SD), median (interquartile range) unless otherwise indicated.

The  $\chi^2$  test was used for comparison of categorical variables and Student's *t*-test or Mann-Whitney U test was used for comparison of continuous variables.

<sup>a</sup>( $P < 0.05$ )

<sup>b</sup>( $P < 0.001$ )

Low sleep duration: 1<sup>st</sup> plus 2<sup>nd</sup> quintile.

High sleep duration: 4<sup>th</sup> plus 5<sup>th</sup> quintile.

Low sleep variability: 1<sup>st</sup> plus 2<sup>nd</sup> quintile.

High sleep variability: 4<sup>th</sup> plus 5<sup>th</sup> quintile.

Abbreviations: *PA* physical activity, *HOMA-IR* homeostasis model of insulin resistance, *IL-6* interleukin 6, *E* total energy intake.

**Table 2.** Metabolites ranked from the highest to the lowest elastic net positive or negative regression coefficients for high sleep duration (4<sup>th</sup> plus 5<sup>th</sup> quintile)

<i>Metabolite</i>	<i>Coefficient</i>	<i>Metabolite</i>	<i>Coefficient</i>
Undecanoylcarnitine/Glutaconylcarnitine/ isoundecanoylcarnitine	0.0862	Tiglylcarnitine/3-methylcrotonylcarnitine	-0.2791
Stearic acid	0.0814	2-Hydroxybutyric acid	-0.1169
Glycerol	0.0436	SM 42:1	-0.0971
Docosahexaenoic acid	0.0008	Butyrylcarnitine	-0.0772
Serotonin	0.0003	SM 33:1	-0.0217
		PC 34:2	-0.0183
		3-Hydroxybutyric acid	-0.0072

Abbreviations: *PC* phosphatidylcholine, *SM* sphingomyelin.

**Table 3.** Metabolites ranked from the highest to the lowest elastic net positive or negative regression coefficients for high sleep variability (4<sup>th</sup> plus 5<sup>th</sup> quintile)

<i>Metabolite</i>	<i>Coefficient</i>	<i>Metabolite</i>	<i>Coefficient</i>
Octadecenylcarnitine	0.2809	PC 40:6	-0.2695
Sucrose	0.1529	Heptanoylcarnitine	-0.0561
Hydroxyhexadecanoylcarnitine	0.1313	PC 37:4	-0.0319
LPC 20:1	0.1143	TAG 52:4	-0.0019
PC 42:5	0.0277		
Glutamic acid	0.0201		

Abbreviations: *LPC* lysophosphatidylcholine, *PC* phosphatidylcholine, *PE* phosphatidylethanolamine, *SM* sphingomyelin, *TAG* triacylglycerol.

**Table 4.** Metabolites ranked from the highest to the lowest elastic net positive or negative regression coefficients for sleep duration (continuous)

<i>Metabolite</i>	<i>Coefficient</i>	<i>Metabolite</i>	<i>Coefficient</i>
Glycerol	0.0001	Tiglylcarnitine/3-methylcrotonylcarnitine	-0.0665
		3-Hydroxybutyric acid	-0.0122

Accepted Manuscript