

Title

The effects of a participatory intervention in mental health units on nurses' perceptions of the practice environment: a mixed methods study

Abstract

Background: The practice environment influences the quality of care and the nursing outcomes achieved in their workplaces.

Aim: To examine the perception of the clinical practice environment among nurses working in mental health units in the context of their participation in an action research study aimed at improving the nurse-patient relationship.

Methods: An explanatory sequential mixed-methods study was designed. The data were collected in three phases in 18 mental health units (n= 95 nurses). Quantitative data were collected through the Practice Environment Scale of the Nursing Work Index, and qualitative data were collected through reflective diaries and focus groups in the context of Participatory Action Research.

Results: The nurses' assessment of their practice environment shifted from positive to negative. Nurse manager leadership was the aspect that worsened the most. In addition, the perception of their participation in the affairs of the centre and nursing foundations for quality of care decreased. The nurses considered it essential to be able to influence decision-making bodies and that the institution should promote a model of care that upholds the therapeutic relationship in actual clinical practice.

Conclusions: Nurses perceived that they should be involved in organisational decisions and required more presence and understanding from managers. Furthermore, nurses stated that institutions should promote nursing foundations for quality of care. This study contributes to understanding how nurses in mental health units perceive their

Mental health nurses' perceptions on the practice environment

work environment and how it affects the improvement of the nurse-patient relationship in clinical practice.

Key words

Evidence-Based practice, Inpatient Psychiatric Care, Patient-Centered Care, Quality Improvement

Manuscript text

INTRODUCTION

The practice environment is a key element that influences the quality of nursing care, job satisfaction and the retention of nurses in health services (Oshodi, Crockett, Bruneau, & West, 2017; Roche et al., 2016). From a theoretical perspective, it was Kanter (1993) who, in his theory of structural empowerment, argued that conditions in the work environment such as access to information and resources, support from managers and opportunities to learn and develop were necessary for the proper development of employees' work (Kanter, 1993). In addition, other authors have also conceptualized the "working environment" as "the organizational characteristics of a work setting that facilitate or constrain professional nursing practice" (Lake, 2002, p. 178). Thus, the practice environment involves factors such as leadership, teamwork, control and autonomy in clinical practice, conflict management, professional relationships, patient communication, work motivation and cultural awareness (Erickson, Duffy, Ditomassi, & Jones, 2009). All of these can facilitate or constrain nursing practice (Lake, 2002). Consequently, bearing in mind that these factors are modifiable in work environments (Lake & Friese, 2006), it is important to identify them in specific contexts, such as acute mental health units, in order to improve the clinical practice of nurses.

The working environment is a major determining factor for improving the quality of patient care, while improving outcomes for nurses (Aiken et al., 2011). In this regard, positive work environments are associated with a number of patient outcomes including: greater satisfaction (Aiken et al., 2012), improved quality of care (Aiken et al., 2011) and a lower rate of adverse events (Copanitsanou, Fotos, & Brokalaki, 2017). Indeed, even at the primary care level, the nursing practice environment has been

observed to affect the quality of chronic disease care (Poghosyan, Norful, Liu, & Friedberg, 2018). Conversely, unfavourable working environments have been associated with higher readmissions, higher health care costs and associated complications (Copanitsanou et al., 2017).

In relation to the impact of the clinical environment on nurses, it is known that positive work environments contribute to greater job satisfaction (Lin, Lai, Huang, Huang, & Hsieh, 2020) and lower burnout among professionals (Nantsupawat et al., 2017). In general, studies indicate that nurses perceive most aspects of their practice environment positively (Al-Maaitah, AbuAlRub, & Al Blooshi, 2018; Dorigan & Guirardello, 2017; Numminen et al., 2016). Specifically, nurses tend to report positive perceptions of team relationships with physicians (Al-Maaitah et al., 2018; Numminen et al., 2016) and, conversely, less positive perceptions tend to be related to staffing and adequacy of resources (Al-Maaitah et al., 2018; Nantsupawat et al., 2017; Numminen et al., 2016). In fact, nurses' positive perceptions of their practice environment are also known to be associated with increased professional competence (Numminen et al., 2016), an increased perception of the quality of care (Al-Maaitah et al., 2018; Numminen et al., 2016), greater job satisfaction (Lin et al., 2020; Nantsupawat et al., 2017) and a reduced intention to leave the job or profession (Nantsupawat et al., 2017; Numminen et al., 2016). In fact, nurses describe a more positive perception of the quality of nursing care perceived by their patients when the practice environment is healthy (Numminen et al., 2016; Roche et al., 2016). Thus, nurses believe that the role of nurse leaders is fundamental to the establishment and improvement of a healthy work environment (Ducharme, Bernhardt, Padula, & Adams, 2017; Numminen et al., 2016).

Specifically, in the field of mental health units, where the therapeutic nurse-patient relationship is key for the provision of care (Moreno-Poyato et al., 2016; Zugai, Stein-

Parbury, & Roche, 2015), understanding that the creation of protected and therapeutic spaces is of particular importance (McAllister, Robert, Tsianakas, & McCrae, 2019; Molin, Graneheim, Ringnér, & Lindgren, 2019; Moreno-Poyato, El Abidi, Rodríguez-Nogueira, Lluch-Canut, & Puig-Llobet, 2021). In fact, it has been shown that the nurses' perception of a better practice environment is related to the establishment of a higher quality nurse-patient relationship (Roviralta-Vilella, Moreno-Poyato, Rodríguez-Nogueira, Duran-Jordà, & Roldán-Merino, 2019). However, some studies indicate that mental health nurses perceive their practice environment more negatively compared to other settings (Roche & Duffield, 2010). It has even been observed that mental health nurses who use more restrictive measures with patients are less satisfied with their jobs (Kurjenluoma et al., 2017). In fact, mental health nurses' job satisfaction is related to how they perceive their work environment (Gillet et al., 2019). In addition, they experience higher levels of emotional exhaustion when they perceive their practice environment to be worse (Gabrielsson et al., 2016; Gillet et al., 2019; Wyder et al., 2017).

Consequently, the available evidence shows that positive work environments improve the quality of care, nurses' job satisfaction, and the quality of their work (Kurjenluoma et al., 2017; Wyder et al., 2017), as well as nurses' work commitment and outcomes in their workplaces (Huang, Wang, Dong, Li, & Wan, 2021). However, there is limited knowledge regarding the influence of the environment on the design and improvement of practice in the field of mental health inpatient units (Moreno-Poyato et al., 2021). Based on the knowledge that certain environmental factors are modifiable (Lake & Friese, 2006), further exploration of the involvement of the environment in the clinical practice of nurses in acute mental health units is warranted. In this sense, an approach

from a global perspective is important to obtain a better understanding of the phenomenon of nurses' perceptions and perspectives.

The aim of this study was to examine the perception of the clinical practice environment among nurses in mental health units in the context of their participation in an action research study aimed at improving the nurse-patient relationship.

METHODS

Design

This study was conducted following an explanatory sequential mixed-methods design with the intention of obtaining complementary views on the same phenomenon and integrating quantitative and qualitative components to form inferences and conclusions (Tashakkori & Teddlie, 2008; Teddlie & Tashakkori, 2012). This study is part of a project with the main aim of improving the clinical practice of the nurse-patient relationship in mental health units in Catalonia (Spain). This study was divided into three phases, with quantitative data collected in the first and third phases. In the second phase, qualitative data was collected using the Participatory Action Research (PAR) method, following the model by Kemmis and McTaggart (2008) (Fig. 1). To this end, a two-cycle process was designed, consisting of four stages in each cycle. The PAR method is based on the notion that participants' actions must be filtered through experience and reflection before their practice can be improved or changed. Thus, PAR is performed by following a sequence of stages comprising a spiral of self-reflective cycles (Kemmis & McTaggart, 2008): planning the desired change, action and observation of the process and consequences of change, reflection on these processes and their consequences, re-planning the change, action and re-observation and further reflection. During this process, participants are asked to act and reflect, in order to

simultaneously gain awareness of and transform their practice (Baum, MacDougall, & Smith, 2006). This study adheres to the GRAMMS guidelines for reporting mixed methods research.

Sample/Participants

All nurses (n=235) in the 21 acute units of the Catalan Mental Health Network were invited to participate in the study. Only resident nurses in training were excluded. Within each unit, a nurse coordinated the study and oversaw the recruitment process. 198 nurses from 18 acute care units initially participated in the study. Of these, 95 completed all three phases. For the quantitative aspect, the power of the study was calculated to detect a correlation coefficient of at least 0.3 at baseline and follow-up measurements with an alpha of 0.05, which was 80% in a bilateral contrast.

Data collection

The nurse coordinators of each unit were responsible for collecting informed consent documents and nurses' email addresses, and the principal investigator provided an individual confidential participant code to each nurse. Data collection for study phase I and III took place between January 2018 and June 2019, whereas qualitative data were collected in phase II, between April 2018 and November 2018.

Quantitative data

Quantitative data collection for both phase I and phase III was conducted via an electronic form sent by email to the participants. This form comprised a questionnaire which gathered nurses' socio-demographic and professional data, together with the Practice Environment Scale of the Nursing Work Index (PES-NWI): Lake 2002), which was the instrument used to measure the perceived nursing practice environment. This instrument consists of 31 items assessed on a Likert scale ranging from 1 to 4 points (1=

strongly disagree, 4= strongly agree). Values above 2.5 indicate higher agreement, whereas values below 2.5 indicate disagreement. These items are grouped into five subscales: (1) Nurse participation in hospital affairs, (2) Nursing foundations for quality of care, which emphasises the nursing principles for high quality patient care, (3) Nurse manager ability, leadership, and support for nurses, (4) Staffing and resource adequacy, and (5) Collegial nurse-physician relationships. The PES-NWI has been validated in the Spanish population by Pedro-Gómez et al (2009). In the case of our sample, the internal consistency obtained as Cronbach's alpha values ranged from 0.77 to 0.92 for the five factors that constitute the scale.

Qualitative data

Qualitative data were collected as the PAR was being carried out through reflective diaries and focus groups. Specifically, the qualitative data came from 248 reflective diaries collected at the different stages of action and observation and 30 focus groups that took place during the reflective stages of the PAR (Fig. 1).

In relation to the reflective diaries, in order to further structure the reflection and its recording (Bolg, Dwyer, Doherty, Pignataro, & Renaud, 2020), the research team sent a self-observation guide by e-mail to each nurse, indicating how they were to record the self-observation data. The diary was to contain a description and reflection on the types of interactions to be self-observed in practice. As its purpose was to monitor the process of change triggered by the PAR, for each interaction, the nurses were to record the description of the situation, the type of verbal and non-verbal language they had used, their reflective intervention and a reflection on the influence of the environment on the interaction and how they had felt in the context of the interaction (Kemmis & Mctaggart, 2008).

The focus groups were conducted with the participating nurses from each unit and were led by the first author, a mental health nurse specialist (ARMP), with the collaboration of an observer, the second author (DTM) or the third author (FGP), depending on the unit where the group was held. The groups were conducted in a space set up in each participating centre. The focus groups were audio recorded and lasted between 1 and 2 hours. At the beginning of each session, participants were provided with a working document containing the preliminary results obtained from the reflective diaries corresponding to the previous stage of the PAR.

Ethical considerations

The study was approved by the referring ethical committees of the 18 participating units. Nurses signed the informed consent for participation. All data were collected in accordance with national and international privacy and confidentiality guidelines.

Data analysis

The quantitative data were analysed using SPSS version 27. Descriptive data are presented as the mean and standard deviation (SD) for continuous outcomes and number and percentage (%) for categorical outcomes. The paired Student's t-test was used to compare the PES-NWI between baseline and follow-up. Statistical significance was set at $P < 0.05$.

The qualitative data were analysed using NVivo version 12. The data were analysed using thematic content analysis (Crowe, Inder, & Porter, 2015). During an initial stage, the text obtained was fragmented into descriptive codes assigned according to their semantic content. In the second stage, these codes were grouped into more analytical sub-themes, by classifying the codes according to the meaning of the linguistic units and their combinations. Finally, during a third hierarchical stage, the semantic analysis

of the previous sub-themes was considered, and the codes were classified deductively according to the dimensions of the PES-NWI.

Rigour

It should be noted that reflexivity was continuous throughout the study process. The research team included both researchers with academic backgrounds and others involved in clinical practice. Therefore, this enabled a reflective and equidistant position throughout the processes of data collection and analysis. The groups were led by the first author, who had qualitative doctoral training and experience as a nurse in acute units, although without any employment relationship with the mental health services. This facilitated the creation of a peer relationship with the participants, and an atmosphere of trust. The credibility and confirmability of the data is supported by the triangulation of techniques and of the researchers in the analysis process, together with the constant auditing of the results by the participants throughout the qualitative process.

RESULTS

Description of the sample

The study was completed by 95 nurses aged between 22 and 62 years with a mean age of 33.4 years (SD=9.3). Almost 70 % of the nurses were female. Their experience in mental health was 7.7 years on average (SD=8.0). Only 23.2 % of the nurses had the official title of specialist in mental health nursing and only 22 % of the nurses had a PhD or master's degree. Up to 80 % of the nurses had a permanent employment contract. All working shifts of the centres were equally represented in the sample.

Nurses' perceptions about the clinical practice work environment

As shown in Figure 2 and Table 1, the nurses' overall assessment of the practice environment worsened significantly after the participatory process. In this regard, the scores on 4 of the 5 dimensions of the scale decreased significantly. It should be noted that, prior to the participatory process, the nurses positively scored the overall practice environment and all its dimensions, except for the adequacy of resources and staffing, which was the only item scored below neutral. However, at the end of the process, both the overall assessment of the practice environment and most of the dimensions were rated negatively.

[FIGURE 2. Differences in scores of the PES-NWI]

Nurse participation in hospital affairs

Nurses scored this dimension significantly lower following their participation in the process of change. Specifically, the items that decreased significantly were related to nurses' perceptions of being involved in management and feeling listened to by management, where the lowest rated item was having opportunities in decision-making bodies (Table 2). Although no changes were noted, one of the items that received the lowest scores in both the baseline and follow-up measures was the perception that the nurse managers had equal levels of authority and power as other management staff at the centre. In this sense, it was important for the nurses to feel the support of the organisation, not only through the resources provided to them, but also in terms of their ability to influence structural changes or plan certain actions aimed at improving their interventions and the therapeutic relationship.

"... if there is a proposal that we jointly see that it would be better for it to be done differently, then you can explain it and they say go ahead, we'll do it and

we'll try it ... I don't know... having this amount of ease to be able to change things". (12FG4P02).

However, the nurses questioned the management of the nursing directors. They felt that they did not defend the role of nurses and did not strive to give nurses their rightful place at the institutional level. Likewise, the nurses perceived that the managers did not give them the opportunity to raise the problems that the nurses faced in their daily work, even though they were the professionals who were present at all hours of the day and those who accompanied the people admitted to the acute units.

“There are many things for which nurses should be consulted more, because in the end we nurses are the ones who are there the most. On Saturdays and Sundays, you spend all your time with patients and there are no doctors anywhere, so in the end they should consider nurses' opinions more.”

(12FG4P04).

Nursing foundations for quality of care

The score given by the nurses to this dimension was also significantly lower after the participatory process. For the participants, the type of allocation of patients to individual nurses in the centres, the existence of a common nursing philosophy and the concern on behalf of managers that nurses provide quality care were the items that significantly decreased (Table 2). However, although unchanged, the lowest rated item in this dimension was the existence of a welcome and mentoring programme for new nurses. For the nurses, the position and discourse of their institutions regarding the therapeutic relationship was highly ambiguous. They indicated that their institutions did not have a clear philosophical model that defended the therapeutic relationship, stating that the nursing staff did not prioritise this in their day-to-day work either and that ultimately, what they prioritised were instrumental tasks, rather than relational ones.

“The therapeutic relationship is not strengthened, when there is no technique involved, or there is no specific need, it is not strengthened by the institution. This is why some of the staff spend a lot of time on administrative tasks instead of spending time at the bedside.” (12RD2P06).

Moreover, the nurses recognised specific training limitations in mental health, due to the difficulty of adapting the theoretical content on which the TR is based to the reality experienced in the acute units, generating insecurity in their approaches.

“Although the therapeutic relationship is highly valued, I miss refresher training courses on the therapeutic relationship.” (03RD2P09).

In addition, the nurses expressed their discomfort at the overload of having to supervise or redirect the actions of the new professionals, as they were sent to work in units they were unfamiliar with, without having prior information about the unit or sufficient knowledge to work in these environments.

“...personally, at this moment, due to my lack of experience, I feel that I still lack the resources, training and experience to resolve these situations better” (10RD2P11) “...yes, because normally, we are very busy, because with someone who has no experience, you have to explain how to do things and at the same time you have to be with this patient who is pre-agitating...” (15FG4P01).

Nurse manager ability, leadership and support of nurses

This dimension, despite continuing to be the most highly rated by the nurses after the participatory process, was the one that decreased the most. As shown in table 2, the scores for all the items that conform the dimension decreased significantly, with the item with the lowest score being the recognition and praise of work well done by

managers. The nurses expected their managers to be motivated and present in the unit for them to understand and be involved in what was happening on a day-to-day basis. However, they questioned the role of the managers, as they considered them to be absent figures, who did not defend the interests of the service, the people admitted, or the nursing staff themselves.

"Involvement is important, he should spend more time with us, he should ask how it's going, he should be involved, obviously. But I think it should be important for a supervisor to spend x hours on a shift, to see how it's going....".
(10FG4P02).

Staffing and resource adequacy

This dimension received the worst rating by the nurses at both baseline and follow-up and decreased significantly after the participatory process. Specifically, the items that worsened significantly were adequacy of staffing and time to discuss care-related problems with other nurse colleagues (Table 2). In this regard, nurses reported that lack of human resources was one of the greatest and most important constraints they had to face. This severely limited the type, quality, and quantity of interactions that nurses were able to have with the people they cared for. The lack of adequate ratios for the needs of the unit generated an overload and a risk, which affected the safety of the users, and that of the staff themselves. This often meant that many of the interactions they carried out were more extensive with those users who were more demanding or who presented more behavioural alterations, to the detriment of those who were isolated or who did not make explicit demands and who might need it just as much as the other more demanding people.

Mental health nurses' perceptions on the practice environment

"In the end, when you are saturated with work, you dedicate yourself to the most behavioural things and the person who is very sick...who is in his own world, but is calm, on the medical record you may state that he is calm when you haven't even seen him...he is still a sick person, a sick person who is suffering. I don't know how this can be solved!" (12FG4P05).

Moreover, the nurses stated that time was one of the most important and most necessary factors for the development of the TR, yet it lacked the most during their working days. The nurses perceived that this lack of time greatly affected the TR, given that not having the time they needed to dedicate to the people they cared for put the quality of the relationship and the care provided at risk.

"The lack of time to perform each of the tasks is evident as the ratio of patients per nurse is very high and the distribution of direct interventions with patients is limited." (07RD2P02).

Thus, the nurses identified that this lack of time was also related to factors of the nursing activity itself. Clinical nursing practice involves many activities that are highly varied and with different levels of complexity. Thus, the nurses pointed out that there were some tasks that were fixed and had to be carried out at certain times. Fixed schedules that often marked the tempos of the whole working day and took precedence over any relational activity.

"...you are not always able to treat patients according to their needs, due to the number of tasks (many of them bureaucratic) that have to be carried out and that take time away from being able to have a deeper interaction with the patient." (03RD2P10).

Collegial nurse-physician relationships

This dimension showed very little change after the nurses' participatory process, remaining close to a neutral value. The item that decreased the most and was the worst rated by the nurses in this dimension was the perception of performing teamwork with the doctors (Table 2). In this sense, for the nurses it was important to carry out the first assessments together. They felt that it favoured the collection of information. They valued and assigned great importance to the cohesion between them, as this was the way to be able to carry out coherent and therapeutic actions.

"When new people come, I tell them: see if you can be in the first interview that the psychiatrist does... the attitude he has if you are there or not, sometimes you find that they say no, not with this nurse. But the first interview already gives you a lot of information and a relationship with the patient, it's like a more direct link, he knows that you are his nurse and that you are going to monitor him. etc."
(10FG4P01).

However, the nurses expressed difficulties due to a lack of communication and cohesion. For them, this lack of communication ended up affecting the nurse-patient relationship. In this sense, they stated that, at times, the medical teams made decisions without considering the nurses, generating an important loss of professional authority in the eyes of the people they cared for. For them, this loss of authority also occurred when certain members of the medical team undermined their authority in front of the patients or breached the established rules of the unit, putting their relationship with the users at risk. When these actions occurred, they felt a lack of support and professional recognition for the contributions and assessments they made as nurses.

“... You saw a person smoking twice, you took away his permission to go out and the psychiatrist comes by and says ‘no, no, he can go out’. Now, excuse me! If there are rules and the nurse has told him that he can't go out because those are the rules, how can you come up with this now? Are you discrediting us?”
(12FG4P02).

[TABLE 1. Differences on scores for items of the PES-NWI]

DISCUSSION

This study aimed to explore the effects on the perception of the practice environment among nurses in mental health units in the context of their participation in a process for improving the nurse-patient relationship. The quantitative results of the study showed a significant worsening of the nurses' perception of their practice environment.

Additionally, the qualitative findings of this study enabled us to identify the most relevant aspects of the environment that affected the nurses in order to establish a better nurse-patient relationship in clinical practice. The first fact that should be highlighted is the general worsening perceived by the nurses in relation to their environment. The nurses shifted from a positive assessment of their environment before starting the action research to a negative perception of the same. This finding can be explained by the nurses' greater awareness of the importance of the environment in order to establish a quality therapeutic relationship (Roviralta-Vilella et al., 2019). This may have led to nurses acquiring a more critical sense of the context of care through their participation in the action research process (Bush et al., 2018; Salzmann-Erikson, 2017).

More specifically, the dimension that suffered the greatest decrease was that of nurse manager leadership although it continued to be one of the dimensions most highly valued by nurses (Farmakas et al., 2014; Kurjenluoma et al., 2017; Redknap et al.,

2016; Roche et al., 2011; Van Bogaert, Clarke, Wouters, et al., 2013). The nurses pointed out that managers should motivate staff, and should have a presence on the units to understand and advocate for the needs of the team and the patients (Kanter, 1993; Wyder et al., 2017).

However, as in other studies, the lowest rated dimension by nurses was Staffing and resource adequacy (Gabrielsson et al., 2016; Hanrahan & Aiken, 2008; Hanrahan et al., 2010). Although this dimension did not suffer a significant decline in the context of the action research process, as this negative assessment already existed for the nurses, the process served to identify that the lack of resources and time put the quality of the relationship at risk (Kanter, 1993; Moreno-Poyato et al., 2016). In this sense, for nurses, fixed tasks took precedence over any relational activity (Busch, Moretti, Travaini, Wu, & Rimondini, 2019), primarily affecting the safety of both patients and staff (Vahidi et al., 2018).

Another notable finding was the worsening of the nurses' perception of their involvement in the affairs of the centre. While it was important for them to be involved in management, to have influence to be able to make changes and to feel heard and supported by management, the nurses stated that the management did not advocate for them or consider them to be able to raise issues or discuss the need for change (Roche et al., 2011; Wyder et al., 2017).

The perception of Nursing foundations for quality of care also decreased for the nurses. In this regard, they pointed out that there was no clear philosophical model advocating for the TR in their practice, such as patient allocation per referring nurse, and that this directly affected the fragmentation of care (Busch et al., 2019). In addition, they identified limitations in terms of training and orientation of new staff (Hooper, Browne, & O'Brien, 2016).

Finally, the doctor-nurse relationship did not undergo significant changes. The process allowed the nurses to confirm the importance of cohesion and teamwork between them, as this was the way to be able to carry out coherent and therapeutic actions with the patients (Gabrielsson et al., 2016; Wyder et al., 2017).

Strengths and limitations

This study has several strengths and limitations. Regarding the limitations, from a quantitative point of view, the single-group design does not allow the effect of the changes to be compared with a control group. Secondly, it is important to bear in mind the duration of the intervention. This could have influenced the rate of nurse dropout, since there were a significant number of service transfers during the study. Likewise, the changes in the work environment of each unit beyond those produced directly by the intervention should be considered. However, given the purpose of the study, the use of mixed methods enables the ability to complement and gain a deeper understanding of the phenomenon studied by integrating and inferring results. Thirdly, it should be considered that the source of data collection was only from the perspective of the nurses in the units. One of the strengths of the study was the representativeness obtained by the number of units and the number and characteristics of the nurses that participated, which enables these results to be generalized to the Catalan and Spanish context. Finally, regarding future lines of research, the findings suggest the need for more in-depth studies on the factors in the practice environment that influence the quality of the therapeutic relationship. Such studies should address the perspective of the directors and managers of the institutions, as well as research the relationship between the environment and indicators of quality of care and examine the perspective of the patients themselves.

CONCLUSIONS

Nurses consider that they are hardly involved in decision-making processes.

Furthermore, they require more presence and understanding from managers in order to identify real needs and to participate in decisions and thus improve the day-to-day clinical practice in the units. For them, the practice environment, more specifically, the lack of time and resources, affects their safety and that of their patients and conditions the therapeutic relationship by reducing the quality and quantity of interactions. Finally, nurses consider that the institutions should promote nursing foundations for quality of care, where, for example, priority is given to plans for welcoming the recruitment of new staff and ongoing training on the therapeutic relationship.

By using a mixed approach, this study contributes to a deeper understanding of how nurses in mental health units perceive their work environment and how it affects the improvement of the nurse-patient relationship in clinical practice. Thus, while the quantitative results provide information on the changes in the nurses' assessment of the environment, the qualitative data shed light on the most important aspects of the environment according to the nurses for the establishment of a quality therapeutic relationship.

REFERENCES

- Aiken, L. H., Sermeus, W., Van Den Heede, K., Sloane, D. M., Busse, R., McKee, M., ... Kutney-Lee, A. (2012). Patient safety, satisfaction, and quality of hospital care: Cross sectional surveys of nurses and patients in 12 countries in Europe and the United States. *BMJ (Online)*, *344*(7851). <https://doi.org/10.1136/bmj.e1717>
- Aiken, L. H., Sloane, D. M., Clarke, S., Poghosyan, L., Cho, E., You, L., ... Aunguroch, Y. (2011). Importance of work environments on hospital outcomes in nine countries. *International Journal for Quality in Health Care*, *23*(4), 357–364. <https://doi.org/10.1093/intqhc/mzr022>
- Al-Maaitah, R., AbuAlRub, R. F., & Al Blooshi, S. (2018). Practice environment as perceived by nurses in acute care hospitals in Sharjah and North Emirates. *Nursing Forum*, *53*(2), 213–222. <https://doi.org/10.1111/nuf.12245>
- Baum, F., MacDougall, C., & Smith, D. (2006). Participatory action research. *Journal of Epidemiology and Community Health*, *60*(10), 854–857. <https://doi.org/10.1136/jech.2004.028662>
- Bolg, J. R., Dwyer, P. A., Doherty, D. P., Pignataro, S. J., & Renaud, A. M. (2020). The Impact of Critical Reflective Inquiry Education on Experienced Nurses' Insights into Practice. *Journal for Nurses in Professional Development*, *36*(2), 68–73. <https://doi.org/10.1097/NND.0000000000000606>
- Busch, I. M., Moretti, F., Travaini, G., Wu, A. W., & Rimondini, M. (2019). Humanization of Care: Key Elements Identified by Patients, Caregivers, and Healthcare Providers. A Systematic Review. *The Patient*, *12*(5), 461–474. <https://doi.org/10.1007/s40271-019-00370-1>

- Bush, P. L., Pluye, P., Loignon, C., Granikov, V., Wright, M. T., Repchinsky, C., ... Macaulay, A. C. (2018). A systematic mixed studies review on Organizational Participatory Research: Towards operational guidance. *BMC Health Services Research, 18*(1), 1–15. <https://doi.org/10.1186/s12913-018-3775-5>
- Copanitsanou, P., Fotos, N., & Brokalaki, H. (2017). Effects of work environment on patient and nurse outcomes. *British Journal of Nursing, 26*(3), 172–176. <https://doi.org/10.12968/bjon.2017.26.3.172>
- Crowe, M., Inder, M., & Porter, R. (2015). Conducting qualitative research in mental health: Thematic and content analyses. *Australian & New Zealand Journal of Psychiatry, 49*(7), 616–623. <https://doi.org/10.1177/0004867415582053>
- de Pedro Gómez, J., Morales Asencio, J. M., Sesé Abad, A., Bennasar Veny, M., Artigues Vives, G., & Pericàs Beltran, J. (2009). Validación y adaptación al español de la escala del entorno de práctica enfermera del Nursing Work Index (Spanish validation and adaptation of the Nursing Work Index scale of nursing practice). *Metas de Enfermería, 12*(7), 65–75.
- Dorigan, G. H., & Guirardello, E. de B. (2017). Nursing practice environment, satisfaction and safety climate: the nurses' perception. *Acta Paulista de Enfermagem, 30*(2), 129–135. <https://doi.org/10.1590/1982-0194201700021>
- Ducharme, M. P., Bernhardt, J. M., Padula, C. A., & Adams, J. M. (2017). Leader Influence, the Professional Practice Environment, and Nurse Engagement in Essential Nursing Practice. *The Journal of Nursing Administration, 47*(7–8), 367–375. <https://doi.org/10.1097/NNA.0000000000000497>
- Erickson, J. I., Duffy, M. E., Ditomassi, M., & Jones, D. (2009). Psychometric evaluation of the Revised Professional Practice Environment (RPPE) scale. *The*

Journal of Nursing Administration, 39(5), 236–243.

<https://doi.org/10.1097/NNA.0b013e3181a23d14>

Farmakas, A., Papastavrou, E., Siskou, O., Karayiannis, G., & Theodorou, M. (2014).

Challenges in mental health nursing: Working in institutional or community settings? *Journal of Psychiatric and Mental Health Nursing*, 21(1), 39–45.

<https://doi.org/10.1111/jpm.12045>

Gabrielsson, S., Sävenstedt, S., & Olsson, M. (2016). Taking personal responsibility:

Nurses' and assistant nurses' experiences of good nursing practice in psychiatric inpatient care. *International Journal of Mental Health Nursing*, 25(5), 434–443.

<https://doi.org/10.1111/inm.12230>

Hanrahan, N. P., & Aiken, L. H. (2008). Psychiatric nurse reports on the quality of

psychiatric care in general hospitals. *Quality Management in Health Care*, 17(3), 210–217. <https://doi.org/10.1097/01.QMH.0000326725.55460.af>

Hanrahan, N. P., Aiken, L. H., McClaine, L., & Hanlon, A. L. (2010). Relationship

between Psychiatric Nurse Work Environments and Nurse Burnout in Acute Care General Hospitals. *Issues in Mental Health Nursing*, 31(3), 198–207.

<https://doi.org/10.3109/01612840903200068>

Hooper, M. E., Browne, G., & O'Brien, A. P. (2016). Graduate nurses' experiences of

mental health services in their first year of practice: An integrative review.

International Journal of Mental Health Nursing, 25(4), 286–298.

<https://doi.org/10.1111/inm.12192>

Huang, X., Wang, L., Dong, X., Li, B., & Wan, Q. (2021). Effects of nursing work

environment on work-related outcomes among psychiatric nurses: A mediating model. *Journal of Psychiatric and Mental Health Nursing*, 28(2), 186–196.

<https://doi.org/10.1111/jpm.12665>

Kanter, R. M. (1993). *Men and women of the corporation*. New York: Basic Books.

Kemmis, S., & Mctaggart, R. (2008). Participatory Action Research. In Y. S. L. Norman K. Denzin (Ed.), *Strategies of Qualitative Inquiry* (3rd ed., pp. 271–330). London.

Kurjenluoma, K., Rantanen, A., McCormack, B., Slater, P., Hahtela, N., & Suominen, T. (2017). Workplace culture in psychiatric nursing described by nurses. *Scandinavian Journal of Caring Sciences*, *31*(4), 1048–1058.
<https://doi.org/10.1111/scs.12430>

Lake, E. T. (2002). Development of the practice environment scale of the Nursing Work Index. *Research in Nursing & Health*, *25*(3), 176–188.
<https://doi.org/10.1002/nur.10032>

Lake, E. T., & Friese, C. R. (2006). Variations in Nursing Practice Environments. *Nursing Research*, *55*(1), 1–9. <https://doi.org/10.1097/00006199-200601000-00001>

Lin, C.-F., Lai, F.-C., Huang, W.-R., Huang, C.-I., & Hsieh, C.-J. (2020). Satisfaction With the Quality Nursing Work Environment Among Psychiatric Nurses Working in Acute Care General Hospitals. *Journal of Nursing Research*, *28*(2), e76.
<https://doi.org/10.1097/jnr.0000000000000350>

McAllister, S., Robert, G., Tsianakas, V., & McCrae, N. (2019). Conceptualising nurse-patient therapeutic engagement on acute mental health wards: An integrative review. *International Journal of Nursing Studies*, *93*, 106–118.
<https://doi.org/10.1016/j.ijnurstu.2019.02.013>

- Molin, J., Graneheim, U. H., Ringnér, A., & Lindgren, B.-M. (2019). Patients' experiences of taking part in Time Together – A nursing intervention in psychiatric inpatient care. *International Journal of Mental Health Nursing*, 28(2), 551–559. <https://doi.org/10.1111/inm.12560>
- Moreno-Poyato, A. R., Montesó-Curto, P., Delgado-Hito, P., Suárez-Pérez, R., Aceña-Domínguez, R., Carreras-Salvador, R., ... Roldán-Merino, J. F. (2016). The Therapeutic Relationship in Inpatient Psychiatric Care: A Narrative Review of the Perspective of Nurses and Patients. *Archives of Psychiatric Nursing*, 30(6), 782–787. <https://doi.org/10.1016/j.apnu.2016.03.001>
- Moreno-Poyato, A. R., Roviralta-Vilella, M., Abidi, K. El, Bordera-Antequera, S., Triviño-Briones, S. R., Olives, C. U., ... Puig-Llobet, M. (2021). Nursing Practice Environment in Psychiatric Units: A Scoping Review. *Journal of Psychosocial Nursing and Mental Health Services*, 1–8. <https://doi.org/10.3928/02793695-20210526-01>
- Moreno-Poyato, A. R., El Abidi, K., Rodríguez-Nogueira, Ó., Lluch-Canut, T., & Puig-Llobet, M. (2021). A qualitative study exploring the patients' perspective from the 'Reserved Therapeutic Space' nursing intervention in acute mental health units. *International Journal of Mental Health Nursing*, 30(3), 783–797. <https://doi.org/10.1111/inm.12848>
- Nantsupawat, A., Kunaviktikul, W., Nantsupawat, R., Wichaikhum, O. A., Thienthong, H., & Poghosyan, L. (2017). Effects of nurse work environment on job dissatisfaction, burnout, intention to leave. *International Nursing Review*, 64(1), 91–98. <https://doi.org/10.1111/inr.12342>
- Numminen, O., Ruoppa, E., Leino-Kilpi, H., Isoaho, H., Hupli, M., & Meretoja, R.

- (2016). Practice environment and its association with professional competence and work-related factors: perception of newly graduated nurses. *Journal of Nursing Management*, 24(1), E1–E11. <https://doi.org/10.1111/jonm.12280>
- Oshodi, T. O., Crockett, R., Bruneau, B., & West, E. (2017). The nursing work environment and quality of care: A cross-sectional study using the Essentials of Magnetism II Scale in England. *Journal of Clinical Nursing*, 26(17–18), 2721–2734. <https://doi.org/10.1111/jocn.13783>
- Poghosyan, L., Norful, A. A., Liu, J., & Friedberg, M. W. (2018). Nurse Practitioner Practice Environments in Primary Care and Quality of Care for Chronic Diseases. *Medical Care*, 56(9), 791–797. <https://doi.org/10.1097/MLR.0000000000000961>
- Redknap, R., Twigg, D., & Towell, A. (2016). What interventions can improve the mental health nursing practice environment? *International Journal of Mental Health Nursing*, 25(1), 42–50. <https://doi.org/10.1111/inm.12187>
- Roche, M. A., Duffield, C., Friedman, S., Twigg, D., Dimitrelis, S., & Rowbotham, S. (2016). Changes to nurses' practice environment over time. *Journal of Nursing Management*, 24(5), 666–675. <https://doi.org/10.1111/jonm.12371>
- Roche, M. A., & Duffield, C. M. (2010). A comparison of the nursing practice environment in mental health and medical-surgical settings. *Journal of Nursing Scholarship*, 42(2), 195–206. <https://doi.org/10.1111/j.1547-5069.2010.01348.x>
- Roche, M., Duffield, C., & White, E. (2011). Factors in the practice environment of nurses working in inpatient mental health: A partial least squares path modeling approach. *International Journal of Nursing Studies*, 48(12), 1475–1486. <https://doi.org/10.1016/j.ijnurstu.2011.07.001>

- Roviralta-Vilella, M., Moreno-Poyato, A. R., Rodríguez-Nogueira, Ó., Duran-Jordà, X., & Roldán-Merino, J. F. (2019). Relationship between the nursing practice environment and the therapeutic relationship in acute mental health units: A cross-sectional study. *International Journal of Mental Health Nursing*, 28(6).
<https://doi.org/10.1111/inm.12648>
- Salzmann-Erikson, M. (2017). Using Participatory Action Research to Develop a Working Model That Enhances Psychiatric Nurses' Professionalism: The Architecture of Stability. *Administration and Policy in Mental Health and Mental Health Services Research*, 44(6), 888–903. <https://doi.org/10.1007/s10488-017-0806-1>
- Tashakkori, A., & Teddlie, C. (2008). Quality of Inferences in Mixed Methods Research: Calling for an Integrative Framework. In M. M. Bergman (Ed.), *Advances in Mixed Methods Research* (pp. 101–119).
<https://doi.org/10.4135/9780857024329.d10>
- Teddlie, C., & Tashakkori, A. (2012). Common “Core” Characteristics of Mixed Methods Research: A Review of Critical Issues and Call for Greater Convergence. *American Behavioral Scientist*, 56(6), 774–788.
<https://doi.org/10.1177/0002764211433795>
- Vahidi, M., Ebrahimi, H., Areshtanab, H. N., Jafarabadi, M. A., Lees, D., Foong, A., & Cleary, M. (2018). Therapeutic Relationships and Safety of Care in Iranian Psychiatric Inpatient Units. *Issues in Mental Health Nursing*, 39(11), 967–976.
<https://doi.org/10.1080/01612840.2018.1485795>
- Van Bogaert, P., Clarke, S., Wouters, K., Franck, E., Willems, R., & Mondelaers, M. (2013). Impacts of unit-level nurse practice environment, workload and burnout on

nurse-reported outcomes in psychiatric hospitals: A multilevel modelling approach.

International Journal of Nursing Studies, 50(3), 357–365.

<https://doi.org/10.1016/j.ijnurstu.2012.05.006>

Wyder, M., Ehrlich, C., Crompton, D., McArthur, L., Delaforce, C., Dziopa, F., ...

Powell, E. (2017). Nurses experiences of delivering care in acute inpatient mental

health settings: A narrative synthesis of the literature. *International Journal of*

Mental Health Nursing, 26(6), 527–540. <https://doi.org/10.1111/inm.12315>

Zugai, J. S., Stein-Parbury, J., & Roche, M. (2015). Therapeutic Alliance in Mental

Health Nursing: An Evolutionary Concept Analysis. *Issues in Mental Health*

Nursing, 36(4), 249–257. <https://doi.org/10.3109/01612840.2014.969795>

Mental health nurses' perceptions on the practice environment

TABLE 1. Differences in scores for items on the PES-NWI

Variable	Baseline mean (SD)	Follow up mean (SD)	Difference mean (SD)	95% CI		P-value†
				Lower	Upper	
Nurse participation in hospital affairs	2.5 (0.6)	2.3 (0.5)	-0,2 (0.5)	-.262	-.085	.004
1. Staff nurses are formally represented in the management of the Centre (boards, decision-making bodies).	2.3 (0.9)	2.1 (0.9)	-0.2 (0.9)	-.452	-.053	.014
2. Nurses at the Centre enjoy opportunities to participate in decisions that affect policies implemented at the Centre.	2.2 (0.9)	2.0 (0.8)	-0.2 (0.8)	-.407	-.077	.004
3. Nursing staff are offered many opportunities for developing their professional training.	2.4 (0.9)	2.3 (0.9)	-0.1 (0.9)	-.296	.065	.206
4. The Management listens and offers answers to issues raised by nurses.	2.4 (0.9)	2.1 (0.8)	-0.3 (0.9)	-.462	-.107	.002
5. The Head of hit nursing department is accessible and readily “visible”.	2.7 (1.0)	2.5 (0.9)	-0.2 (0.9)	-.364	.007	.059
6. Room for professional development; opportunities for promotion.	2.7 (0.9)	2.5 (0.8)	-0.2 (0.9)	-.369	.032	.099
7. Nurse managers consult with nurses over problems arising in the day to day working routine.	2.6 (0.9)	2.4 (0.9)	-0.2 (0.9)	-.376	.018	.075
8. Staff nurses enjoy opportunities to participate in committees at the Centre, such as the Research Commission, Ethics board, etc.	2.7 (0.8)	2.7 (0.8)	0.01 (0.9)	-.176	.197	.911
9. Nurse managers have equal levels of authority and power as other management staff at the Centre.	2.3 (0.8)	2.3 (0.7)	0.01 (0.9)	-.186	.186	1.000
Nursing foundations for quality of care	2.7 (0.6)	2.6 (0.5)	-0,1 (0.5)	-.224	-.028	.012
10. Nurses’ diagnosis are taken into account.	2.5 (1.1)	2.5 (0.9)	0.01 (0.9)	-.197	.197	1.000

Mental health nurses' perceptions on the practice environment

11. An active quality assurance and enhancement programme is in place.	2.8 (0.9)	2.7 (0.7)	-0.05 (0.9)	-.237	.131	.572
12. A newcomers; welcome briefing and sponsorship programme is in place for nurses.	2.4 (1.0)	2.4 (1.0)	0.02 (0.9)	-.178	.220	.834
13. Care provided by nurses is based on a nursing model rather than on a biomedical model.	3.0 (0.8)	2.8 (0.8)	-0.1 (0.9)	-.340	.045	.132
14. Patient assignment promotes care continuity (e.g. the same nurse cares for the patient along the time).	2.7 (1.1)	2.4 (1.0)	-0.3 (1.2)	-.572	-.080	.010
15. A common, well-defined nursing philosophy is in place, which impregnates the patient care environment.	2.7 (0.8)	2.5 (0.6)	-0.2 (0.8)	-.358	-.021	.028
16. A written nursing care plan is drafted and updated for each patient.	2.6 (1.0)	2.5 (1.0)	-0.1 (0.9)	-.322	.069	.202
17. Managers at the Centre ensure that nurses provide high quality care.	2.7 (0.9)	2.5 (0.9)	-0.2 (0.9)	-.443	-.041	.019
18. Continuing training programmes for nurses are offered.	2.9 (0.9)	2.8 (0.9)	-0.1 (0.9)	-.255	.108	.422
19. Nurses employed at the Centre possess adequate clinical competencies.	3.1 (0.8)	2.9 (0.7)	-0.2 (0.7)	-.279	.026	.103
Nurse manager ability, leadership and support of nurses	3.1 (0.7)	2.8 (0.7)	-0.3 (0.8)	-.517	-.177	<.001
20. The Head Nurse is a good manager and leader.	3.2 (0.9)	2.9 (0.9)	-0.3 (0.9)	-.472	-.076	.007
21. The Head Nurse makes decisions that will support the staff, even in the case of conflict with the Medical staff.	3.3 (0.9)	2.8 (0.9)	-0.5 (1.0)	-.654	-.230	<.001
22. The Head Nurse sees errors as opportunities to extract lessons learned and to improve, not to criticize.	3.2 (0.8)	2.8 (0.9)	-0.4 (0.9)	-.563	-.173	<.001
23. The Head Nurse is understanding, able to advise and gives support to nurses.	3.3 (0.8)	2.9 (0.9)	-0.4 (0.9)	-.548	-.168	<.001

Mental health nurses' perceptions on the practice environment

24. Due recognition and praise is given for work well done.	2.8 (0.9)	2.5 (0.8)	-0.3 (1.1)	-.517	-.073	.010
Staffing and resource adequacy	2.1 (0.7)	2.0 (0.7)	-0.1 (0.6)	-.255	-.008	.037
25. Personnel numbers are sufficient to perform the workload adequately.	2.0 (0.9)	1.9 (0.9)	-0.1 (0.7)	-.310	-.005	.042
26. The number of qualified nurses is sufficient to provide quality care.	2.1 (1.0)	2.0 (0.9)	-0.1 (0.9)	-.285	.074	.247
27. Supporting services (hospital attendants, administrative staff, etc.) are adequate and allow nurses to spend more time with patients.	2.3 (0.9)	2.2 (0.9)	-0.1 (0.9)	-.258	.131	.520
28. There are sufficient opportunities and time to discuss care related problems with nurse colleagues.	2.2 (0.8)	2.0 (0.8)	-0.2 (0.8)	-.372	-.028	.023
Collegial nurse-physician relationships	2.5 (0.7)	2.4 (0.6)	-0.1 (0.6)	-.238	.027	.117
29. Teamwork regularly involves both Medical and nursing staff.	2.3 (0.9)	2.1 (0.8)	-0.2 (0.9)	-.346	.009	.063
30. Good working relations between doctors and nursing staff are in place.	2.7 (0.7)	2.6 (0.7)	-0.1 (0.8)	-.290	.037	.128
31. Practice among nurses and doctors is based on the appropriate principles of collaboration.	2.5 (0.7)	2.5 (0.7)	-0.02 (0.7)	-.167	.124	.775
PES-NWI total	2.6 (0.5)	2.4 (0.4)	-0.2 (0.4)	-.262	-.085	<.001

The values are expressed as mean (SD). In bold, statistically significant differences between baseline and follow-up. † Paired Student's t-test; PES-NWI, Practice Environment Scale of the Nursing Work Index

Figure 1. Mixed methods design

Figure 2. Differences in scores of the Practice Environment Scale of the Nursing Work Index