

## Hydrogen-Bonded Matched Ion Pair Gold(I) Catalysis

Àlex Martí, Gala Ogalla, and Antonio M. Echavarren\*

Cite This: *ACS Catal.* 2023, 13, 10217–10223

Read Online


ACCESS |

Metrics &amp; More

Article Recommendations

Supporting Information

**ABSTRACT:** The enantioselective reaction of 1,6-enynes with *O*-, *N*-, and *C*-nucleophiles has been developed by matched ion pair gold(I) catalysis in which the chiral gold(I) cation and anion are H-bonded through a urea group. Very high levels of enantiocontrol are achieved (up to >99:1 er) for a broad scope of substrates. DFT studies demonstrate the importance of the H-bond donor group in anchoring the matched chiral cation- and anion-favoring additional noncovalent interactions.




**KEYWORDS:** hydrogen bond interaction, chiral counterion, gold(I) catalysis, urea, phosphoramidate, match–mismatch

Although the field of gold(I) catalysis has experienced an exponential growth in past decades,<sup>1–9</sup> the development of enantioselective transformations with broader scope has been more difficult.<sup>10–13</sup> After the pioneering work by Toste et al. on gold asymmetric counterion-directed catalysis (ACDC) for the enantioselective cyclization of allenes using chiral phosphate salts in combination with achiral digold complexes (Figure 1),<sup>14–17</sup> the use of that concept has been used for the activation of alkyne<sup>18–22</sup> or allene-containing substrates<sup>23–27</sup> to circumvent some of the limitations in enantioselective gold(I) catalysis.<sup>28–30</sup>

Our group recently introduced H-bonded counterion-directed catalysis (HCDC) (Figure 1) using achiral gold catalysts containing urea or squaramide H-bond donor motifs.<sup>31,32</sup> In this approach, the H-bond donor facilitates the ligand-substrate exchange step and fixes the chiral information close to the reaction center, thereby allowing for an efficient transfer of the stereochemical information in cyclization reactions.<sup>33–36</sup>

Herein, we present the application of the HCDC approach with chiral binol-based phosphite gold(I) complexes<sup>37</sup> equipped with urea groups together with matched chiral counterions for the enantioselective gold(I)-catalyzed nucleophilic addition to 1,6-enynes, which takes place with excellent enantioselectivities.

A library of phosphite gold(I) complexes carrying a urea was easily prepared in a one-pot, two-step procedure starting from chiral (*R*)-binaphthols with the desired 3,3'-substitution pattern or from achiral resorcinol [4]arenes (Figure 2). Additionally, chiral silver salts were prepared from commercially available chiral BINOLs and related biphenols (Figure 2). Urea groups in the *para*-, *meta*-, or *ortho*-position with




**Figure 1.** Gold(I) in the ACDC and HCDC strategies.

Received: June 8, 2023

Revised: July 7, 2023

Published: July 20, 2023






**Figure 2.** Library of chiral gold(I) complexes and Ag(I) salts. <sup>3</sup>CYLview representations are for X-ray crystallography structures of **Au4** and **Au11**. Solvent molecules are omitted for clarity. **Au4** forms dimers in the solid state. Color code: P, orange; Au, yellow; Cl, green; O, red; N, blue; C, gray; and H, white. anthr = anthracenyl, phenant = phenantraceny.

respect to the phosphite were introduced to test the directing effect of the H-bond donor. We envisioned that chiral phosphoramidate silver salts would form more reactive catalysts since they are less basic than their corresponding phosphoric acid counterparts and, therefore, more easily substituted by the unsaturated substrates from the gold(I) coordination sphere.<sup>31</sup>


Our system proved to be particularly efficient in the addition of nucleophiles to 1,6-enynes.<sup>38–52</sup> We first examined, using high-throughput experimentation (HTE), the addition of indole to **1a**<sup>40,41</sup> with different chiral gold(I) complexes and silver(I) salts.<sup>53</sup> The best combinations were then scaled up to 0.05 mmol (Table 1). Gold(I) complexes with the urea in the *para*-position showed much higher reactivity than those with ureas at *ortho* or *meta*. Substituents in the 3,3'-position of the BINOL scaffolds in both the Au(I) catalyst and Ag salt also had an important impact. Through the use of toluene as solvent, a clear match-mismatch scenario was observed using (R)-**Au4** with the (R)- or (S)-enantiomers of **Ag6**, which led to **1a** with 95:5 *er* and 61.5:38.5 *er*, respectively (Table 1, entries 4 and 7). This observation was also present in other examples. For example, EtOH was used as a nucleophile with (R)-**Au1** or (R)-**Au4** in combination with the (R)- or (S)-enantiomers of the same Ag salt (Table 1, entries 8–11). Better yields and enantioselectivities were obtained in 1,4-dioxane. Thus, the combination of (R)-**Au4** with the (R)-enantiomers of **Ag1**, **Ag6**, or **Ag8** provided **1a** in >99:1, 97.5:2.5, and 95:5 *er*, respectively (Table 1, entries 13–15). Control experiments showed that neither (R)-**Au4** nor (R)-**Ag1** was active on its own (Table 1, entries 17 and 18). Whereas cavitand **Au9**,<sup>52</sup> equipped with *para*-urea groups, gives satisfactory results with (R)-**Ag6** (Table 1, entry 16), achiral Au(I) complex **Au12**, which was found to be the optimal for the intramolecular formal [4 + 2] cycloaddition of arylalkynes with alkenes,<sup>31</sup> together with (R)-**Ag1**, showed

**Table 1.** Enantioselective Gold(I)-Catalyzed Addition of Indole to 1,6-Enyne **1a**<sup>a</sup>



entry	[Au]	[Ag]	NuH	time (h)	yield (%) <sup>b</sup>	<i>er</i> <sup>c</sup>
1	<b>Au1</b>	(R)- <b>Ag6</b>	Ind <sup>d</sup>	44	61	78:22
2	<b>Au2</b>	(R)- <b>Ag6</b>	Ind <sup>d</sup>	44	69	54:46
3	<b>Au3</b>	(R)- <b>Ag6</b>	Ind <sup>d</sup>	44	6	50:50
4	<b>Au4</b>	(R)- <b>Ag6</b>	Ind <sup>d</sup>	44	82	95:5
5	<b>Au5</b>	(R)- <b>Ag6</b>	Ind <sup>d</sup>	44	74	60:40
6	<b>Au6</b>	(R)- <b>Ag6</b>	Ind <sup>d</sup>	44	39	54:46
7	<b>Au4</b>	(S)- <b>Ag6</b>	Ind <sup>d</sup>	44	61	61.5:38.5
8	<b>Au4</b>	(R)- <b>Ag6</b>	EtOH <sup>d</sup>	14	87	98:2
9	<b>Au4</b>	(S)- <b>Ag6</b>	EtOH <sup>d</sup>	14	85	57:43
10	<b>Au1</b>	(R)- <b>Ag6</b>	EtOH <sup>d</sup>	14	67	91:9
11	<b>Au1</b>	(S)- <b>Ag6</b>	EtOH <sup>d</sup>	14	53	47:53
12	<b>Au1</b>	(R)- <b>Ag1</b>	Ind <sup>e</sup>	14	82	91:9
13	<b>Au4</b>	(R)- <b>Ag1</b>	Ind <sup>e</sup>	14	93	>99:1
14	<b>Au4</b>	(R)- <b>Ag6</b>	Ind <sup>e</sup>	14	98	97.5:2.5
15	<b>Au4</b>	(R)- <b>Ag8</b>	Ind <sup>e</sup>	14	90	95:5
16	<b>Au9</b>	(R)- <b>Ag6</b>	Ind <sup>e</sup>	18	78	94:6
17	<b>Au4</b>		Ind <sup>e</sup>	24	0	
18		(R)- <b>Ag1</b>	Ind <sup>e</sup>	24	0	
19	<b>Au12</b>	(R)- <b>Ag1</b>	Ind <sup>e</sup>	24	18	52:48
20	<b>Au4</b>	AgSbF <sub>6</sub>	Ind <sup>e</sup>	24	76	92.5:7.5

<sup>a</sup>Reactions carried out under Ar or N<sub>2</sub> at a 0.05 mmol scale, at 27 °C with (R)-configured Au(I) complexes. <sup>b</sup>Yields determined by <sup>1</sup>H NMR using dodecane as internal standard. <sup>c</sup>*er* determined by supercritical fluid chromatography (SFC) using a chiral stationary phase. <sup>d</sup>Toluene (0.1 M) was used as solvent. <sup>e</sup>1,4-dioxane (0.1 M) was used as solvent. NuH = nucleophile. Ind = Indole.

Scheme 1. Enantioselective Addition of *O*-, *N*-, and *C*-Nucleophiles to 1,6-Enynes<sup>a</sup>

<sup>a</sup>Reaction performed under Ar or N<sub>2</sub> in anhydrous solvent (0.1 M). Yields given for isolated material after purification; *er* values were determined by HPLC or SFC on chiral stationary phase. Products were obtained as single diastereomers. Reaction times are in parentheses. <sup>b</sup>1,3-Cyclohexandione (2.0 equiv) was used as nucleophile. <sup>c</sup>Reaction carried out using (R)-Au7 and (R)-Ag5 at a 3 mol% catalyst loading. <sup>d</sup>Diphenylsulfoxide (1.5 equiv) was used as nucleophile.

poor reactivity and low enantioinduction in the formation of **2a** (Table 1, entry 19). Silver salt AgSbF<sub>6</sub> in combination with (R)-Au4 (Table 1, entry 20) gave lower yields and enantioselectivities than those observed with the dual-matched chiral system.<sup>53</sup>

It is interesting that, whereas in our system the matched pair is achieved with (R)-Au cation and (R)-phosphoramidate anion, in systems based on two BINOL-based cation/anion ion pairs, the matched catalyst resulted from the (R)/(S) ion combination.<sup>17,24,26</sup>

We examined the scope of the reaction using the optimal combination (R)-Au4 and (R)-Ag1 (Scheme 1). The reactions of 1,6-enyne **1a** were performed at 2 mol% catalyst loading in a 0.100 mmol scale, and two conditions were used depending on the nature of the nucleophile and the reactivity of the enyne: conditions A using 1,4-dioxane at 27 °C or conditions B using toluene at 27, 0, or -10 °C. Substituents in the 1, 2, and 5-

positions of indole were well tolerated to give adducts **2b–e**. Electron-rich arenes, such as 1,3,5-trimethoxybenzene, 1,3-dimethoxybenzene, and *N,N*-dimethylaniline, led to **2f–h** in excellent yields and enantioselectivities. Similarly, excellent results were also obtained with heteroatom-centered nucleophiles, such as anilines; a carbamate; alcohols; and water to give **2i–r** in excellent yields and enantioselectivities.


1,6-Enynes with different substitutions at the alkene also gave the corresponding adducts **2t–ad** in good to excellent yields and high enantioselectivities, except for those with a *para*- or *ortho*-anisyl group, which gave **2w** and **2z** in 87:13 and 93:7 *er* (Scheme 1). The absolute configurations of **2v** and **2ai** were determined to be the (*S,S*) and (*R,S*) by X-ray diffraction.<sup>54</sup> Changing the malonate to a dimethyl ether tether favored the formation of the cycloisomerization product, thereby leading to **2ae** in moderate yield. However, increasing the size of the ester from methyl to isopropyl reduced the

enantioselectivity to a 93:7 *er* in **2af**, which suggests that the tether plays an important role in the folding of the 1,6-enyne in the chiral pocket of the catalyst.

Changing the malonate tether in the 1,6-enyne to a *N*-tosyl led to a decrease in the enantioselectivity, which provided **2ag** in 78:22 *er*. (Scheme 1) However, using (*R*)-**Au7** together with (*R*)-**Ag5** gave adducts **2ag** and **2ah**, as well as aldehyde **2ai**, which resulted from the oxidation of the gold(I) carbene intermediate with diphenylsulfoxide,<sup>55</sup> in 98:2 to >99:1 *er*.

To further demonstrate the utility of this enantioselective addition of nucleophiles to 1,6-enynes, selected transformations into more complex products were performed (Scheme 2). Thus, the ring-closing metathesis of **2r** with the second

### Scheme 2. Product Derivatization




generation Grubbs catalyst<sup>56</sup> gave bicyclic derivative **3a** with no erosion on the enantioselectivity. The *ortho*-iodoaniline addition product **2k** underwent an intramolecular Heck reaction with the concomitant formation of a new stereocenter to afford 2,3,3a,9b-tetrahydro-1*H*-cyclopenta[*c*]quinoline **3b**.<sup>50</sup> Finally, the allylic position in pyrrolidine **2ah** was oxidized with  $\text{NaClO}_2$  and *N*-hydroxyphthalimide to give lactam **3c** with 98.5:1.5 *er*.

DFT studies were conducted [B3LYP-D3/6-31G(d) (C, H, P, O, F, N, S)//B3LYP-D3/6-311G(d,p) and SDD (Au), PCM = toluene]<sup>57</sup> to gain insight into the role of the urea and the possible secondary interactions involved in the enantioselective cyclization step using chiral (*R*)-**Au4** and chiral counterion from (*R*)-**Ag6**, which provided **2a** in 95:5 *er*.

Our computations predicted a Curtin–Hammett scenario (Scheme 3a), where the two orientations of the alkenes (**int1a** and **int3a**) are in equilibrium. Although **int3a** is 4.9 kcal/mol more stable than **int1a**, the major product arises from the latter via  $\text{TS}_{\text{int1a-2a}}$  which is lower in energy than  $\text{TS}_{\text{int3a-4a}}$ , thereby giving rise to product **2a** with an *S* configuration by reaction through the *Re* face of the alkene, which agrees with the experimental results.

Noncovalent interaction (NCI) plots were performed to visualize the noncovalent interactions in the two possible transition states (Scheme 3b,c) which revealed that the H-bond interactions from the urea and phosphoramidate group act as anchors of the two parts and favor the additional interactions that stabilize the transition state  $\text{TS}_{\text{int1a-2a}}$ . Apart

### Scheme 3. DFT Calculations for the Enantiodetermining Step<sup>a</sup>



<sup>a</sup>Two most relevant pathways for the enantiodetermining cyclization step of **1a** with (*R*)-**Au4** and (*R*)-**Ag6**. NCI plots and CYLview representations for  $\text{TS}_{\text{int1a-2a}}$  and  $\text{TS}_{\text{int3a-4a}}$ . Hydrogens are omitted for clarity, except relevant ones. Strong attractive interactions are blue, weak attractive interactions are green, and strong repulsive interactions are red. Color code: P, orange; Au, yellow; F, cyan; O, red; N, blue; S, purple; C, gray; and H, white. Energy values are in kcal/mol. HB = hydrogen bond.

from the strong H-bonding interactions, T-shaped- $\pi$  interactions between the *ortho*-C–H of the *N*-phenyl urea and the  $\pi$ -system of the BINOL counterion were observed. A strong extended attractive sandwich  $\pi$ - $\pi$  interaction between the  $\pi$ -system of the cinnamyl alkene of the substrate and one

anthracenyl group of the chiral counterion were observed in  $\text{TS}_{\text{int}1\text{a}-2\text{a}}$ . However, for  $\text{TS}_{\text{int}3\text{a}-4\text{a}}$ , a T-shaped  $\pi$ -attractive interaction was found between the C–H in *para* position to the alkene in the aryl ring of the substrate and the anthracene of the counterion, which helped to stabilize the transition state.

Finally, other reactions were tested by combining BINOL-derived gold(I) catalysts and different silver(I) salts as chloride scavengers, but the optimal ion pair combination could not be found. We discovered that 1,6-enynes bearing internal aryl-substituted alkynes led to formal products of [4 + 2] cycloaddition<sup>31</sup> with poor yields and moderate enantioselectivities, most likely because of the small pocket generated by the ion pair. However, when using chiral gold(I) complexes and small achiral counterions, such as  $\text{AgSbF}_6$ , promising enantioselectivities were found (95:5 *er*). We also tested the [2 + 2] cycloaddition of phenylacetylene with alkenes,<sup>58</sup> but no promising combination was observed.

In summary, we have developed the enantioselective nucleophilic addition of hetero- and carbonucleophiles that proceeds with the broadest scope and highest enantioselectivity using a chiral catalyst with a chirally matched gold(I) phosphitoureia and phosphoramidate, which can be readily prepared from commercially available BINOLs. A model for the enantioinduction has been proposed on the basis of DFT calculations and NCI plots where the urea in the chiral gold(I) cation anchors the chiral counterion in close proximity creating a chiral pocket to fold the unsaturated substrate.

## ■ ASSOCIATED CONTENT

### SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acscatal.3c02638>.

Optimization tables, procedures, characterization, NMR spectra, SFC and HPLC traces, DFT computations, and crystallographic data (PDF)

### Accession Codes

CCDC 2266405–2266408 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: + 44 1223 336033.

## ■ AUTHOR INFORMATION

### Corresponding Author

Antonio M. Echavarren – *Institute of Chemical Research of Catalonia (ICIQ), Barcelona Institute of Science and Technology (BIST), 43007 Tarragona, Spain; Departament de Química Orgànica i Analítica, Universitat Rovira i Virgili (URV), 43007 Tarragona, Spain; [orcid.org/0000-0001-6808-3007](https://orcid.org/0000-0001-6808-3007); Email: [aechavarren@iciq.es](mailto:aechavarren@iciq.es)*

### Authors

Àlex Martí – *Institute of Chemical Research of Catalonia (ICIQ), Barcelona Institute of Science and Technology (BIST), 43007 Tarragona, Spain; Departament de Química Orgànica i Analítica, Universitat Rovira i Virgili (URV), 43007 Tarragona, Spain*

Gala Ogalla – *Institute of Chemical Research of Catalonia (ICIQ), Barcelona Institute of Science and Technology (BIST), 43007 Tarragona, Spain; Departament de Química*

*Orgànica i Analítica, Universitat Rovira i Virgili (URV), 43007 Tarragona, Spain*

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acscatal.3c02638>

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We thank the MCIN/AEI/10.13039/501100011033 (PID2019-104815GB-I00 and CEX2019-000925-S), the MUniv (FPU20/0255 Fellowship to À.M.), the European Research Council (Advanced Grant 835080), the AGAUR (2021 SGR 01256), and CERCA Program/Generalitat de Catalunya for financial support. We also thank Dr. Imma Escofet for assistance with the computational calculations, as well as the ICIQ X-ray diffraction, NMR, and the chromatography and mass spectrometry units.

## ■ REFERENCES

- (1) Hashmi, A. S. K. Homogeneous gold catalysts and alkynes: A successful liaison. *Gold Bull.* **2003**, *36*, 3–9.
- (2) Zhang, L.; Sun, J.; Kozmin, S. A. Gold and platinum catalysis of enyne cycloisomerization. *Adv. Synth. Catal.* **2006**, *348*, 2271–2296.
- (3) Fürstner, A.; Davies, P. W. Catalytic Carbophilic Activation: Catalysis by Platinum and Gold  $\pi$  Acids. *Angew. Chem., Int. Ed.* **2007**, *46*, 3410–3449.
- (4) Fensterbank, L.; Malacria, M. Molecular complexity from polyunsaturated substrates: the gold catalysis approach. *Acc. Chem. Res.* **2014**, *47*, 953–965.
- (5) Dorel, R.; Echavarren, A. M. Gold (I)-catalyzed activation of alkynes for the construction of molecular complexity. *Chem. Rev.* **2015**, *115*, 9028–9072.
- (6) Asiri, A. M.; Hashmi, A. S. K. Gold-catalyzed reactions of diynes. *Chem. Soc. Rev.* **2016**, *45*, 4471–4503.
- (7) Boyle, J. W.; Zhao, Y.; Chan, P. W. H. Product Divergence in Coinage-Metal-Catalyzed Reactions of  $\pi$ -Rich Compounds. *Synth.* **2018**, *50*, 1402–1416.
- (8) Mato, M.; Franchino, A.; García-Morales, C.; Echavarren, A. M. Gold-catalyzed synthesis of small rings. *Chem. Rev.* **2021**, *121*, 8613–8684.
- (9) Campeau, D.; Rayo, D. F. L.; Mansour, A.; Muratov, K.; Gagosz, F. Gold-catalyzed reactions of specially activated alkynes, allenes, and alkenes. *Chem. Rev.* **2021**, *121*, 8756–8867.
- (10) Milcendeau, P.; Sabat, N.; Ferry, A.; Guinchar, X. Gold-catalyzed enantioselective functionalization of indoles. *Org. Biomol. Chem.* **2020**, *18*, 6006–6017.
- (11) Jiang, J. J.; Wong, M. K. Recent advances in the development of chiral gold complexes for catalytic asymmetric catalysis. *Chem.—Asian J.* **2021**, *16*, 364–377.
- (12) Zuccarello, G.; Escofet, I.; Caniparoli, U.; Echavarren, A. M. New-Generation Ligand Design for the Gold-Catalyzed Asymmetric Activation of Alkynes. *ChemPlusChem.* **2021**, *86*, 1283–1296.
- (13) Escofet, I.; Zuccarello, G.; Echavarren, A. M. Gold-catalyzed enantioselective cyclizations and cycloadditions. *Adv. Organomet. Chem.* **2022**, *77*, 1–42.
- (14) Hamilton, G. L.; Kang, E. J.; Mba, M.; Toste, F. D. A powerful chiral counterion strategy for asymmetric transition metal catalysis. *Science* **2007**, *317*, 496–499.
- (15) LaLonde, R. L.; Wang, Z. L.; Mba, M.; Lackner, A. D.; Toste, F. D. Gold(I)-Catalyzed Enantioselective Synthesis of Pyrazolidines, Isoxazolidines, and Tetrahydrooxazines. *Angew. Chem., Int. Ed.* **2010**, *49*, 598–601.
- (16) Mourad, A. K.; Leutzow, J.; Czekelius, C. Anion-Induced Enantioselective Cyclization of Diynamides to Pyrrolidines Catalyzed

- by Cationic Gold Complexes. *Angew. Chem., Int. Ed.* **2012**, *51*, 11149–11152.
- (17) Miles, D. H.; Veguillas, M. F.; Toste, F. D. Gold(i)-catalyzed enantioselective bromocyclization reactions of allenes. *Chem. Sci.* **2013**, *4*, 3427–3431.
- (18) Spittler, M.; Lutsenko, K.; Czekelius, C. Total synthesis of (+)-mesembrine applying asymmetric gold catalysis. *J. Org. Chem.* **2016**, *81*, 6100–6105.
- (19) Zhang, Z.; Smal, V.; Retailleau, P.; Voituriez, A.; Frison, G.; Marinetti, A.; Guinchard, X. Tethered counterion-directed catalysis: merging the chiral ion-pairing and bifunctional ligand strategies in enantioselective gold(I) catalysis. *J. Am. Chem. Soc.* **2020**, *142*, 3797–3805.
- (20) Pedrazzani, R.; An, J.; Monari, M.; Bandini, M. New Chiral BINOL-Based Phosphates for Enantioselective [Au(I)]-Catalyzed Dearomatization of  $\beta$ -Naphthols with Allenamides. *Eur. J. Org. Chem.* **2021**, *2021*, 1732–1736.
- (21) Yu, Y.; Zhang, Z.; Voituriez, A.; Rabasso, N.; Frison, G.; Marinetti, A.; Guinchard, X. Enantioselective Au(I)-catalyzed dearomatization of 1-naphthols with allenamides through Tethered Counterion-Directed Catalysis. *Chem. Commun.* **2021**, *57*, 10779–10782.
- (22) Zhang, Z.; Sabat, N.; Frison, G.; Marinetti, A.; Guinchard, X. Enantioselective Au(I)-Catalyzed Multicomponent Annulations via Tethered Counterion-Directed Catalysis. *ACS Catal.* **2022**, *12*, 4046–4053.
- (23) Aikawa, K.; Kojima, M.; Mikami, K. Axial chirality control of gold (biphep) complexes by chiral anions: application to asymmetric catalysis. *Angew. Chem., Int. Ed.* **2009**, *48*, 6073–6077.
- (24) Aikawa, K.; Kojima, M.; Mikami, K. Synergistic Effect: Hydroalkoxylation of Allenes through Combination of Enantiopure BIPHEP-Gold Complexes and Chiral Anions. *Adv. Synth. Catal.* **2010**, *352*, 3131–3135. Aikawa, K.; Kojima, M.; Mikami, K. *Adv. Synth. Catal.* **2011**, *353*, 2882–2883.
- (25) Barreiro, E. M.; Brogini, D. F. D.; Adrio, L. A.; White, A. J. P.; Schwenk, R.; Togni, A.; Hii, K. K. Gold (I) complexes of conformationally constricted chiral ferrocenyl phosphines. *Organometallics* **2012**, *31*, 3745–3754.
- (26) Handa, S.; Lippincott, D. J.; Aue, D. H.; Lipshutz, B. H. Asymmetric gold-catalyzed lactonizations in water at room temperature. *Angew. Chem., Int. Ed.* **2014**, *53*, 10658–10662.
- (27) Zi, W.; Toste, F. D. Gold (I)-Catalyzed Enantioselective Desymmetrization of 1, 3-Diols through Intramolecular Hydroalkoxylation of Allenes. *Angew. Chem., Int. Ed.* **2015**, *54*, 14447–14451.
- (28) Loh, C.; Enders, D. Merging organocatalysis and gold catalysis—a critical evaluation of the underlying concepts. *Chem.—Eur. J.* **2012**, *18*, 10212–10225.
- (29) Inamdar, S.; Konala, A.; Patil, N. T. When gold meets chiral Brønsted acid catalysts: extending the boundaries of enantioselective gold catalysis. *Chem. Commun.* **2014**, *50*, 15124–15135.
- (30) Bao, M.; Zhou, S.; Hu, W.; Xu, X. Recent advances in gold-complex and chiral organocatalyst cooperative catalysis for asymmetric alkyne functionalization. *Chin. Chem. Lett.* **2022**, *33*, 4969–4979.
- (31) Franchino, A.; Martí, C.O.; Echavarren, A. M. H-Bonded Counterion-Directed Enantioselective Au(I) Catalysis. *J. Am. Chem. Soc.* **2022**, *144*, 3497–3509.
- (32) Martí, C.O.; Montesinos-Magraner, M.; Echavarren, A. M.; Franchino, A. H-Bonded Counterion-Directed Catalysis: Enantioselective Gold(I)-Catalyzed Addition to 2-Alkynyl Enones as a Case Study. *Eur. J. Org. Chem.* **2022**, *2022* (38), No. e202200518.
- (33) Sandee, A. J.; van der Burg, A. M.; Reek, J. N. H. UREAphos: Supramolecular Bidentate Ligands for Asymmetric Hydrogenation. *Chem. Commun.* **2007**, 864–866.
- (34) Meeuwissen, J.; Kuil, M.; van der Burg, A. M.; Sandee, A. J.; Reek, J. N. H. Application of a Supramolecular-Ligand Library for the Automated Search for Catalysts for the Asymmetric Hydrogenation of Industrially Relevant Substrates. *Chem.—Eur. J.* **2009**, *15*, 10272–10279.
- (35) Meeuwissen, J.; Reek, J. N. H. Supramolecular Catalysis beyond Enzyme Mimics. *Nat. Chem.* **2010**, *2*, 615–621.
- (36) Reek, J. N. H.; de Bruin, B.; Pullen, S.; Mooibroek, T. J.; Kluwer, A. M.; Caumes, X. Transition Metal Catalysis Controlled by Hydrogen Bonding in the Second Coordination Sphere. *Chem. Rev.* **2022**, *122*, 12308–12369.
- (37) Delpont, N.; Escofet, I.; Pérez-Galán, P.; Spiegl, D.; Raducan, M.; Bour, C.; Sinisi, R.; Echavarren, A. M. Modular chiral gold(i) phosphite complexes. *Catal. Sci. Technol.* **2013**, *3*, 3007–3012.
- (38) Nieto-Oberhuber, C.; Muñoz, M. P.; Buñuel, E.; Nevado, C.; Cárdenas, D. J.; Echavarren, A. M. Cationic Gold(I) Complexes: Highly Alkynophylic Catalysts for the Exo- and Endo-Cyclization of Enynes. *Angew. Chem., Int. Ed.* **2004**, *43*, 2402–2406.
- (39) Nieto-Oberhuber, C.; Muñoz, M. P.; López, S.; Jiménez-Núñez, E.; Nevado, C.; Herrero-Gómez, E.; Raducan, M.; Echavarren, A. M. Gold(I)-Catalyzed Cyclizations of 1,6-Enynes: Alkoxylation and Exo/Endo Skeletal Rearrangements. *Chem.—Eur. J.* **2006**, *12*, 1677–1693.
- (40) Toullec, P. Y.; Genin, E.; Leseurre, L.; Genet, J.-P.; Michelet, V. Room-Temperature Au-Catalyzed C-C Bond Formation through a Tandem Friedel-Crafts-Type Addition/Carbocyclization Reaction. *Angew. Chem., Int. Ed.* **2006**, *45*, 7427–7430.
- (41) Amijs, C. H. M.; Ferrer, C.; Echavarren, A. M. Gold(I)-Catalyzed Arylation of 1,6-Enynes: Different Site Reactivity of Cyclopropyl Gold Carbenes. *Chem. Commun.* **2007**, 698–700.
- (42) Amijs, C. H. M.; López-Carrillo, V.; Raducan, M.; Pérez-Galán, P.; Ferrer, C.; Echavarren, A. M. Gold(I)-Catalyzed Intermolecular Addition of Carbon Nucleophiles to 1,5- and 1,6-Enynes. *J. Org. Chem.* **2008**, *73*, 7721–7730.
- (43) Chao, C. M.; Genin, E.; Toullec, P. Y.; Genêt, J. P.; Michelet, V. Towards Asymmetric Au-Catalyzed Hydroxy- and Alkoxylation of 1,6-Enynes. *J. Organomet. Chem.* **2009**, *694*, 538–545.
- (44) Leseurre, L.; Chao, C. M.; Seki, T.; Genin, E.; Toullec, P. Y.; Genêt, J. P.; Michelet, V. Synthesis of Functionalized Carbo- and Heterocycles via Gold-Catalyzed Cycloisomerization Reactions of Enynes. *Tetrahedron* **2009**, *65*, 1911–1918.
- (45) Matsumoto, Y.; Selim, K. B.; Nakanishi, H.; Yamada, K.; Yamamoto, Y.; Tomioka, K. Chiral Carbene Approach to Gold-Catalyzed Asymmetric Cyclization of 1,6-Enynes. *Tetrahedron Lett.* **2010**, *51*, 404–406.
- (46) Pradal, A.; Chao, C. M.; Vitale, M. R.; Toullec, P. Y.; Michelet, V. Asymmetric Au-Catalyzed Domino Cyclization/nucleophile Addition Reactions of Enynes in the Presence of Water, Methanol and Electron-Rich Aromatic Derivatives. *Tetrahedron* **2011**, *67*, 4371–4377.
- (47) Wang, W.; Yang, J.; Wang, F.; Shi, M. Axially Chiral N-Heterocyclic Carbene gold(I) Complex Catalyzed Asymmetric Cycloisomerization of 1,6-Enynes. *Organometallics* **2011**, *30*, 3859–3869.
- (48) Yamada, K. I.; Matsumoto, Y.; Selim, K. B.; Yamamoto, Y.; Tomioka, K. Steric Tuning of C 2-Symmetric Chiral N-Heterocyclic Carbene in Gold-Catalyzed Asymmetric Cyclization of 1,6-Enynes. *Tetrahedron* **2012**, *68*, 4159–4165.
- (49) Gung, B. W.; Holmes, M. R.; Jones, C. A.; Ma, R.; Barnes, C. L. Structure-enantioselectivity Correlation in NHC-Au(I) Catalysis for 1,6-Enynecyclizations. *Tetrahedron Lett.* **2016**, *57*, 3912–3915.
- (50) Miller, R.; Carreras, J.; Muratore, M. E.; Gaydou, M.; Camponovo, F.; Echavarren, A. M. Broad Scope Aminocyclization of Enynes with Cationic JohnPhos-Gold(I) Complex as the Catalyst. *J. Org. Chem.* **2016**, *81*, 1839–1849.
- (51) Tugny, C.; Del Rio, N.; Koohgard, M.; Vanthuyne, N.; Lesage, D.; Bijouard, K.; Zhang, P.; Meijide Suárez, J.; Roland, S.; Derat, E.; Bistri-Aslanoff, O.; Sollogoub, M.; Fensterbank, L.; Mourès-Mansuy, V.  $\beta$ -Cyclodextrin-NHC-Gold(I) Complex ( $\beta$ -ICyD)AuCl: A Chiral Nanoreactor for Enantioselective and Substrate-Selective Alkoxylation Reactions. *ACS Catal.* **2020**, *10*, 5964–5972.

(52) Martín-Torres, I.; Ogalla, G.; Yang, J. M.; Rinaldi, A.; Echavarren, A. M. Enantioselective Alkoxy cyclization of 1,6-Enynes with Gold(I)-Cavitands: Total Synthesis of Mafaicheenamine C. *Angew. Chemie. Int. Ed.* **2021**, *60*, 9339–9344.

(53) See the [Supporting Information](#) for more details.

(54) (a) Comparison of the reported optical rotations of **2a**–**2d**<sup>43</sup> with those of our products on the basis of the absolute configurations of **2v** and **2ai** determined by X-ray diffraction shows that the originally assigned configuration in ref 43 should be reversed.<sup>53</sup> (b) Similarly, the absolute configuration of two of the products (**5a,b**) reported in ref 52 should be reversed.

(55) Witham, C. A.; Mauleon, P.; Shapiro, N. D.; Sherry, B. D.; Toste, F. D. Gold (I)-catalyzed oxidative rearrangements. *J. Am. Chem. Soc.* **2007**, *129*, 5838–5839.

(56) Chao, C. M.; Toullec, P. Y.; Michelet, V. Synthesis of polycyclic heterocycles via sequential Au-catalyzed cycloisomerization and Ru-catalyzed metathesis reactions. *Tetrahedron Lett.* **2009**, *50*, 3719–3722.

(57) See the [Supporting Information](#) for details. A data set collection of computational results is available in the ioChem-BD repository and can be accessed <https://iochem-bd.iciq.es/browse/handle/100/60425>. Álvarez-Moreno, M.; De Graaf, C.; Lopez, N.; Maseras, F.; Poblet, J. M.; Bo, C. Managing the Computational Chemistry Big Data Problem: The ioChem-BD Platform. *J. Chem. Inf. Model.* **2015**, *55*, 95–103.

(58) García-Morales, C.; Ranieri, B.; Escofet, I.; López-Suarez, L.; Obradors, C.; Konovalov, A. I.; Echavarren, A. M. Enantioselective Synthesis of Cyclobutenes by Intermolecular [2 + 2] Cycloaddition with Non-C2 Symmetric Digold Catalysts. *J. Am. Chem. Soc.* **2017**, *139*, 13628–13631.

## Recommended by ACS

### Synthesis of Hexamethyl-1,1'-spirobiindane-Based Chiral Spiro Cp Ligands and Their Application in Rhodium-Catalyzed Enantioselective Aryl C–H Addition to Nitroal...

Hui Yang, Shu-Li You, *et al.*

JUNE 20, 2023  
ACS CATALYSIS

READ 

### Diastereoselective Rotaxane Synthesis with Pillar[5]arenes via Co-crystallization and Solid-State Mechanochemical Processes

Keisuke Wada, Tomoki Ogoshi, *et al.*

JULY 06, 2023  
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

### Design and Synthesis of Planar-Chiral Oxazole–Pyridine *N,N*-Ligands: Application in Palladium-Catalyzed Asymmetric Acetoxylation Cyclization

Yu-Qing Bai, Yong-Gui Zhou, *et al.*

JULY 11, 2023  
ACS CATALYSIS

READ 

### Force-Modulated Selectivity of the Rhodium-Catalyzed Hydroformylation of 1-Alkenes

Yichen Yu, Ross A. Widenhoefer, *et al.*

OCTOBER 28, 2022  
ACS CATALYSIS

READ 

Get More Suggestions >